

Orthovisc®

Hyaluronic Acid Multi-Injection

Medical Information Brochure

Structured product education for osteoarthritis pain management, designed for the Platinum Pharma website brochure download section.



HA helps lubricate, cushion and mechanically support the joint.




3 weekly injections



Up to 6 months*



Ultra-pure High MW HA



Product overview: Orthovisc® is an ultra-pure, high molecular weight hyaluronic acid viscosupplement administered as a multi-injection treatment course to help relieve symptoms of osteoarthritis through lubrication and mechanical support of the joint. Claims and indication details should be verified against the Egypt-approved IFU and local registration documents before publication.

*Pain relief duration and clinical claims must be verified against the Egypt-approved IFU and local registration documents before publication.

Understanding knee osteoarthritis and viscosupplementation

A healthy knee contains synovial fluid, a viscous liquid that lubricates and absorbs shocks. In osteoarthritis, changes in synovial fluid and degeneration of joint structures may contribute to pain, swelling, reduced flexibility, grating sensation and reduced mobility.


1

Lubrication

HA supplementation is intended to improve synovial-fluid lubrication and help reduce friction within the joint.

2

Cushioning

The gel-like viscoelastic nature of HA supports cushioning during movement and weight-bearing activities.

3

Mechanical support

Orthovisc® is described as providing lubrication and mechanical support as a viscoelastic supplement or replacement.

What is Orthovisc®?

Orthovisc® is a sterile, non-pyrogenic, clear, viscoelastic solution of high molecular weight hyaluronan supplied in a single-use syringe. It is used as a hyaluronic acid viscosupplement for osteoarthritis-related joint pain, according to the applicable regional IFU.



Product facts and administration



Key product facts

- **Category:** hyaluronic acid multi-injection viscosupplement
- **Composition:** ultra-pure high molecular weight sodium hyaluronate
- **Presentation:** sterile viscoelastic solution in a single-use syringe
- **Strength:** 30 mg / 2 mL per syringe
- **Source:** non-animal-derived hyaluronic acid produced by bacterial fermentation
- **Intended use:** symptomatic osteoarthritis knee pain relief according to local approval



Administration summary

- One intra-articular injection into the selected knee joint at each visit
- Administered by a qualified healthcare professional using strict aseptic technique
- Remove joint effusion, if present, before injection
- Treatment is typically given as three weekly injections according to product labeling
- Product selection and suitability should follow the approved IFU and physician judgment
- Post-procedure monitoring and patient counseling should follow local practice

Treatment course visual



Three weekly intra-articular injections (one injection per week)



Post-injection patient advice


As with any invasive joint procedure, patients are advised to avoid strenuous activity or prolonged weight-bearing activity, such as running or tennis, within 48 hours after the intra-articular injection. Patients should follow their treating physician's instructions.

Evidence highlights and safety

Clinical evidence summary

Anika Orthovisc brochure materials describe less pain and improved mobility after a 3-injection treatment course.

Source materials also describe pain relief within 1 week of injection and pain relief lasting up to 6 months, while helping reduce reliance on prescription pain medication.



*Duration of effect is individual and subject to clinical response, local labeling, and medical approval.



Long-term relief

Brochure materials describe pain relief that may continue for up to 6 months, subject to local labeling and medical approval.



Improved mobility

Orthovisc® materials describe less pain and improved mobility after the treatment course.



Medication-use reduction

Source brochure materials describe reduced use of prescription pain medication after treatment.



Essential safety and suitability notes

Do not administer to patients with known hypersensitivity to hyaluronate preparations or gram positive bacterial proteins, or in knees with infections or skin disease at the injection site or joint. Transient pain or swelling may occur after intra-articular injection. Orthovisc® should be injected only by medical professionals trained in accepted intra-articular injection technique. Final publication should be checked against the Egypt-approved IFU and local registration file.



Core sources used: Anika Orthovisc brochure and Orthovisc IFU references; KneePainRelief.ca and Anika Orthovisc product information. **Regulatory note:** final publication should be checked against the Egypt-approved IFU/registration file and Platinum Pharma/Anika medical approval.