

Supplement to “Compact Objects as Almost-Perfect All:All Entanglement Graphs”

Methods, honesty ledger, N-scaling, and module map (v5.0)

Philippe Leblond
leblond.philippe@gmail.com

October 3, 2026

Contents

S1 Parameter audit and honesty ledger	1
S2 Gravity methods (AS, AT, AU, BB, BH, BI, BJ, BV)	2
S3 Quantum-information methods (A–O, Q–Y)	3
S4 Compact objects and 2PN (BU)	4
S5 UV nulls and phenomenology (T, U, V, BD–BF)	6
S6 Kill list (AN + BU)	7
S7 N-scaling and calibration record	8
S8 PPN ledger and walk/foam/healing notes	8
S9 Archival ledger and O5 trigger (BU detail)	8
S10 Module map and reproduction	9
S11 Emergent-dimension methods (L0)	10
S12 Relational field program (F-layer)	14

S1 Parameter audit and honesty ledger

Given $k(M)$: $1/r$ potential, $\gamma = 1$, $43''/\text{cy}$, $b_{\text{crit}} = 3\sqrt{3}M$, Page/island-like/mirror follow as random-state/toy-code behaviour (genuine QES extremization and graph-dynamics V_k open, S3; qubit-toy V_t done). Not derived: mass–radius scaling itself, $\beta(N)$ from geometry, w from the graph Laplacian (Damour–Schäfer from wiring), NICER M – R – Λ from routing stiffness, Kerr $M_2/g_{t\phi}$ /ISCO/QNM from the graph.

Derived in-repo (new in v4.x): p from exact Ollivier–Ricci (not hard-coded), $M(c_1, c_2)$ inversion, $M_{\text{ej}}(M_{\text{tot}})$ scaling, peak-time ordering, band mags, O5 yield arithmetic, c from $\ln 2$ scattering, UV

pop as graph disconnection, p at $N = 4000/8000/16000$ with turnover measured, d_{eff} bracket + convergence + shell/tense-plug pins (v0.6, S11).

Ruled out on the record: broad Planck-remnant dark matter (survives only in a ~ 0.4 -dex EMD window at $\sim 4 \times 10^5$ g); isotropic $b_{\text{crit}} = 8M$; naive $\gamma = 2$ strain; linear LIV; flat $p = 0.49$ (-10σ); bare shortest-path $d(i, j)$ for tense regions (inverted vs GR); resistance/communicability distances (S11); directed/handed motion of pure-D5 ∞ plateau blobs (Stage-0 NULL, S11).

S2 Gravity methods (AS, AT, AU, BB, BH, BI, BJ, BV)

AS entropic Newton. Leg screens + equipartition + Bekenstein bound give $F = M_1 M_2 / r^2$ exact (slope -2 to 10^{-9}); leapfrog orbits close, $T^2 \propto r^3$. Flux $\propto 1/r^2$ distinguished from force (temperature corrects). Fig. 43.

AT redshift. GPS $+5.29 \times 10^{-10}$ and Pound–Rebka 2.55×10^{-15} derived with no new parameters. Congestion fronts give tortoise divergence; $\alpha = 1$ matches Schwarzschild dr/dt exactly. Fig. 44.

AU Einstein–Hilbert routes. Heat-kernel a_1 responds to curvature; Ollivier signs exact (flat/neg/pos on line/tree/ K_6); Clausius fixes $\varepsilon = \kappa/8\pi$, $G = 1$. Raychaudhuri-for-leg-bundles open. Fig. 45.

BB lensing + Shapiro + chroma. Fermat $n = 1/c_{\text{eff}}$ gives full $4M/b$ (naive $2M/b$ bet lost and recorded). Shapiro log matches GR to 5%. Chromatic extension uses *phase* velocity $1/24$, not group $1/8$ (conflating the two triples the prediction): r -independent factor drops out of Born, leaving 10^{-56} – 10^{-32} . Figs. 49, 55.

BH strain. Naive $dl = dr/(1 - \sqrt{\chi})$ gives $\gamma = 2$ (Cassini-excluded); tortuosity $dl = (1 + \sqrt{\chi}/2)dr$ gives $h = (1 + x/2)^2$, $\gamma = 1$ exactly, Mercury $42.99''/\text{cy}$. Peel-off $-0.75 M/a$ measured. PPN: $\gamma = 1$ derived, β unresolved (coordinate-confused, not quoted), $\alpha_{1,2}, \xi$, Nordtvedt untouched. Figs. 58, 60.




Figure 1: Gravity battery I: Fermat bending matches GR $4M/b$ to first order (left), Shapiro log pile-up matches (right); the middle panel records the pre-BH orbit gap (0 vs $43''/\text{cy}$) that the strain sector closes. **lensing, shapiro.**

BI/BQ weak-field OR. Grid+hub with flux stubs: radial OR negative in all configs (sign derived). Direct mode $p = 0.92 \pm 0.48$ ($L = 9$); $1/r$ collapse beats $1/r^2$ (CV 0.47 vs 0.60). Green-function walk fails by leg shorting (h -steepness $9.2 \rightarrow 4.1$); AS temperature triply load-bearing. Figs. 59, 64.

BV UV tortuosity-as-scattering. Legs as radial line defects ($\sigma = 4 \ln 2 l_p^2$), soft cost $\ln 2$ per encounter: dilute $c = 0.44$ – 0.60 (target 0.456 , geometric $1/\sqrt{\pi} = 0.564$) with zero tuning; $p = 2c$, $\gamma = 2c$. Mixed mode ($0.75 r_e$ core) rises then disconnects at k_{crit} — the pop without metric input.

Fig 58 — BH: tortuosity strain closes Mercury




Figure 2: Gravity battery II: naive strain gives $\gamma = 2$ (excluded); tortuosity $h = (1 + x/2)^2$ gives $\gamma = 1$ exactly and Mercury $42.99''/\text{cy}$ by direct geodesic integration. **strain**.

Boltzmann ladder (Dijkstra vs MC walks vs rays) shows routing worth $3\times$; graph- distance c agrees with Euclidean- r c (coordinate-free check). Figs. 70–72.




Figure 3: BV tortuosity from $\ln 2$ scattering: dilute $c = 0.44\text{--}0.60$ with zero tuning against the BU-implied 0.456 ± 0.025 band and the geometric $1/\sqrt{\pi}$ line. Right axis shows $p = 2c$. Hard cylinders (red $\times$) overshoot; soft and mixed modes agree. **uvscatter**.

S3 Quantum-information methods (A–O, Q–Y)

Log law $t_*(N) \approx \log_2 N / \log_2(1+p)$; MaxEnt $k \geq N s_{\text{node}} / s_{\text{leg}}$ plus fixed point $k^*(N) = 16\pi(\varepsilon N / l_p)^2 / 4 \ln 2$ (stable; BS gives $\varepsilon = c / \sqrt{N}$ and $\alpha = k^* / N = 1$ exactly); QES crossing iff $s_{\text{leg}} > l_p^2 / 4$ (Sec. 3 dies otherwise); Page wiring-only vs N -shrinks identical $A(t)$; QEC 99% at $k \geq N/2 + 1 + \log_2 100$; Kerr $k_{\text{eff}} = A / l_p^2$ monotonic in spin; exact Page $H(mn) - H(n) - (m-1)/2n$ with 0.72-bit dip; CKW verified; OTOC $\lambda + \text{MSS } 0.50\text{--}0.68$ ($\alpha \in [9.0, 12.4]$, 11.24 inside); TN $S_{\text{bdy}} = \min(N \log d, k \log D)$ fixing $\varepsilon \approx 0.047 l_p^{-1}$; SYK $N = 6, 8, 10$ vs Ising chain; GWTC-3 32 BBH $k_f > k_1 + k_2$; hardware

head-to-head 7.9 vs 14.6 steps. Figs. 7–20, 23–25, 30–33. Healing $dA/dt = (4 \ln 2 k l_p^2 - A)/\tau$, $\tau = 11.24M$ (3.5 ms at GW150914 mass); ladder healing/scrambling/Page/ evaporation separated by $\sim 66\times$ to 10^{80} s.

What Page vs. island means here (honesty note). The Page curve is derived as random-state behaviour: idealised leg surgery $k \rightarrow k - 1$ is imposed and $S_{\text{rad}} = \min(t, N_{\text{eff}} - t)$ holds by construction (module `evaporation`), but the qubit toy now has an explicit unitary completion (module `evaporation_unitary`, Fig. 5): per-step V_t with $V_t^\dagger V_t = I$ verified, S_{rad} computed from ρ_{rad} tracks the finite-dimensional Haar/Page expectation (mean dev. 0.002 bits, $N = 8$), and finite-depth all:all circuits converge to Page by depth $\sim 5/\text{step}$ without assuming Haar. The QES/island crossing $S_{\text{no}} = k s_{\text{leg}}$ vs $S_{\text{isl}} = k \ln 2 + \max(S_0 - k s_{\text{leg}}, 0)$ is a two-saddle competition plus a discrete min-cut analogue (module `qes`), not an extremization of $S_{\text{gen}} = A/4G + S_{\text{matter}}$ from a gravitational path integral. The graph-dynamics evaporation isometry $V_k : \mathcal{H}_{\text{graph},k} \rightarrow \mathcal{H}_{\text{graph},k-1} \otimes \mathcal{H}_{\text{leg}}$ is closed at small- N exact-diagonalisation level (module `graphvk`: per-step $V_k = \exp(-iH_{\text{graph}}dt)$ from the hole adjacency, $V^\dagger V = I$ proved, all:all tracks exact Page to < 0.25 bits at $N = 8$ while the chain sags > 0.4); the remaining missing theorems are the large- N limit, k -backreaction, and the emission spectrum. Tracked as D1 in `docs/DEFERRED.md`.

What Kerr means here (honesty note). Only the Kerr–Newman area $A(M, a, Q)$ is assumed to set the leg budget $k_{\text{eff}} = A/4 \ln 2$ (module `kerr`; extremal Kerr half, extremal RN quarter; ratios patch-independent). Not derived from the graph: spin-induced quadrupole $M_2 = -Ma^2$, frame dragging $g_{t\phi}$, ISCO(M, J), or the Kerr QNM spectrum (the overtone toy is Schwarzschild-like; fundamental damping $\tau = 11.24M$ calibrated, footprint corrections $1/2k, 1/\sqrt{k}$ unobservable). Echoes are covered by a dimensional lattice-scattering estimate only: amplitude $R \sim (\omega/\omega_P)^2 \sim 10^{-80}$ at 100 Hz, energy $\sim 10^{-160}$ vs $O(0.01)$ detectability (~ 158 orders of margin, module `gwdata`) — not a leg S-matrix derivation. The exterior perturbation problem ($\delta\omega_{nlm}$ vs Kerr) is open. Hence GW250114 (SNR 80 area law + Kerr ringdown) and GW241011 (strongest $\delta\kappa$ bound to date, Kerr to factor ~ 2 resp. $\sim 10\%$ by parametrization) are currently consistent by construction, not passed predictions. GW250114 provides the QNM benchmark ($\delta f_{220} \sim 2\%$, $\delta\tau_{220} \sim 10\%$, $\delta f_{221} \sim 30\%$, $\delta f_{440} \sim \text{tens}\%$): a future graph-derived $|\delta\omega/\omega|$ above those thresholds fails. A future graph derivation $Q = -Ma^2(1 + \delta_Q)$ with $|\delta_Q| \gtrsim 0.17$ would be ruled out by GW241011. Tracked as D2 in `docs/DEFERRED.md`.

Kerr thermodynamics (conditional check). With $S = k \ln 2$ (saturated legs) and $A = 4 \ln 2 \cdot k l_p^2$, $S = A/4$; importing $A(M, J)$ gives $S(M, J) = 2\pi[M^2 + \sqrt{M^4 - J^2}]$, whose derivatives return $T_H = \sqrt{M^4 - J^2}/4\pi M(M^2 + \sqrt{M^4 - J^2})$ and $\Omega_H = J/2M(M^2 + \sqrt{M^4 - J^2}) = a/(r_+^2 + a^2)$, i.e. $dM = T dS + \Omega_H dJ$ (module `thermo`: identities $< 10^{-6}$, independent finite-step $\Delta M = T \Delta S + \Omega_H \Delta J$ to $O(d^2)$; super-extremal $|J| > M^2$ returns NaN, never clamped). Per-leg reading $T_H = (dM/dk)/\ln 2$; finite-step $T_k = (M(k+1) - M(k))/\ln 2$ shifts $-1/4k$ ($\sim 10^{-77}$ stellar-mass). Conditional consistency, not derivation: $M(k, J)$ from graph dynamics is the open debt (D2).

S4 Compact objects and 2PN (BU)

Gradient shells. $p_{\text{adj}}(s) = 0.85 + 0.015s$, 8–10 shells, exact EMD on full neighbourhoods, deterministic bridges $n \propto r^{\beta(N)}$. Measured: $N = 1020$ (10×102), $\beta = 1.24$, 80 graphs exact Floyd+LP (~ 545 s): $p = 0.913 \pm 0.049$ per graph, SEM 0.0055, stacked 0.911 ($R^2 = 0.956$), 80/80 fits ok, range 0.79–1.05. Distance to target $|0.913 - 0.92| = 0.007$ (0.24σ in p , $\sim 0.09\sigma$ in $\dot{\omega}$), $5\times$ precision margin (0.0055 vs 0.028). Flat control $p = 0.49$ is -10σ dead.

N-scale campaigns (BV). CSR-direct shell graphs + Sinkhorn OR (no NetworkX): $N=1020$ reproduced (0.9107 vs 0.9134); $N=4000$ $p = 0.9315 \pm 0.0032$ ($\beta = 0.99$); $N=8000$ $p = 0.9382 \pm 0.0030$

Fig 10 — D: leg-surgery evaporation + Page curve




Figure 4: Leg-surgery evaporation: radiation entropy follows the Page curve (left, turnover at half evaporation); horizon area is identical whether N shrinks with k or stays fixed (right) — area tracks wiring alone, the distinctive Claim 2 prediction. **evaporation.**

($\beta = 0.87$, pods); $N=16000$ $p = 0.9137 \pm 0.0022$ ($\beta = 0.74$, GPU farm). Exact shell distance oracle + torch backend; $\beta(N)$ log-linear over 6 points. Local- p turnover $0.63\text{--}0.69 \rightarrow 1.26\text{--}1.46$ measured (inner windows flatten — the UV turnover). Kill wire: p outside 0.92 ± 0.056 at $N = 1024$ class fails.

2PN and pulsars. $h = (1+x/2)^2$ gives $c_1 = 3.36$ vs GR 1.94 (73% excess); naive fixed- M gives B1913 6.2σ , J0737 9.7σ vs real TOAs. With $c_2 = p(2p-1)$, $w = 1.953$: $c_{\text{tot}} = 4.8693$ at $p = 0.92$ vs GR 4.8695 (0.00σ fixed- M). Self-consistent $R + \dot{\omega} \rightarrow M$: $\Delta M = -11.2$ ppm naive (-4.5 ppm with GR g_{rr}), $\Delta \sin i = 3.7 \times 10^{-6}$, s_{obs} passes at 0.36σ ; B1913 0.01σ . Bending/Mercury 2PN pieces hide below VLBI/astrometry (Fig. 69, right). The companion toy-generator route (dense layers, $k_0 - c_2/r^2$ fits) crosses zero in κ and needs a κ -tweak fudge; power-law fits better on our profiles (R^2 0.91 vs 0.81) but the two $\kappa \rightarrow c_2$ maps disagree (0.82 vs 3.14) cross-applied — hence the open-map label. e_{int} weighting *lowers* our p ($0.94 \rightarrow 0.84$ at 0.9, 0.78 at 0.99): uniform measure is both principled (continuum theorems) and required.

Leg-shedding kilonovae. $M_{\text{ej}} = \Delta k m_{\text{leg}} \times 0.1$, blue ($v \sim 0.3c$, $t_{\text{peak}} \sim 0.6\text{--}1$ d) + red ($v \sim 0.1c$); $L \propto t^{-1.3}$ tail. Band mags assume BC = 0 and per-component peaks; POSSIS refinement queued. g -band verdicts robust; i -band NOT claimed (one-zone $\kappa = 10$ over-traps: $t_{\text{red}} \sim 10$ d vs observed ~ 4 d decline, m_i faint by $\sim 2\text{--}3$ mag — conservative for gap- g detectability). Standard 2–28% band is a literature EOS input, not derived.

O5 protocol. Trigger: $m_1 \in [2.5, 5.0] M_{\odot}$, < 200 Mpc, 90% area $< 100 \text{ deg}^2$ (multi-detector); Rubin ToO epochs 0.5/2/5 d in $g+r+i+z$ (DECam/Gemini backup). Kill rule: 10 qualifying non-detections kill BU. Archival ledger: GW230529 uninformative (single-detector, 93% unsearched); S250206dm unverified (check GCNs before citing). Full plan: docs/observation-protocol.md. Figs. 66–69.

Mass-gap audit (module massgaps). Lower gap: $k(M)$ log-log curvature $< 10^{-9}$ over $0.7\text{--}150 M_{\odot}$ (no forbidden masses); $e_{\text{int}} \sim e_{\text{ext}}$ location descriptive (calibration inherited from AT2017gfo). Upper gap: spin-leg ratio $1 \rightarrow 0.857 \rightarrow 0.5$ at $a = 0 \rightarrow 0.7 \rightarrow 1$ with no jump on $[0, 0.99]$ plus exact extremal continuity; Love k_2 slope -2 everywhere; leg-band entry $\sim 32 M_{\odot}$ at 10 Hz with $dM/M \sim 10^{-81}$; congestion ladder $\chi \sim 10^{-8}$ (He core), 10^{-14} (envelope), 0.12 (NS surface), ~ 1 (gap masses at 10 km footprints, footprint-conditional), 1 at horizons. Shedding ex-

Fig 74 — D2: unitary evaporation ($V^d V = I$, S from ρ_{rad})




Figure 5: Unitary completion of the qubit toy: sequential V_t (left) tracks the finite-dimensional Haar/Page expectation including the 0.72-bit dip, computed from ρ_{rad} rather than $\min()$; finite-depth all:all circuits (right) converge to Page by depth $\sim 5/\text{step}$ without assuming Haar. Module `evaporation_unitary` (D2).

Fig 66 — BU: 2PN cancellation at $p=0.92$ (J0737 0.00 sigma)




Figure 6: 2PN cancellation: model $c_{\text{tot}}(p)$ crosses GR at $p = 0.92$ (left); fixed-mass J0737 residual vanishes there (middle, green 2σ band); self-consistent $\sin i$ passes for both GR and $p = 0.92$ (right). pulsar.

actly flat in q (machine precision) — but q -independence is prescription (shed fraction $\times$ efficiency fixed once on AT2017gfo), not derived from the graph-dynamics evaporation isometry V_k (open, S1/S3/D1): $q \sim 0.11$ (GW190814) flashing at the same fraction as $q = 1$ (GW170817) is the model's biggest theory bet. GW190814 epoch audit: CFHT g-band (Vieira+2020) and GROWTH i-band detection limits (Andreoni+2020) with per-epoch coverage $\times$ depth against distance (241 ± 43 Mpc) and ± 1 mag theory errors; combined $P \approx 0.68$ (g-only) / 0.88 (g+i at $g - i = 0.7$). Erratum on record: the first audit used a nominal ZTF depth, but ZTF ran no targeted GW190814 follow-up; superseded by the CFHT/GROWTH epoch tables above. Fig. 75 (main text). Analytic systematics (round 4): viewing $A_g \sin^2 \theta$ ($A_g = 1.25$) + opacity-rescaled blue mags give the P grid in Fig. 75b; the hiding window is bounded, not open-ended.

S5 UV nulls and phenomenology (T, U, V, BD–BF)

TeV: $M_D = 1$ TeV gives $k \approx 11$ (3 TeV) – 17 (13 TeV) vs $k_{\text{crit}} \approx 50$ — thermal nulls expected, onset ~ 550 TeV. Echoes: $\Delta t = 4M \log(M/l_p)$, 0.03 – 0.3 s (GW150914 0.11 s), in searched windows but $R \sim 10^{-160}$ unobservable. EHT: $l_p/2M \approx 10^{-48}$, 47 orders below sensitivity. Dispersion:

Fig 67 — BU: gradient OR gives $p=0.92$ (exact EMD)




Figure 7: Gradient-shell Ollivier–Ricci: stacked $|k|(r)$ steepens with the gradient (left); per-graph p histogram centres on target (middle); 80-graph SEM extrapolation clears the $1\sigma/2\sigma$ precision bars (right). Production $N=1020$: $p = 0.913 \pm 0.049$, SEM 0.0055. **orici**.

$E_{QG,2} = \sqrt{8} E_P$, Fermi-safe by 10^8 ; universal GW extrapolation shifts 100 Hz by $\sim 10^{-82}$ (LVK $\alpha = 4$, safe by $\sim 10^{60}$ — a recorded null, not a falsifier; no GW sector claimed). Chromatic lensing 10^{-56} – 10^{-32} . Figs. 21–22, 39, 51–56.

Fig 51 — BD: quadratic lattice dispersion, Fermi-safe




Figure 8: Lattice dispersion: band bending at the Brillouin edge (left) with group velocity even in k (middle, no linear term by symmetry); GRB arrival delays sit orders of magnitude below detectability across GeV–TeV (right). **dispersion**.

S6 Kill list (AN + BU)

AF quench ratio < 1.3 ; α outside $[9.0, 12.4]$; $A \propto N$ in any TN calculation; thermal LHC excess below k_{crit} ; $s_{\text{leg}} \leq l_p^2/4$ in any physical state class; any linear-LIV signal; p outside 0.92 ± 0.056 at $N = 1024$; graph-predicted $|\delta_Q| \gtrsim 0.17$; simulator d_{eff} shape-mismatch after the cost mapping is fixed (D10/P0' wire); 10 gap non-detections; O5 nearby-BBH ToO sample clean (universal shedding, Eq. 2 main). Field-program wires (S12): a second admissible M_O with robustly non-3D IR refutes reconstruction universality; $I > 0$ in any collision cell beats the linearity null; a firing trigger derived from the instantaneous ontology overturns BR27-NO-MODE (strong stop: no rate shopping; TRIGGER-0 census: 19 conditions, 0 implications); an earned local covariant rewire selector overturns REWIRE0-DEGENERATE; a tension geometry with no U-shape under the fixed cost rule, or amplitude anti-scaling in χ , kills the tension-imprint conjecture. One sub-claim (broad remnant DM) already ruled out on the record. Figs. 37–39.

S7 N-scaling and calibration record

Drift check: $\beta = 1.5$ calibrated at $N = 300$ drifts high at $N = 600$ (1.16 ± 0.06); recalibration to 1.28/1.24 recovers $p \approx 0.92$. Pre-run $1/N$ extrapolation predicted $\beta(16000) \approx 1.18$; measured 0.74 — drift faster than $1/N$, now fit log-linear. $N=1020$ cross-check: CSR-direct 0.9107 vs NetworkX 0.9134 (agree to 3×10^{-3}). Local- p turnover (inner windows flatten, 0.63–0.69 $\rightarrow$ 1.26–1.46) is measured at 4k/8k/16k, not extrapolated. Fig. 73.

Fig 73 — BV: $N=4000/8000/16000$ campaigns (80 graphs, CPU+pod+farm)




Figure 9: N-scale campaigns: $\beta(N)$ recalibration drifts faster than $1/N$ (left); stacked $|\kappa|(r)$ profiles agree in shape across 1020–16000 (middle); local- p windows show the UV turnover with SEM error bars (right). 80 graphs per production row. **shellscale**, **sinkor**.

Reproduce ($N=1020$, ~ 9 min):

```
measure_p(per_shell=102,n_shells=10,n_graphs=80,
gradient=True,beta=1.24,seed0=0,max_per_shell=8)
```

S8 PPN ledger and walk/foam/healing notes

Walk no-go (BP/BQ). All smooth weight rules give drift slope -2.99 ; only the absurd $e^{2\sqrt{d-d_0}}$ escapes. Constant- μ persistence keeps -3 analytically; $\mu(\chi)$ fluctuation walk gives $-1.95 \approx -2$ modulo the labelled $\sqrt{\chi}$ assumption. Harmonic/hitting-probability walks fail by leg shorting. Conclusion: temperature (AS), not bare graph diffusion, carries Newton. Figs. 63–64.

Foamgrid FDTD (BF). Beams on defective fabric: centroids unbiased (jitter, average straight) with deficit $\propto \omega^2$. Bounds assume independent edges; correlated patches of size ξ relax them by $\sqrt{\xi}$. Fig. 56.

Healing ladder (AG/AH). $\tau_{\text{heal}} = 11.24M$ ($1/\text{Im } \omega_{220}$); GW150914-mass ladder: healing 3.5 ms, scrambling 0.23 s, Page $\sim 10^{79}$ s, evaporation $\sim 10^{80}$ s. MSS $\lambda/2\pi T = 0.50$ –0.68; LVK $\alpha \in [9.0, 12.4]$ contains 11.24. Figs. 30–31.

S9 Archival ledger and O5 trigger (BU detail)

Trigger (full plan docs/observation-protocol.md): $m_1 \in [2.5, 5.0] M_\odot$, < 200 Mpc, 90% $< 100 \text{ deg}^2$; Rubin ToO 0.5/2/5 d in $g+r+i+z$ (DECam/Gemini backup). Sought: fast blue ($v \sim 0.3c$, $t_{\text{peak}} \sim 0.6$ –1 d) $\rightarrow$ slow red with $L \propto t^{-1.3}$ tail. Yield: 1.5 gap/yr $\times 70\% \approx 1.05$ /yr ours vs ≤ 0.3 /yr standard. Yield arithmetic in `collapse_gap_o5_yield`; brightness in `collapse_peak_apparent_mags`; GW190814 epoch audit + systematics grid in `massgaps.gw190814_combined_pdetect` / `massgaps.gw190814_systematics_table`. Figs. 68–69, 75 (main text), 75b.

Fig 30 — AG: healing lag + timescale ladder




Figure 10: Healing lag: wiring jumps at merger while horizon area catches up on the ringdown time (left); timescale ladder from milliseconds to 10^{80} s (right). The final sigh carries $\sim 10^{-78}$ of the mass. healing.

BBH extension (this work): the same shedding law predicts $m_g \sim 22$ transients for nearby BBH, so the ToO program extends to well-localized ($< 100 \text{ deg}^2$) BBH within 200 Mpc with identical epochs and kill accounting kept separate from the gap sample (a clean BBH sample kills Eq. 2 main even if gap events flash). Full thresholds in docs/observation-protocol.md.

Fig 69 — BU: O5 detectability + GR-battery 2PN margins




Figure 11: O5 protocol: gap kilonovae visible in g -band to 200 Mpc with margin over Rubin/DECam depths (left); yield $\sim 1/\text{yr}$ vs $\leq 0.3/\text{yr}$ standard (middle); the same p keeps bending/Mercury 2PN pieces below VLBI/astrometry (right). collapse, pulsar.

S10 Module map and reproduction

Modules follow the appendix letters of the v4.1 extended record: `graphs`, `scrambling`, `emergent_dim` (L0 dimensions), `horizon`, `micro`, `circuits` (A), `maxent` (B), `qes` (C), `evaporation` (D), `evaporation_unitary` (D2), `qec` (F), `robustness` (G), `kerr` (H), `thermo` (H2), `haar` (I), `monogamy` (J), `otoc+pheno` (L), `tn` (M), `kerrpage` (N), `syk` (O), `data` (Q), `litcompare` (R), `tev` (T), `echoes` (U), `posteriors` (W), `ds` (X), `krylov` (Y), `collapse` (AA), `cosmic` (AB), `lunch+remnant` (AC), `bounds` (AE), `healing` (AG), `mss` (AH), `big syk` (AI), `mp+greybody` (AJ), `congestion` (AK), `charge` (AL), `bandwidth` (AM), `gridcirc+monitor+selfattack+lhc` (AN), `concentration` (AO), `ps` (AP), `scatter` (AQ), `emd+viability` (AR), `entropic` (AS), `redshift` (AT), `heatker+orici+jacobson` (AU), `fission` (AV), `klanguage` (AW), `tension` (AX), `gw250114` (AY), `overtones+tensionvol` (AZ), `sparse24` (BA), `lensing+chroma+shapiro` (BB), `bcrit` (BC), `dispersion` (BD), `qnmfoot+qnmlegs` (BE), `foamgrid` (BF), `perwalk` (BG), `strain` (BH,BJ), `weakfield` (BI,BQ), `legham` (BL), `pulsar+orici gradient shells+collapse leg-shedding` (BU), `uvscatter+sinkor+shellscale` (BV), `massgaps` (mass-gap null + GW190814 epoch audit). Update-rule and dimension programs: `update_rule` (D1 tournament), `blind_u` (D1 blind- U), `weighted` (D10/D14 pricing), `spectral`

Fig 75b — GW190814 systematics: equatorial + kap=2 hides ($P \sim 0.18$)




Figure 12: GW190814 analytic systematics (round 4; full POSSIS queued, D7). Left: $P(\text{detect})$ g -only over viewing angle $\times$ blue opacity (CFHT epochs, no color term): fiducial face-on $\kappa = 0.5$ gives 0.68; equatorial + lanthanide-mixed ($\kappa_{\text{blue}} = 2$) drops to ~ 0.18 . Right: face-on hiding bar per CFHT g epoch (g 1.7 d needs ~ 1.6 mag). Module `massgaps`.

(D10 second leg), `formation` (D14 C2/SSB-1/Stage-0). Field program (S12): `ballistic` (P1 waves), `coherence` + `slit` + `tunnel` (interference), `potential` (POT-0) + `driven` (POT-1), `continuum` (EM-0) + `falsification` (EM-1), `malus` (sheet sectors), `obs0` + `obs0r` + `obs1` + `obs1_reveal` + `quot` (rulers, blind observer, quotient), `backreaction` (BR-0) + `phase` (BR-2) + `rigidity` (BR-1), `contraction` (BR-2.5) + `conservation` + `accounting` (CONS-0/BR-2.6) + `stability` (BR-2.7), `u0` + `time0` + `rand0` + `measure0` + `sym0` (dynamics completion + state census), `field0` (linearity null), `response` + `bgresp` (kernel + vacuum susceptibility), `vac0` + `vacfield` + `vacexc` + `vaccomp` + `vacselect` (Final MIXED + vacuum family), `hidden` + `hiddenbr` (dead-sector info + response), `zero` (nodal census), `grav0` (graph-only null), `split0` + `rewire0` (inverse fiber + rewire null), `merge0` + `reservoir0` + `trigger0` (update + account + conditions), `fiber0` + `info0` (fiber debt + exact books), `vacdomain` + `vactexture` + `vacstab` (vacuum trio), `source0` (boundary data). Side apparatus: `spectroscopy` (SPEC0 null), `stern_gerlach` (SG gate fails), `fep` (FEP0 null), `mergershed` (BU2 q -shape derived, flat law kept as O5 falsifier), `graphvk` (D1 graph instance closed at small- N ED).

Reproduce:

```
pipinstall-e".[dev]"
pytesttests/-q
pythonscripts/generate_figures.py
streamlitrunapp.py
```

S11 Emergent-dimension methods (L0)

Protocol. For a graph state, pick an information-transfer distance $d(i, j)$ (NOT bare adjacency: shortest-path is the documented failure mode on K_N , diameter 1 at every N), build balls $B(r) = \{j : d(i, j) \leq r\}$, measure capacity $V(r)$, and fit $d_{\text{eff}}(r) = d \ln V / d \ln r$. Observed dimension is the infrared conjecture $d_{\text{obs}} = \lim_{r \rightarrow \text{IR}} d_{\text{eff}}(r) \simeq 3$ (module `emergent_dim`; 12 tests). Total emergence requires the two-distance bracket: shortest-path balls (adjacency scaling) and diffusion balls (information-spreading scaling) must agree at 3 on the same graph with nothing dimensional imposed.

Ruler validation (imposed lattices, circular for emergence). Shortest-path converges from below (exact BFS shells): chain $0.92 \rightarrow 0.96 \rightarrow 0.98$ ($N = 60/200/600$), 2D $1.67 \rightarrow 1.77 \rightarrow 1.82 \rightarrow 1.87$ ($L = 20/40/60/100$), 3D $2.25 \rightarrow 2.46 \rightarrow 2.56 \rightarrow 2.63$ ($L = 7/11/15/21$), gaps $\sim 1/\sqrt{L}$ (Manhattan-diamond shape + open box). Diffusion recovers dimension from above at tuned t (chain $t = 20 \rightarrow \sim 1$, 2D $t = 5 \rightarrow \sim 2$, 3D $L = 7, t = 5 \rightarrow \sim 3.3$); the 3D overshoot shrinks with size

(3.56 $\rightarrow$ 3.34 $\rightarrow$ 3.14 at $L = 5/7/9$, $t = 5$). Degree-3/4/6 lattices all scale 2D: dimension comes from 2D-ness, not coordination.

Rejections (negative pins). Effective resistance fails in 2D (log-growth $\rightarrow$ spurious large p); communicability fails on chains (similarity, not separation). Radial shells fail both sides (shortest ~ 1.4 , diffusion ~ 0.6): radial-only wiring gives $\sim 1D$ with no 2-sphere factor — the missing-area diagnosis that motivates P0'. On K_N , shortest is trivial and diffusion uniform, by design (T1: no interior geometry).

Tense plug (D10a). A 5×5 clique plug at the centre of a 40×40 grid: tense-centre mid-window ($r = 8\text{--}20$) scaling dips below fabric (~ 1.52 vs ~ 1.89 control) and recovers from below — *inverted* vs the GR far side (above 3). Bare shortest-path is therefore *rejected* as $d(i, j)$ for tense regions: shortcuts shrink balls while GR mass stretches rulers. Costs must be congestion-weighted (D10b gates D10a), consistent with the light sector ($c_{\text{eff}} \rightarrow 0$, tortuosity), which never counts hops. Companion pin: far-source near balls ($r = 2\text{--}6$) are bit-identical to control — relaxed fabric next to tension measures exactly relaxed. Far-field tail (D11, closed): $\chi \sim 1$ decays $1/r^2$ ($E \cdot r^2$ flat to $10R_c$), $\chi \sim 2$ decays $1/r$ ($E \cdot r$ flat to $10R_c$), $\chi \sim 5$ slow crossover (q 1.19 $\rightarrow$ 1.10 to $20R_c$; $1/r$ supported, not firm) — the exponent is tension-dependent. κ -profile (D10b, closed): corner-free disk plug reproduces the boundary-negative/core-positive pattern quantitatively (+0.90/−0.86/0.00 vs clique +0.89/−0.93/ ~ 0); mild-disk matches mild-square (−0.05/−0.23/0.00).

GR fingerprint (D10 target, computed). Weak-field uniform star, proper balls of proper radius l : $d_{\text{eff}} \approx 2.990$ at $l = 30M$ (near dip), 3.030 at $100M$ (overshoot), 3.010 at $1000M$, 3.0017 at $10000M$ ($\rightarrow 3^+$ as $\sim 1/l$); amplitude $O(M/l)$, $\sim 10^{-9}$ at Earth's surface. The far-field overshoot is structural: a localised mass leaves $V(l) = \text{flat} - \text{deficit}$, so V/V_{flat} rises toward 1, which *is* $d_{\text{eff}} > 3$. Close criterion (D10a): a tensed simulator region reproduces this *shape* under info-side d_{eff} once the cost mapping is fixed; shape failure refutes P0'. Reproduce: `pytesttests/test_emergent_dim.py-q`.

Reconstruction universality (D12, open). The observer map M_O is not unique on its face: closeness, resistance, diffusion, and communicability distances are all candidate reconstructions. D12 demands an access class $\mathcal{A}_{\text{macro}}(G)$ defined from G alone — the information-access floor is part of the definition, and neither “3D” nor “Lorentzian” may occur in it — such that every admissible M_O recovers the same infrared M up to coordinate/coarse-graining equivalence. Falsifier: a second admissible M_O giving robustly non-3D IR on relaxed fabric refutes the P0' reconstruction program. Partial credit runs criterion stated $\rightarrow$ two instances agree on vacuum $\rightarrow$ agreement extends to tensed regions. P4 (uniform Ollivier) audited against the five: criteria 1, 2, 3, 5 pass (permutation-covariance and drift-localization tested, the latter on flat and plug backgrounds), 4 partial. Observer-indexed extension filed ($M_O(G, o)$: sky chart + relational depth, vantage-covariance as a second universality axis); static depth bake-off negative (best principled $d_{\text{obs}} = 2.28$ vs tuned-null 2.73; MDS readout likewise negative, logC gap shrinking with N). Adopted reframing: holographic DOFs + dynamical relational distance $\rightarrow$ emergent localization, M_O demoted to readout; rank-3 claim needs an N -stable λ_3/λ_4 gap with non-radial v_3 . Tier-1 MDS calibration pinned: lattice/gabriel/kNN hop show λ_1, λ_2 dominance, while Delaunay hop carries a radial tortuosity bowl ($v_3 \leftrightarrow r^2$ at 0.93) — the MDS bar is per-substrate, and eigenvector geography is part of the test. Shell-counting dilemma sketched: honest per-shell counts on 2D fabric (nodes $\sim r$, cut $8r+4$, MDS rank $O(1)$ with $\text{GoF}_2 = 0.772$) never reach r^2 , while cumulative-ball r^2 violates shell-independence by construction — $d_H=2$ XOR independence, with non-geometric C_O or non-shell composition left as escapes. The dilemma binds hop shells only: the MDS-rank route is independent, and dynamical metrics redefine shells (first shipped dynamical distance: SI first-passage, MDS null cleanly 2-dominant with $\lambda_2/\lambda_3 \approx 17.5$).

Staged causal order (D13). Time undergoes reconstruction exactly as space does: the mi-

croscopic update index n , the causal order $\prec_U$, and clock time τ are three distinct orderings. The static control (D13.0) is measured: the analytic Manhattan law $V_G(r) = 1 + 2r(r + 1)$ bit-pinned at $L = 40/80$, the exact diamond perimeter cut law $\text{cut}(r) = 8r + 4$ on unclipped $r \in [5, 18]$, disk-boundary cut capacity deficit at $r \leq 4$ with exact recovery from $r = 5$, and the hop/weighted plug sign pattern ($35 < 38$ hops vs $+5.0$ weighted delay) pre-registered for dynamical confirmation. Substrate family measured under one frozen rule: triangular/hexagonal/noisy-grid controls reproduce $d_G \rightarrow 2$ (exact shell/cut laws on the lattices, statistical pins under disorder); Tier-1 extension: Poisson-Delaunay (reference member), Lloyd-relaxed, Gabriel, k-NN, and Delaunay-medial quadrangulation all give $d_G \rightarrow 2$ with linear cuts, die on levered swaps, and survive $q = 0.05$ deletion, while span-limited swaps preserve 2D-ness — degeneracy across topology/degree/order is the expected universality signature, and the physical object is the class $[G]_{\sim_O}$, not a winning tessellation. Spectral dimension is the second leg: torus heat-trace $d_s(t)$ holds $(1.95, 2.05)$ over $t \in [10, 100]$, and Weyl counting fits land every member in $(1.90, 2.15)$. Exact-UIPQ sampling stays queued (free-walk labels are not conditioned well-labeled trees); Lloyd is excluded from spectral comparison (unclipped relaxation coalesces points above $N \approx 1600$). gated-wall negative control keeps $\sim r^2$ balls with collapsed cuts; rewired-grid fragility control plus rewiring-fraction sweeps give measured $N_{\text{med}}^* = (20, 10, 10)$ at $L = 40/60/80$ ($N^* = O(1)$ consistent, $\alpha \simeq 0$ over the tested range; asymptotic $f^* \rightarrow 0$ hypothesis) with hand-placed dose-response dissolving the placement lottery. RG blocking (2×2 , $L40 \rightarrow 5$) flows rewired long-edge density $0.013 \rightarrow 0.31$ against plain 0.0 (rough $y_\lambda \approx 1.5$). Headline: low-dimensional locality is not generic; it must be protected — vacuum as a dynamically protected locality class whose preservation ($U(G_{\text{vac}}) \in \mathcal{C}_{\text{vac}}$, self-healing vs amplification under shortcut injection) is now a D1 falsifier. BFS first-passage is hop distance, so this control is static geometry, not propagation. Open stages: (1) $U \rightarrow T_U$, $\prec_U$ via counterfactual-influence arrival times; (2) observed cone under the *same* M_O producing R^3 ; (3) internal clocks $d\tau/dt(\chi)$ reproducing the T9 profile unimported; (4) interval $g_{\mu\nu}$ from $(\prec_U, V_U, M_O)$, where the spacetime-volume observable V_U (e.g. calibrated event counts in causal diamonds) is independently defined — spatial $V(r)$ is reachability, not event volume. Malament’s theorem is motivation conditional on the continuum correspondence (order fixes the conformal class; V_U the factor), not an earned mapping. Companion constraint on U (D1, first pass): six update rules tournamented against shortcut damage all stall with residual longs (best single-move: slide rule, $30 \rightarrow 5$) while p -healing proves gameable (greedy perfects p to 10^{-6} while tripling longs) — judged on (p, longs) jointly. Second pass: coordinated double-swap breaks the floor ($30 \rightarrow 1$, pair-locked residual) and the visibility-chained triple endgame fully heals (longs $\rightarrow 0$, two seeds plus torus $2/2$; triple-lock localizes to boundaries) — locality protection demands coordination order 3 with detour-aware proposals, order 4 with a strict census gate for corners (ungated order-4 scrambles: $10/10$ accepts while longs $1 \rightarrow 5$). Gate duality filed and split by a temperature knob: strict order-3 clears dm4 in 0.5 s but stalls seed7, $T_0=2$ annealing clears seed7 — neither pole dominates (stall topology differs). The knob replicates at order 2 (pair_anneal matches-or-beats pair on every stream); gate-default adoption rejected (worse-on-s5 default is bad science). Reproduce: `pytesttests/test_scrambling.pytests/test_substrate_family.pytests/test_spectral.py-q` (D13); `pytesttests/test_update_rule.pytests/test_blind_u.pytests/test_weighted.pytests/test_j2.py-q` (D1/D14/J2).

Dynamics coherence (C1–C5, over-arching). $M_O \circ U \simeq U_{\text{eff}} \circ M_O$ in five levels: coherence $\rightarrow$ locality $\rightarrow$ autonomy $\rightarrow$ universality across $\mathcal{A}_{\text{macro}} \rightarrow$ GR limit in the IR, each presupposing the previous. Autonomy (C3) is defined by the twin-histories criterion $M_O(G_a) = M_O(G_b) \Rightarrow M_O(U(G_a)) \simeq M_O(U(G_b))$; factoring M_O through future-equivalence classes does *not* imply it (two-bit swap counterexample: singleton classes, failing twins).

Update-rule tournament, third pass (D1). Blind- U reframing adopted: stationary-ensemble vacuum with U -admissibility (drift / square / triangle first entrants; Metropolis no-

window, grid-not-optimum, and kappa frozen-descent negatives; triangle-landing misdirection plus degree-fiber reachability). Cross-candidate battery opened (triangular healing generalizes $24 \rightarrow 13$; hex needs fabric-relative radius). Coordination need is substrate-dependent: triangular pair alone fully heals ($24 \rightarrow 0$, p bit-plain); hex floor resolved (leaf edges, interior census 0, guillotine $54 \rightarrow 17$); Delaunay reference heals at order 2 (pair600 $21 \rightarrow 0$); Gabriel hits a broad-spectrum census limit (descent overshoots below plain); medial mixed (strict gate $6 \rightarrow 0$ in one accept); k-NN mostly-heals (floor 16); Lloyd heals via chain (pair grinds to 1, strict clears). Self-calibrating census (median-relative; exact on gapped, Gabriel unseparated). Reproduce: `pytesttests/test_update_rule.py`, `pytesttests/test_blind_u.py-q`.

Pricing-hierarchy origin (D14). Weighted-graph audit adopted (event sweep + edge census; 3-vertex cross-checks; weight-tolerance recovery at $L_w=20$; T15 cites Prop. 1). Weighted RG: $L_w=10$ washes out by level 3 (pricing irrelevant above span scale; vacuum RG-protected, no pumping needed); Φ -apparatus with tolerance contour; knot pilot (dual ruler, no bundle mixing, ruler-trap law). Experiment A (Form-1 relaxation): control bit-exact, $\beta=350$ recovers the banked slice ($(36, 23, 7, 0)$ vs $(36, 23, 5, 0)$), Δy_w split (substitution at price level, span-selection at Φ level). Walk-atrophy killed (0/8 horizons plus inverted signal; avoidance is conductance math). Betw-congestion closure: feedback attenuates ($m=1$: factor 0.684, in-basin cross at $k=3$; $m=4$ exit is a cadence artifact), concave- D holds — static χ closed. Stateful-scale derivation: linear memory equals static (LTI steady map), nonlinearity buys shape not magnitude (characteristic-scale lemma), YES-prong criterion (gain must be a structural number; unoccupied). Gain-free discriminator fails (weak median 6.2, rows $(39, 30, 20, 4)$, no crossing; threshold narrowed to $(6.2, 11.0)$): the gain is debt by elimination, and magnitude origin passes to formation/dynamic topology. Reproduce: `pytesttests/test_weighted.py-q`.

D14 formation design (C2 machine). E-conserving relocation on ER/RR soup: C2-PILOT-1 topological verdict DARK (54/54, 0/54 WEAK over a $20 \times$ floor sweep; D1 null holds as Poisson equilibrium; D3 hope dead — 13 dust-shedding arrests + 23 stillborns, zero active; no concentration pathway, so the D5 triangle-closure lead). C2-PILOT-2 geometric verdict NO-coexistence (66 + 6-repair runs, bit-identical cross-machine): CONDENSATION 9 ($D5\kappa_2$ @1600 single-clique) vs FRUSTRATED 15 ($D5\infty$ all- N , kmax-4 churn limit, N -independent, active-7% EXCHANGE) vs POISSON+ 39; $\kappa_c \in (1, 2)$, nucleation window $N^* \in (1600, 3600]$, kinetics mirror-broken, refined D3 ... $D5\infty$... $D5\kappa_2$ bracket (frustration-as-selector lead). SSB-1: core selection SPONTANEOUS (same-soup Jaccard $\approx$ random), so D15 reopens with symmetry restated as statistical- S_N -of-law. Stage-0: blob kinematics NULL (0/6 directed, 0/6 persistent-handed, achiral plateau) — the debt-free pure- $D5\infty$ route stops and polarity pauses (individuals yes, circulation-sign no). Reproduce: `pytesttests/test_formation.py`, `pytesttests/test_closure.py-q`.

J2 micro/macro probe (D10). $J_2 = \mathbb{Z}^2 \rtimes \mathbb{Z}_2$ walk graph (degree 8; substrate of D'Ariano–Erba–Perinotti, Phys. Rev. A 100, 012106 (2019)), pre-registered as a probe, not Tier-1-track (index 2 over $\mathbb{Z}^2$, hence quasi-isometric to the square lattice: long-scale $d=2$ is a control). Quotient verified computationally (1861 cells, every edge square-adjacent, 1741 interior cells square-4 with micro-multiplicity 4, quotient shells exactly $4r$); exact micro laws (shells $8r$ for $r \geq 3$, cuts $32r+16$ for $r \geq 2$); bipartite correction filed ($q=x+y$ bipartition, odd returns vanish exactly — a pre-registered aperiodic prediction falsified with mechanism); 4-cycle census 26072 at $R=18$ ($\sim 20 \times$ square-equivalent); perturbation family-typical (swaps $2/3$ kill, q -deletion robust) — no tier claim. Orientation program (D14/D15, 32 runs on the J2 torus): cores form despite the pristine triangle-free substrate (kmax-4, triangle-free bootstrap), radial SSB preserved, but orientation absent (random-axis elongation, edge $x/y \leq 8\%$, quadrant signs null-consistent, sheet 0.49–0.50) — verdict (c)-ISOTROPIC, so $D5\infty$ -radial is the wrong kind for compass consumption (D15 stays closed on this candidate). J2 is adopted as the canonical working vacuum substrate (`docs/j2-status.md`;

evidence-based, non-uniqueness explicit). Reproduce: `pytesttests/test_j2.py-q`.

S12 Relational field program (F-layer)

Postulates and anatomy. FP1: two real scalars per node, $\psi_u = r_u + is_u$ (no vector, momentum, or coordinate register). FP2: $i\partial_t\psi = H\psi$ with $H = -A$ (frozen $J = 1$), the one-way $G \rightarrow \psi$ map, postulated as the working law. Granted only this, VAC-0A derives the anatomy for every simple undirected graph (A1–A6 identities on hostile graphs $\rightarrow$ LAW): conserved $\rho_u = |\psi_u|^2$, $Q_\psi = \sum_u |\psi_u|^2$; bond energy $B_{uv} = \text{Re}(\psi_u^* \psi_v)$; continuity current $J_{u \rightarrow v} = 2 \text{Im}(\psi_u^* \psi_v)$ with $d\rho_u/dt + \sum_v A_{uv} J_{u \rightarrow v} = 0$; energy $E_\psi = \langle \psi | H | \psi \rangle = -2 \sum_{(uv) \in E} B_{uv}$ and the conjugacy $\partial E_\psi / \partial A_{uv} = -2B_{uv}$ (first pinned on the J2 stack inside EM-0). Quadrature $B \sim \cos \Delta\theta$, $J/2 \sim \sin \Delta\theta$; some readouts store the bare imaginary part (exactly half the current — convention, not a second current). Reproduce: `pytesttests/test_vac0a_identities.pytests/test_vac0_battery.py-q`.

Propagation, response, rejected EM reading. POT-0 separates direction from any node arrow (POT0-COLLECTIVE: symmetric source spreads, coherent packet goes ballistic, scrambling kills direction, coherence restores it). POT-1 + EM-0 give one field with a static regime and a propagating front on source change (EM-0 continuum reading: J2 Bloch bands, long-wave Schrödinger sector, continuity to numerical zero; no Maxwell/charge/photon claim). EM-1 rejects the electromagnetic identification (EM1-FALSIFIED): static range saturates ($\xi \approx 0.53$, range 3), one propagating scalar mode, local phase moves $B/J/E$ by order one, nodal drift tracks $v \cdot q$ not $v|q|$; the conserved signed charge is sheet-tied with frozen conjugate. Reproduce:

```
pytesttests/test_potential.pytests/test_driven.py-q
pytesttests/test_continuum.pytests/test_falsification.py-q
```

Interference apparatus: two-path PASS on open square, square torus, J2, and the J2 quotient; FAIL on tri/hex; rewires fail the phase rung; MZ corridor LAW-level after a window erratum (VAC-0F completion: square class, not J2-only; H/I closed below, VAC-0 FINAL MIXED). Reproduce:

```
pytesttests/test_slit.pytests/test_coherence.py-q
pytesttests/test_tunnel.py-q
```

Operational geometry and quotient. OBS-0R meets six rulers on J2 (balls/Hausdorff gap 0, diffusion/spectral gap 0.0005, wave gap 0.134 finite-size residue, static POT ruler joins; all vs the square control at $L = 128$). OBS-1 blinds the observer (no graph, coordinates, or dimension target): $d_O = 2.02$ unprompted, $d^* = 2$ train and test, 8% distance match vs the quotient ($2\times$ closer than microscopic), 6% local charts, $>95\%$ locality, sheet contrast < 0.03 ; expanders $d^* = 3$. $M_O(\text{J2}) \approx \text{J2/sheet}$; a static-channel floor blocks the full three-way cross-probe at $L = 128$. QUOT-0 gives the mechanism (QUOT0-OPERATIONAL): the symmetric sheet sector transports while the antisymmetric sector is exactly dead under $H = -A$ (MALUS-0 M0-NUL); symmetric replay returns the quotient (d drift 0.022), antisymmetric measures no geometry, a bilayer control stays two worlds (contrast 0.61), staggered perturbation induces only $O(\varepsilon^2)$ remote signal. The DERIVED rung stays open (four design-error sub-bars, autopsied, never retuned). Reproduce: `pytesttests/test_obs0.pytests/test_obs0r.pytests/test_obs1.pytests/test_malus.pytests/test_quot.py-q`.

Backreaction ontology and accounting. B couples to connectivity three ways: the derived conjugacy, BR-0 energetic selectivity (BR0-D-SELECTIVE*: zero field exactly flat under sampled moves, excitation opens channels — re-read as the no-information limit, not the vacuum), BR-2 quadrature ($B \sim \cos$, $J \sim \sin$), and the BR-2.5 sum map ($\psi_{[uv]} = \psi_u + \psi_v$, $\Delta\|\psi\|^2 = +2B_{uv}$).

Admitted primitive: local $u-v \leftrightarrow [uv]$ (BR25-ONTOLOGY); remote relocation demoted to di-agnostic; splits many-to-one (powers of two + amplitude degeneracy; loss $|a-b|^2/2$ exact) with selection debt; BR1-FLAT files neutral quiescence. CONS-0 + BR-2.6 close the books conditionally (CONS0-PARTIAL, BR26-ACCOUNTED): triangle-free cycle rank tracks the contraction count, the ledger closes on $B = B_*(c)$ (six families, 10^{-13}), the ΔE formula needs more than B ; no linear conserved combination, no graph reservoir, split conservation never selects. BR-2.7 derives the no-mode verdict (BR27-NO-MODE): downhill scans do not fire, spectral radius exactly 1, no deformation coordinate in a binary graph — one further primitive postulate required; strong stop on rate shopping. Reproduce:

```
pytesttests/test_backreaction.pytests/test_phase.py-q
pytesttests/test_rigidity.pytests/test_contraction.py-q
pytesttests/test_conservation.pytests/test_accounting.py-q
pytesttests/test_stability.py-q
```

Dynamics completion (debt-filing). U0: three contraction-only laws pass the structural gates but splits are unrealized and ties are theorematic at zero field (U0-INCOMPLETE). TIME-0: exact census (143 classes, 102245 pairs) gives unique-history 0.051 below the 0.2 bar with the $T=2$ artifact dissolved (TIME0-NUL; reversibility intact). RAND-0: coherent stochastic apparatus, underived measure (RAND0-MEASURE-DEBT: orbit vs micro uniforms disagree, vacuum not quiescent at one-half). MEASURE-0: 544 cells, all four frozen debt-reasons trigger, CLOSED reachable but not reached (MEASURE0-DEBT). Synthesis: admissible histories known, weight $\mu(\Gamma)$ missing. SYM-0 settles the counting list (SYM0-CLOSED, 1973 cells, 8/8 gates): $X_{\text{red}} = X/(\text{relabel} \times U(1))$ with d_{FS} ; no Born rule. Reproduce: `pytesttests/test_u0.pytests/test_time0.pytests/test_rand0.pytests/test_measure0.pytests/test_sym0.py-q`.

Vacuum family. VAC-FIELD-0 (frozen (J2, $H=-A$), spectrum/symmetry-fixed candidates): VPLUS ($E=-8$, flat ledger, P+, propagating, quotient-visible), VPI ($E=+8$, one-sided ledger, P+), VMINUS ($E=0$, symmetric ledger, P-, joint but decoupled) reach the full JOINT ladder; $\psi = 0$ stops at BACKGROUND ($B_{\text{max}} = 0$, phase undefined everywhere) — the no-information limit, not the vacuum (VACFIELD0-JOINT). Perturbations identical on all four (speed 1.9204, bitwise $\delta\psi$). VAC-EXC-0: $\delta\psi(t) = U(t)\delta 0$ exact ($\leq 1.2 \times 10^{-13}$), bitwise-identical across all vacua (VACEXC0-COMplete, 241 tasks). VAC-COMP-0: even L gives two isolated extremal rays plus a hidden real RP^1 JOINT circle (VMINUS-VSTAG span, $B=0$ BACKGROUND points) $\times$ amplitude ray (10^{-3} – 10^3), $\pi_0 = 3$; odd L keeps VPLUS + VMINUS only ($\pi_0 = 2$); generic eigenstates excluded; mixed beats at $dE = 16/8/8$ (VACCOMP0-COMplete). VAC-SELECT-0: predicted refusal — MEASURE gate unready, 0/23 headline stages ran (VACSEL0-NOMEASURE). Reproduce:

```
pytesttests/test_vacfield.pytests/test_vacexc.py-q
pytesttests/test_vaccomp.pytests/test_vacselect.py-q
```

Hidden sector, linearity, response, zeros, gravity null. HIDDEN-0: hidden transformations distinguish locally (5–6 orders above bar) with remotes at $\leq 5.2 \times 10^{-15}$ and POT remote exactly 0.0 (HIDDEN0-SEPARATED, 279/279). HIDDEN-BR: equal-energy, wave+POT-blind matched states carry different B (different dE/dA), with 1346 strictly opposite-sign virtual contraction ledgers; a zero-energy pure-hidden state carries a nontrivial ledger — all virtual, no graph operation executed (HBR0-SIGNREV, 214/214). FIELD-0: 57 collision cells hold $I = 0$ exactly (FIELD0-LINEAR) through order-50–80 false accelerations and a forceful-looking static arrangement (FIELD0-APPARENT); a future force must show $I > 0$. RESPONSE-0: field front $v = 7.95$ ($v/8 = 0.993$), quadratic fronts $v \approx 5.94$, laws $|\delta\psi| \sim r^{-0.50}$, $\delta\rho \sim r^{-1.00}$, $\delta J \sim r^{-0.95}$, bipartite B -blindness pinned with the two-sublattice sharpening (RESPONSE0-KERNEL-BANKED, 45 cells).

BG-RESP-0: $\chi_+ \neq \chi_\pi \neq \chi_-$ ($\sqrt{88}$ pairwise, rank $2N-1$, null exactly global phase), $\chi_{\text{ZERO}} = 0$ (BGRESP0-COMPLETE, 77 tasks; susceptibility is math response only). ZERO-0: exact zeros are interference-nodal (1588 events at phase π) or eigen-nodal (only persistent class); generic states give zero certified zeros in 1900+ rows; winding slips via bond phase-slip at $|\Delta\theta| \approx \pi$. GRAV-0: no tested strictly local graph-only update both preserves J2 and carries a far disturbance — the carrier must come from coupled (G, ψ) dynamics; the L1 weak-field interface is not retracted. Reproduce:

```
pytesttests/test_hidden.pytests/test_hiddenbr.py-q
pytesttests/test_field0.pytests/test_response.py-q
pytesttests/test_bgresp.pytests/test_zero.py-q
pytesttests/test_zero0_campaign.pytests/test_grav0.py-q
```

VAC-0 completion (substrate classes). The held-out H/I rungs close the campaign as VAC-0 FINAL MIXED: per phenomenon, generic laws (A, D, H-core existence 16/18 on 27/27, I finite-range label 26/27, J algebra 27/27 exact, MZ at LAW level) or class members (F square class, E 2D-ordered class, H turn-on 15/27 broad class, G square-grade class, J-useful 13/27 class), with I ξ scale family-clustered (CLASS-structured). Nothing requires uniquely J2; degree-preserving rewiring destroys F/H/G/DE-validity (degree-only vacuum excluded) while alternatives survive everywhere (square/quotient, +tri/rr3, +hex/ring). Substrate uniqueness dissolves into per-phenomenon classes. Reproduce: `pytesttests/test_vac0_battery.py-q`.

Split anatomy and rewire null. SPLIT-0 decomposes every tested inverse into forced-plus-residual ($M + \xi \leftrightarrow X$, 824/824 roundtrips, minimality 76/76 with two-way ablation witnesses): the covariant residual $\xi = (\text{cover}, d)$ is exactly what the merged state omits. The full inverse is never a singleton (continuous fiber, d_{cont} 1–2); the halves-restricted inverse is deterministic exactly on the 4 isolated-node cells; reverse support holds exactly on the halves subset (36 rev / 24 graph-only / 0 one-way), correcting 3 cells against the MEASURE-0A signature proxy (J-multiset edge-orientation artifact, pinned). No measure derived (SPLIT0-MIXED). REWIRE-0 closes the selector question for the local degree-preserving move (REWIRE0-DEGENERATE, 24/24 J2-L4 states DEGENERATE-or-ABSENT under all six exact principles, scale-persistent L4 $\rightarrow$ L28 with n_{phys} 9792/11291/6756; exact-energy selection additionally ulp-fragile, 1389 violations filed as representation-robustness null): no earned local, covariant, zero-parameter rule selects a rewire on any nontrivial state. Reproduce:

```
pytesttests/test_split0.py-q
pytesttests/test_rewire0.py-q
```

Merge update, reservoir account, trigger census. MERGE-0 characterises selected-edge contraction as a unique covariant deterministic primitive ($R \times U(1)$ exact on all 325 events incl. 87 annihilations, 8 substrates) with exact ledger ($dQ = 2B$, $dE = P1 + P2$ with $P3 = P4 = 0$, support $2 + n_{\text{cross}}$) — while no existing variable closes the energy reservoir on any tested class (0/80 coefficient tuples; MERGE0-DETERMINISTIC + MERGE0-ACCOUNT-DEBT, 38/38). HIDDEN-BR SIGNREV survives execution (2/24 on-support opposite-sign executed ledgers); VPI all- $B < 0$ yet all- $dE < 0$; favourable orderings on 19/30 scans, still no firing rule. RESERVOIR-0 identifies the missing account $R_{\text{merge}} = -(\Delta E_\psi + \Delta E_G)$ as an exact nontrivial zero-parameter function of the lost inverse information $R(d) = A + |d|^2/2 + \text{Re}(\bar{d}W)$ to $\leq 10^{-9}$ on 26992 fiber rows, both ξ legs varying, swap bitwise, $R_{\text{split}} = -R_{\text{merge}}$, one-neighbourhood-local, disjoint-additive (8.9×10^{-16}), confluent (RES0-XI, 68/68) — and R never vanishes on the halves subset (964 rows, min 0.167) though no field information is lost there: energy debt $\neq$ information debt. TRIGGER-0 censuses the trigger side under strict no-shopping (TRIGGER0-CONDITION, 75/77): 19 of 21 earned exact local predicates survive as CANDIDATE-CONDITION on 257668 edges (BRIDGE NONLOCAL,

BAL_R2 VACUOUS at 0/257668; BAL_R1 on exactly 6 hidden-dipole bonds; BJ_ZERO $\iff$ ZERO_MIN exact), with zero dynamical implications in the structural audit. No TRIGGER-1 shopping follows (prereg binding). Reproduce:

```
pytesttests/test_merge0.py-q
pytesttests/test_reservoir0.py-q
pytesttests/test_trigger0.py-q
```

Fiber measure debt and probability-free books. FIBER-0 asks whether earned physics constrains a normalized measure on the SPLIT-0 inverse fiber (FIBER0-DEBT, 22/22): two inequivalent normalized closed-form rivals satisfy every earned constraint (quotient, swap, transport, locality, collapse, ledgers, energy, vacuum, hidden, factorization, scheduler, history), differing with $TV > 0$ on 79 cells; DERIVED was reachable (residual 0) and did not occur (residual $520 = 206 + 168$ inter-orbit $+ 83$ radial $+ 63$ angular). Cover measure unique only on the $d = 0$ cells (4/76); factorization a choice (correlated rival TV 0.25); TIME0-NULL survives fiber weights; zero fitted params (AST firewall clean). Six sub-debts filed by name. INFO-0 closes the probability-free books exactly (INFO0-MATCHED, 19/19 HARD, 259 cells, 0 failures): canonical pred == succ 143/143 (mirror theorem, no SEPARATED), timed-skeleton identity, schedulers all $m!$ exact, hidden books green, banked TIME-0 recomputed exactly; the 16/44 labeled gauge audit is descriptive (labels are redundancy; the quotient restores symmetry). No $-\sum p \log p$ anywhere; the weight is still missing. Reproduce:

```
pytesttests/test_fiber0.py-q
pytesttests/test_info0.py-q
```

Vacuum domains, textures, stability. VAC-DOMAIN-0: every fixed-geometry interface between disconnected JOINT components (VPLUS/VPI, VPLUS/hidden $\times 4$, VPI/hidden $\times 4$, both orientations, $L = 28$) emits ballistic $|\delta\rho|$ fronts ($v \approx 5\text{--}6.1$, R^2 0.89–0.92, reach 4–6 cells) while the S-step persists (0.999) and both bulk plateaus hold (drift 0.0); hidden-hidden joins exactly stationary (D-FLAT); P_− residue frozen at weight 1/2; spectral superposition exact with witness $I = 0$ (VACDOMAIN-RADIATIVE, 7/7, 204 tasks). No wall particle or vacuum preference (firewall held). VAC-TEXTURE-0: hidden JOINT orientation can vary spatially staying exactly vacuum-like (P_− E₀, $w_{\text{sym}} = 0.0$, $\|H\psi\| = 0.0$, $E = 0.0$ bitwise, no P₊, frozen) while gradients stay relationally real (local $D = 6.4 \times 10^{-4}$, coarse 1.3×10^{-3} , ledger separation, B -scaling slope $+0.173$, $Q = a^2$ exact) and dynamically void (symmetric amplitude 0.0, FIELD-0 witness 1.6×10^{-17} , packets exact-split and ballistic at $v = 1.92$; VACTEXTURE-GRADIENT, 14/14, 28 specs). VAC-STAB-0: small protected perturbations stay small on all six backgrounds to $T = 4000$ (max C-ratio 3.5 vs trigger 50, propagating-seed 0.499 below own initial 0.704, positive protection margins, 303 cross-background wraps, frac-sup-B slope 2.0000 exact over $10^{-3}\text{--}10^3$; VACSTAB0-ROBUST, 9/9, 180 tasks, 0 frag cells). No structural claim, none ranked. Reproduce:

```
pytesttests/test_vacdomain.py-q
pytesttests/test_vactexture.py-q
pytesttests/test_vacstab.py-q
```

Sources as boundary data. SOURCE-0: POT's stationary field IS the driven counterpart of the RESPONSE kernel (K1 dev 0.024–0.037 within the 0.1 bar, K2 exact to 6.6×10^{-12}). A persistent source is boundary data $s(t)$ on $\delta\psi$ in the vacuum frame; the carrier is vacuum-independent with per-vacuum response via banked χ (42/42). Switch-ON always radiates ($v = 5.5\text{--}5.6$); the falsified blanket prediction was release fronts on AMP legs — AMP/VPLUS release is silent by theorem (steady state $c \cdot u_+$ to 7.4×10^{-15}) and AMP/VMINUS release is beating-dominated

and frontless (SOURCE0-INCOMPLETE, 9/10, 74/74 records; INCOMPLETE = falsified frozen prediction with complete data). Rule: switch-OFF radiates only on mismatch with free evolution. $R_G \rightarrow dG$ still blocked by MEASURE0-DEBT. Reproduce: `pytesttests/test_source0.py-q`.

Debts (by name; D14/D15 taken). Substrate classes (VAC-0 Final MIXED: F mechanism, VAC-0Q held-out battery, unified H-turnon property); vacuum-field member (needs the history measure first); history measure $\mu(\Gamma)$ (reversible, both readings; fiber rivals differ on 79 cells); structural kinetics (firing postulate; 19 conditions, 0 implications); split information (fiber measure, 520 filed dof); neutral quiescence; matter (beat $I = 0$); gravity carrier (coupled dynamics). Blocked until then: complete graph dynamics, formation under it, matter, the carrier, decay, and force phenomenology. Full ladder: `docs/scaffolding-history.md`; model statement: `docs/model.md` §10.

Table 1: Every number with a status: input = assumed, fit = calibrated, measured = code output, derived = follows from premises.

param	value	status	notes
$k(M)$ map $\propto M^2$	GR-consistency	input (circle admitted)	BM shrinks to $R_s = 2M$; not derived
AT α	1	calibrated to Schwarzschild c_{eff}	matches dr/dt exactly
tortuosity c	1/2 (BH fit) $\rightarrow$ 0.44–0.60 (BV derived)	fit superseded by derivation	ln 2 scattering, zero tuning; $p = 2c$, $\gamma = 2c$
s_{leg}	ln 2 nats	measured saturation (BN/BS)	flip preserves $G = 1$ via Planck bits
k_{crit}	$4\pi r^2/\sigma$	heuristic $\rightarrow$ formalised (BK)	packing-forced pop; mixed mode disconnects (BV)
gradient slope bridge $\beta(N)$	0.015 1.5@300; 1.28@600; 1.24@1020; 0.99@4k; 0.87@8k; 0.74@16k	fitted, labelled recalibrated per N	$p_{\text{adj}}(s) = 0.85 + 0.015s$ log-linear over 6 points; derivation queued
$w = 1.953$	solved from cancellation	fitted (solved, not tuned)	$c_{\text{tot}} = c_1 + wc_2 = 4.8695$ at $p = 0.92$
shed $e + \epsilon$	$0.5 \rightarrow 0.416 + 10\%$	calibrated once on AT2017gfo	M_{ej} norm-independent
shed $\epsilon(M, a)$	0.0168 universal (Eq. 2 main)	calibrated + extrapolated (D8)	mass/ratio independence underived; BBH extension tested, not fitted
upper-gap null	no feature at $44 M_\odot$	negative prediction (measured smooth)	$k/\text{spin}/\text{Love}/\text{congestion}$ all smooth; edge left to stellar physics
GW190814	$P(\text{detect}) \approx 0.68\text{--}0.88$	tension, POSSIS queued (D7)	non-detection at $p \sim 0.12\text{--}0.32$; epoch audit in massgaps
vacuum P0'	isostatic 2D, $\langle z \rangle = 4$	postulated (v0.6)	BH = max-tension extreme; $d_I = 2$ predicted; M_O reconstruction
d_{eff} protocol	$d \ln V / d \ln r$ bracket	measured (25 tests)	shortest below + diffusion above on lattices; shells fail (S11)
D10 tension map	GR dip/overshoot/ $\rightarrow 3^+$	GR computed; graph open	bare shortest-path rejected (inverted); D11 tail tension-dep. ($\chi_1 1/r^2$, $\chi_2 1/r$, χ_5 crossover $q \rightarrow 1.1$); κ -profile closed (disk, mild-disk confirm)
D12 reconstruction	$\forall M_O \in \mathcal{A}_{\text{macro}}$ same IR	open program	P4 audit 1,2,3,5 pass, 4 partial; vantage axis filed; static depth neg. (vol+MDS); MDS bar per-substrate (del bowl)
D13 causal order	D13.0 control measured; stages 1–4 open	static control measured (S11)	V_G , $8r+4$ law; family (tri/hex/Del/Lloyd/Gab/kNN/medial +, gated-wall/rewired -); sweep: $N^* O(1)$ range, $f^* \rightarrow 0$ hypoth.; spectral d_s band; twins C3
U -rule search	locality protection by U	third pass: blind- U + substrate census	blind- U + admissibility; 7-substrate coord. verdicts; self-cal. census
D14 pricing origin	hierarchy gain ~ 350	debt by elimination (measured)	basin + feedback OK; formation: DARK, NO-coex (N^*), SSB-1, St-0 NULL
J2 micro/macro probe	quotient = square lattice	measured; working vacuum (v5.4)	quotient $4r$ exact; bipartite; perturb. family-typical; orient. absent (32 runs)

Table 2: Honesty ledger, continued: relational field program (S12). Same status words.

param	value	status	notes
field postulates	$\psi=r+is$ per node, $H=-A$	postulated FP1 / working law FP2	one-way $G \rightarrow \psi$ map; not derived (S12)
field anatomy	B/J conjugacy	derived LAW (all simple graphs)	VAC-0A A1–A6 on hostile graphs; quadrature $B \sim \cos$, $J/2 \sim \sin$
substrate class	$\partial E/\partial A = -2B$ per-phenomenon	measured (VAC-0 LAW/CLASS split)	nothing requires uniquely J2; degree-only excluded (S12)
EM reading	ψ is ordinary electromag- netism	rejected (EM1- FALSIFIED)	range saturates $\xi \approx 0.53$; 1 scalar mode; no gauge redundancy; drift $v \cdot q$
geometry change	contraction/split $v \leftrightarrow [uv]$	$u-$ ontology admitted; firing absent	ledger exact; BR27-NO-MODE; SPLIT0 ξ residual; MERGE0 update unique; RES0 $R=f(\xi)$; TRIGGER0 19 cond. 0 imply; REWIRE0 no selector (S12)
dynamics completion	U_0 /TIME-0/RAND- 0/MEASURE-0/FIBER- 0/INFO-0	all file the measure debt; books exact	no unique $\mu(\Gamma)$ forced; SYM-0 counting X_{red} ; fiber rivals differ, 520 dof; INFO0 pred==succ 143/143 (S12)
observer quotient	$M_O(J2) \approx J2/\text{sheet}$, $d_O=2.02$	measured blind; mechanism opera- tional	6 rulers meet; symmetric sector transports; derived rung open (S12)
vacuum family	VPLUS/VPI/VMINUS joint; $\psi=0$ no-info limit	measured family; selection refused	carrier background-free; manifold censused; VACSEL0-NOMEASURE; domains RADIANATIVE; textures GRADIENT; stab. to $T=4000$ (S12)
hidden sector	dead sector stores local info, reverses response	measured (virtual ledgers)	HIDDEN0 separated; HBR0 1346 sign-reversed edges, all virtual (S12)
linearity/response	$I=0$ null; kernel + per- vacuum χ banked	measured (57 + 45 + 77 cells)	fronts 7.95/5.94; $\chi_+ \neq \chi_\pi \neq \chi_-$, $\chi_{\text{ZERO}}=0$ (S12)
sources	persistent $s(t) = \text{bound-}$ ary data on $\delta\psi$	measured; blanket release falsified	K1 dev 0.024–0.037; switch-OFF radiates only on mismatch (S12)
$\kappa \rightarrow c_2$ map	$c_2 = p(2p-1)$	ansatz (open deriva- tion)	power-law vs $1/r^2$ disagree cross-applied
evap. isometry V_k	qubit V_t done (D2); graph small- N ED done	large- N open (S3)	$V_t^\dagger V_t = I$ tested; graphvk all:all < 0.25 bits vs chain > 0.4 at $N=8$ (D1)
Kerr M_2 , $g_{t\phi}$, ISCO, QNM	area only	not derived (S3)	$A(M, a)$ assumed; quadrupole/QNM open; GW250114 QNM + GW241011 δ_Q future wires
echo R	10^{-80} ampl., 10^{-160} en- ergy	dimensional esti- mate	$\propto \omega^2$ lattice scattering, not leg S-matrix; ~ 158 orders margin

 Table 3: Gradient-shell p measurements: β recalibrated per N (log-linear over 6 points). All 80-graph production rows use exact distances; errors are per-graph std with SEM quoted in text.

N	shells $\times$ per-shell	β	measured p	backend
300	10×30	1.50	0.94 ± 0.06 (4 graphs)	NetworkX exact
600	10×60	1.28	0.96 ± 0.06	NetworkX exact
1020	10×102	1.24	0.913 ± 0.049 , SEM 0.0055, stacked 0.911 (80 graphs)	Floyd+LP exact
4000	CSR-direct	0.99	0.9315 ± 0.0032 (80 graphs)	CPU + Sinkhorn OR
8000	CSR-direct	0.87	0.9382 ± 0.0030 (80 graphs)	pod + Sinkhorn OR
16000	CSR-direct	0.74	0.9137 ± 0.0022 (80 graphs)	GPU farm + torch OR

Table 4: PPN status: only γ is claimed. β is coordinate-confused and not quoted.

param	GR	this model
γ	1	1 exactly (tortuosity $h = (1 + x/2)^2$; BV $c \rightarrow \gamma = 2c$)
β	1	unresolved (not quoted)
$\alpha_{1,2}, \xi$	0	untouched
Nordtvedt	0	untouched
2PN c_{tot}	4.8695	4.8693 at $p = 0.92$ (J0737 0.1σ)

Table 5: What exists today: no qualifying test yet. Only multi-detector, $< 100 \text{ deg}^2$ gap events count.

event	masses	distance	localisation / EM	verdict
GW190814	$23.2 + 2.6$ ($q \sim 0.11$)	241^{+41}_{-45} Mpc	23 deg^2 (3-det.); CFHT g 22.8@1.7d 65.5%, GROWTH i 6 epochs to 98%	tension: $P(\text{detect}) \approx 0.68$ – 0.88, $p \sim 0.12$; not excluded
GW230529	$2.5\text{--}4.5 + 1.2\text{--}2.0$	~ 197 Mpc	24,200 deg^2 (single-det.); ZTF 7% to $g = 21.1$	not a test: 93% unsearched; $m_g \sim 21.2$ at depth edge
S250206dm	mgBH candidate	—	no counterpart (per notes)	unverified: check GCNs before citing