

Compact Objects as Almost-Perfect All:All Entanglement Graphs: From fast scrambling and weak-field gravity to a testable gap-kilonova prediction

Philippe Leblond

leblond.philippe@gmail.com

Draft v5.0 — main text (companion supplement with methods; 9pp preprint $\approx$ 6pp journal two-column)

`python scripts/generate_figures.py` reproduces every figure; 356 tests

September 27, 2026

Abstract

We study a phenomenological model in which spacetime connectivity is an entanglement graph and a black-hole interior is an *almost-perfect* all:all (complete) subgraph. Interior edges cost no exterior space; each of k exterior legs costs $4 \ln 2 \approx 2.77$ Planck patches of horizon area, so $A(k) = 4 \ln 2 \cdot k l_p^2$ independent of interior size N . From this wiring picture we recover fast scrambling ($t_* \sim \log N$), the Bekenstein–Hawking area law, the exact Page curve with Haar-typical fluctuations, island/QES takeover, Kerr thermodynamics, and Hayden–Preskill mirror recovery. The same leg network yields weak-field gravity to first post-Newtonian order: Newton’s $1/r^2$, Kepler orbits, textbook redshifts, full-strength light bending $4M/b$, Cassini-grade Shapiro delay, $\gamma = 1$ exactly, and Mercury’s $43''/\text{cy}$ by direct geodesic integration. Lattice hopping gives quadratic-only dispersion ($E_{QG,1} = \infty$), safe from Fermi bounds by $\sim 10^8$. At second post-Newtonian order the double pulsar J0737 is matched to 0.1σ via a measured radial exponent $p = 0.913 \pm 0.049$ (80 graphs, $N = 1020$, exact; confirmed at $N = 4000/8000/16000$). This promotes the toy to a testable model of *all* compact objects: pulsars, mass-gap objects, and black holes as one $k \propto M^2$ family with no neutron-matter phase. Calibrated once on AT2017gfo ($0.047 M_\odot$ shed), the model predicts mass-gap mergers $2.5\text{--}5 M_\odot$ are kilonova-bright at $\sim 1/\text{yr}$ in O5 versus $\leq 0.3/\text{yr}$ in the standard picture. Ten well-localized gap non-detections below 200 Mpc kill the proposal; one bright gap kilonova kills neutron stars instead. All claims ship with runnable code, tests, and pre-registered falsifiers.

Roadmap. §1 motivates the model and states how it differs from neighbours. §2 defines the three claims precisely. §3 summarises gravity to 1PN with a 2PN preview. §4 covers UV dispersion and quantum information. §5 unifies compact objects and states the gap-kilonova prediction. §6 lists falsifiers and concludes. The supplement gives methods, the honesty ledger (postulated vs. derived vs. fitted), and the module map. The v4.1 living document (1758 lines, 68 appendices, 77 figures) remains archived as the extended technical record.

1 Introduction

Black holes confront us with three facts that sit uneasily together: they scramble information as fast as any system allowed [1], their entropy scales as area rather than volume [4, 5], and their evaporation — if unitary — must release that information in a Page curve with island/QES turnover [6, 7, 8]. Separately, gravitational-wave astronomy has populated corners of the compact-object mass function the standard picture did not expect: objects in the $2.5\text{--}5 M_\odot$ mass gap (GW190814 at $2.6 M_\odot$ [21], GW230529 at $3.6 M_\odot$ [15]), a central compact object below the minimum neutron-star mass (HESS J1731 at $0.77 M_\odot$ [16]), and pulsars pressing against the maximum Tolman–Oppenheimer–Volkoff mass (J0740 at $2.14 M_\odot$, J0952 at $2.35 M_\odot$). Each anomaly has its own candidate excuse; none was predicted.

This paper explores one idea that addresses both sets jointly: *maximal entanglement is zero-length connectivity*. In the ER=EPR and “spacetime from entanglement” reading [2, 3, 29], a black-hole interior is a region where interior mutual connectivity is so dense that interior distance collapses, while horizon size counts only the *exterior* wiring that must be embedded through ambient space. A perfect all:all graph with zero exterior budget would pinch off entirely (a baby universe); a black hole is the almost-perfect corner with small but nonzero exterior budget. A sufficiently small hole stays pointlike until its exterior budget forces an empty routing surface — a horizon — into existence.

We formalise this as three claims (§2), each implemented as runnable code in `src/bh_graph/`. The model is phenomenological, not a UV-complete quantum gravity: it has no Hamiltonian derivation of the mass map, no Lorentz-invariant dynamics, and several labelled derivation debts (supplement §1). Its value is that it *derives* a large number of textbook numbers from a small number of inputs, fails loudly on pre-registered wires when it is wrong, and makes one sharp sky prediction that the current observing run can kill or confirm.

How this differs from neighbours. Fuzzball [36] and gravastar [37] proposals also remove the singularity but keep compact-star matter phases and do not generically predict bright gap kilonovae; quark/strange-star models [38, 26] populate the gap with a new equation of state (EOS) rather than removing the EOS entirely. Polymer and loop-inspired holes modify the near-horizon metric at a level constrained by EHT [32] and γ [31, 30]; our 1PN sector is exactly GR ($\gamma = 1$) with deviations appearing only at 2PN below current VLBI/astrometry. Linearly-dispersive “spacetime foam” models [39] are already excluded by Fermi GRB 090510 [17]; our lattice implies quadratic-only dispersion by $k \leftrightarrow -k$ symmetry, evading that bound by construction. Table 1 (survival ledger) and Fig. 1 make the comparison explicit: among the surveyed alternatives, only the present model passes the Fermi, LHC-recast, EHT/ γ , and gap/HESS battery simultaneously while predicting a bright-gap rate distinct from the standard one.

Fig V5-1 — Survival matrix: only all:all passes the full battery + predicts a distinct gap-KN rate

	Fermi GRB 090510	LHC thermal null	EHT shadow + gamma	Gap 2.6/3.6 +HESS 0.77	Gap-KN rate distinct?
Linear LIV foam	FAIL	—	—	—	FAIL
TeV thermal BH	—	FAIL	—	—	FAIL
Fuzzball / polymer	—	—	FAIL	—	FAIL
GR + NS EOS	PASS	PASS	PASS	FAIL	FAIL
All:All (this work)	PASS	PASS	PASS	PASS	PASS

Figure 1: Survival matrix (visualisation of Table 1; no new numbers). `scripts/generate_v5_figs.py`.

2 The model: three claims


Let the interior be a graph G_N on N nodes. Exterior legs $k \ll N^2$ connect it to an ambient graph. Write $e_{\text{int}} \in [0, 1]$ for the interior entanglement fraction and e_{ext} for the exterior budget fraction.

Table 1: Survival ledger: headline result per sector and where it is implemented and tested.

sector	headline	status / modules
gravity	Newton, Kepler, redshift, $4M/b$, Shapiro, Mercury $42.99''$, $\gamma = 1$ exactly, J0737 0.1σ	pass (AS, AT, AU, BB, BH)
quantum info	$\log N$ vs N hierarchy, Page dip 0.72 bits, QES pop, mirror, $k_f > k_1 + k_2$ at 100%	derived (A, C, D, F, H, Q, W)
UV	quadratic-only LIV, $E_{QG,2} = \sqrt{8} E_P$, achromatic lensing 10^{-56} – 10^{-32}	nulls held (T, BB, BD, BF)
compact objects	gap 2.6/3.6, HESS 0.77, J0740/J0952, AT2017gfo, gap-KN $\sim 1/\text{yr}$	5 anomalies, one law (BU)
open	$p = 0.913 \pm 0.049$ matches GR; $c \approx 0.44$ – 0.60 derived (BV); $\kappa \rightarrow c_2$ map open	measured, map queued

Claim 1: all:all is no interior space (fast scrambling). The complete graph K_N has diameter exactly 1, mean distance 1, and spectral gap N at every N : every node is adjacent to every other (Fig. 2). A deterministic SI/operator-spread process covers K_N in 1 step for any N , versus $\sim N/2$ for a chain P_N , $\sim 2\sqrt{N}$ for a $\sqrt{N} \times \sqrt{N}$ grid, and $\sim \log N$ for a random 3-regular expander. Imposing finite speed — one 2-qubit interaction per qubit per step with random perfect matching (all:all) versus dimer covering (chain) and per-encounter success p — promotes the toy to a derived Sekino–Susskind law [1] $t_*(N) \approx \log_2 N / \log_2(1+p)$, exactly $\log_2 N$ at $p = 1$, versus ballistic $t_* \sim N/v$ on the chain (measured over 25 trials per N , $N = 8\dots 128$; chain already $> 5\times$ slower at $N = 64$). Modules `graphs`, `scrambling`, `circuits`, `otoc`, `krylov`, `syk`.

Fig 1 — Sec 1: fast scrambling hierarchy


 Figure 2: SI cover time and diameter: all:all stays at 1 while local baselines grow. Finite-speed circuits (supplement §3) promote this to $t_* \sim \log N$.

Claim 2: horizon area counts exterior legs. Postulate

$$A(k) = 4 \ln 2 \cdot k l_p^2, \quad R(k) = \sqrt{A/4\pi}, \quad (1)$$

so interior size N drops out: adding interior nodes without exterior legs tightens binding but buys no horizon area. The patch coefficient $4 \ln 2$ was fixed (BS) from the measured von Neumann leg entanglement $\eta_{vN} = \ln 2$ per leg with Planck-bits equipartition, closing the Jacobson chain at $G = 1$; legs saturate. Each exterior leg must be embedded through ambient geometry with Planck-density bandwidth, so k legs need k Planck patches — the Bekenstein–Hawking / LQG-puncture picture [4, 5, 10] in graph language. Mass enters only through the GR-consistency condition $k(M) = A/4 \ln 2 l_p^2 =$

$(4\pi/\ln 2)M^2/l_p^2 \propto M^2$ in Schwarzschild units (module `horizon.k_from_mass_schwarzschild`). We state plainly that this map is *assumed for consistency, not derived*: the reduction theorem (BM in the extended record; supplement §1) shrinks the circle to the single statement $R_s = 2M$, but the derivation of the mass–radius relation from wiring alone remains open (supplement §1). Monogamy bounds $e_{\text{int}} + e_{\text{ext}} \leq 1$ in the linear toy (Fig. 3), with the exact Coffman–Kundu–Wootters frontier [11] $\tau_{A|BE} - C_{AB}^2 - C_{AE}^2 \geq 0$ verified on a 25-point grid and lying strictly below the linear bound (e.g. $x + y = 0.70$ at $t = 0.3$). At $e_{\text{int}} \rightarrow 1$, $k \rightarrow 0$: the subgraph decouples as a closed graph (baby universe). Black holes live in the almost-perfect corner.




Figure 3: Monogamy frontier: black holes live at $e_{\text{int}} \lesssim 1$ with small nonzero k ; the $k \rightarrow 0$ limit is a decoupled baby universe.

Claim 3: micro-holes pop (point-to-horizon transition). A point region of radius r_{point} embeds k legs without a surface while Planck-density packing succeeds:

$$R_{\text{obs}}(k) = \begin{cases} r_{\text{point}} & k \leq k_{\text{crit}} \\ \sqrt{4 \ln 2 \cdot k l_p^2 / 4\pi} & k > k_{\text{crit}} \end{cases}, \quad k_{\text{crit}} = 4\pi r_{\text{point}}^2 / 4 \ln 2 l_p^2. \quad (2)$$

With an LQG-style minimal-area gap, area is 0 below threshold and $\geq A_{\text{min}}$ above: there is no 0.2-Planck-area hole. Growth $k(N) = k_0 + \alpha N$ stays flat at r_{point} then pops — the computed QES/island crossing $S_{\text{no}} = k s_{\text{leg}}$ versus $S_{\text{isl}} = k \ln 2 l_p^2 + \max(S_0 - k s_{\text{leg}}, 0)$ at $k_{\text{page}} = S_0 / (2s_{\text{leg}} - \ln 2 l_p^2)$, which exists iff $s_{\text{leg}} > \ln 2 l_p^2 / 2$ (an explicit falsifiable assumption; modules `micro`, `qes`). The horizon interior is mostly empty routing buffer. Prediction: micro-holes are *not* scaled-down Schwarzschild holes but point defects until a critical exterior budget.

3 Weak-field gravity to 1PN, with a 2PN preview

Newton and orbits. Microscopic lattice walks alone do *not* give Newton: persistent walks with hop-budget time dilation $\sqrt{1 - v^2}$ (linear $1 - v$ excluded by the 3-4-5 triangle test) and degree-biased hopping on a flux-conserved pileup $d \propto 1/r^2$ yield drift $\propto 1/r^3$ (measured slope ≈ -3), and every smooth weight rule stays cubic (no-go -2.99). The $1/r^2$ force law emerges from the combination with exterior-leg thermodynamics on leg screens (Verlinde equipartition [27], Bekenstein bound, Jacobson route [28] — all postulated): $F = M_1 M_2 / r^2$ exact to 10^{-9} in slope, with closed leapfrog orbits and

$T^2 \propto r^3$. Ollivier–Ricci curvature is radially negative in every tested configuration, attachment mode, and seed — the sign of attraction derived from pure flux with zero tuning.

Light: bending, Shapiro, and why isotropic dies. Fermat ray-tracing with $c_{\text{eff}} = 1 - x$ ($x = R_s/r$) gives the full first-order bending $4M/b$, not the Newtonian $2M/b$. The chromatic extension $n(r, \omega) = n_{\text{geom}}(r) n_{\text{disp}}(\omega)$ with phase-velocity factor $n_{\text{disp}} - 1 = (\omega/\omega_P)^2/24$ contributes only r -independently and factors out of the Born integral: fractional chromaticity $\sim 10^{-56}$ (optical) to $\sim 10^{-32}$ (10 TeV), achromatic by ~ 50 orders. Shapiro delay $\Delta T = \int (1/c_{\text{eff}} - 1) dl$ reproduces $R_s \ln(4r_1 r_2/b^2)$ to 5%, passing Cassini [31] with $\gamma = 1$ exactly. An isotropic reading would give $b_{\text{crit}} = 8M$, 54% above GR’s $3\sqrt{3}M$ and excluded by EHT [32] at $\sim 3.6\sigma$. It is the isotropic reading that dies, not the core: transverse propagation must be essentially unimpeded, as an all:all interior demands.

Mercury and the spatial sector. Purely temporal models give closed Newtonian ellipses (0 vs $43''/\text{cy}$): light never needed g_{rr} , orbits do. Radial tortuosity $dl = (1 + \sqrt{\chi}/2) dr$ with $\sqrt{\chi} = R_s/r$ gives $h = (1 + x/2)^2 \approx 1 + x$, hence $\gamma = 1$, $c_{\text{eff}} = \sqrt{f/h} = 1 - x$ matching the light sector to first order, and b_{crit} back to $3\sqrt{3}M$. Direct geodesic integration (ϕ -domain, complex-step) yields GR $42.99''/\text{cy}$ and model $42.99''/\text{cy}$; a flat- h hybrid gives $28.7''/\text{cy}$ (PPN 2/3). The second-order peel-off (ours–GR)/GR = $-0.75 M/a$ runs from 4.1% at $20M$ to 2×10^{-8} at Mercury. Status of the coefficient: $1/2$ was *fitted* to enforce $\gamma = 1$ in BH (unique coefficient, labelled fit); BV then *derived* $c \approx 0.44$ – 0.60 from $\ln 2$ line-defect scattering with zero tuning, predicting $p = 2c$ and $\gamma = 2c$ (supplement §2). The two determinations agree; the micro-derivation is now the adopted route with the fit preserved in history.

2PN preview (J0737). With $g_{rr} = (1 + U)^{2p}$ and $c_2(p) = p(2p - 1)$, the 2PN periastron weight $c_{\text{tot}} = c_1 + w c_2$ with $w = 1.953$ must equal the GR value 4.8695. The model’s $c_1 = 3.36$ (vs GR 1.94) is cancelled at $p = 0.92$. Gradient-shell Ollivier–Ricci measurements give $p = 0.913 \pm 0.049$ (80 graphs, $N = 1020$, exact, SEM 0.0055) and $0.93/0.94/0.91$ at $N = 4000/8000/16000$, landing J0737 [12] at 0.1σ (B1913 passes at 0.01σ [13]; supplement §4). The flat $p = 0.49$ alternative is -10σ dead. The quantitative $\kappa \rightarrow c_2$ map remains ansatz-dependent (power-law vs $1/r^2$ disagree cross-applied) and is the open micro-derivation, now labelled rather than hidden.

4 UV dispersion and quantum information

Dispersion: quadratic by symmetry. Tight-binding hopping $\omega = 2J|\sin ka/2|$ has group velocity $v_g = Ja \cos ka/2$, even in k : there is *no* linear term, so Fermi’s linear bound $E_{QG,1} > 9.3 \times 10^{19}$ GeV (and GRB 090510’s $E_{QG,1} > 1.2 E_P$ at 31 GeV, $z = 0.903$ [17]) is evaded by symmetry, with $E_{QG,1} = \infty$ predicted. The leading correction is $v(E) = c(1 - E^2/E_{QG,2}^2)$ with $E_{QG,2} = \sqrt{8} E_P \approx 3.4 \times 10^{19}$ GeV, safe by $\sim 10^8$ against Fermi’s quadratic bound 1.3×10^{11} GeV: a 10 GeV GRB photon over 3 Gpc delays by $\sim 10^{-20}$ s. Foamgrid FDTD confirms unbiased centroids with transmission deficit $\propto \omega^2$ ($16\% \rightarrow 4\% \rightarrow 0.2\%$ for $\lambda = 8 \rightarrow 32$ cells at $\varepsilon = 0.25$), bounding per-Planck-edge defect density to $\varepsilon \lesssim 10^{-3}$ (optical/Gpc) and 10^{-14} (TeV/Gpc) for independent edges. Caveats: regular-lattice result, near-horizon running open, scalar sector only (no birefringence claimed).

Scrambling, Page, QES, mirror. Circuits give $C(t) \sim e^{\lambda t}/N$ with $t^* = \log N/\lambda$ versus ballistic N/v ; QEC recovery error $\text{err}(k) = \min(1/2, 2^{N/2+1-k})$ reaches 99% at $k \geq N/2+1+\log_2 100$, sealing at $k \rightarrow 0$. Evaporation as leg surgery ($k \rightarrow k-1$ per step) reproduces the Page curve $S_{\text{rad}} = \min(t, N_{\text{eff}} - t)$ with the exact Page dip 0.72 bits and Haar-typical spread, identically whether N shrinks with k or

stays fixed — area tracks wiring alone. Kerr–Newman ($r_+ = M + \sqrt{M^2 - a^2 - Q^2}$, $A = 4\pi(r_+^2 + a^2)$) costs legs monotonically in spin at fixed M (extremal Kerr half, extremal Reissner–Nordström quarter the Schwarzschild budget, patch-independent ratios); the Kerr Page curve at $a_0 = 0.95$ peaks at 65% of Schwarzschild and turns over at $t/T = 0.73$ versus 0.54 because shedding J grows area early.

Gravitational-wave battery. All 32 confident GWTC-3 [22, 23] BBH medians satisfy $k_f > k_1 + k_2$ (median fractional creation 0.77 at 0.04 radiated; spin neglected but $a_f \sim 0.7$ costs $\sim 13\%$ against a 77% margin and cannot flip the inequality). GW150914 posteriors (8350 samples, spin-aware both ends) give $P(\Delta k > 0) = 100\%$, median 0.57: the area theorem in wiring language. Hardware anchors (Gärttner 2017 [18] all:all ions to $m = 8$ -body OTOC; Mi 2021 [19] Sycamore grid ballistic spread; Jafferis 2022 [20] sparse-SYK teleportation as a Hayden–Preskill mirror primitive [9, 40]) are all consistent, with a head-to-head prediction at $N = 53$ of $t^* \approx 7.9 \pm 0.7$ (all:all) versus 14.6 (grid), ratio $1.9\times$ ($9.9\times$ versus a 1D chain).

Nulls held. No LHC thermal black holes [33] ($k \approx 11$ –17 at 3–13 TeV versus $k_{\text{crit}} \approx 50$), no echoes ($R \sim 10^{-160}$), no EHT shadow shift (10^{-48} , 47 orders below sensitivity). Remnant dark matter is excluded on the record except a narrow ~ 0.4 -dex window at $\sim 4 \times 10^5$ g (photon bounds exceed by 40+ orders elsewhere; induced-GW bounds may close even this).

5 One family of compact objects

The standard picture needs a separate excuse for each of five facts. One law, $k = 1.51 \times 10^{77} (M/M_\odot)^2$ with no matter phases, plus two shedding numbers calibrated *once* on AT2017gfo [14], addresses all five:

Table 2: Five anomalies, one law. Shedding e : $0.5 \rightarrow 0.416$ plus 10% efficiency, fixed on #4.

#	anomaly	standard	this model
1	Gap: GW190814 2.6, GW230529 3.6 $M_\odot$	impossible (NS < 2.3, BH > 5)	continuous $k = 1.0$ – 2.0×10^{78} ; must flash
2	Light: HESS J1731 0.77 $M_\odot$	below min 1.17 $M_\odot$, cannot form	low- k tail 9×10^{76}
3	Heavy: J0740 2.14, J0952 2.35 $M_\odot$	EOS tension with NICER $R_{1.4}$	more legs 7 – 8.4×10^{77} , no EOS
4	AT2017gfo 0.05 $M_\odot$ blue+red	neutron-rich EOS in tension	leg-shedding 0.047 $M_\odot$, m_g 18.0 vs 17.5
5	Next gap merger (prediction)	dark (≤ 0.3 /yr O5)	bright ~ 1 /yr O5; 10 misses kill us

Leg-shedding works as follows: merger-released exterior legs hadronise neutron-rich with ejecta mass $M_{\text{ej}} = \Delta k m_{\text{leg}} \times \epsilon$, exactly independent of k -normalisation ($M_{\text{ej}} = \text{frac} \times M_{\text{tot}} \times \epsilon$), so the factor-3 k -rounding debate cannot move it. The $1.4 + 1.4$ case gives $0.047 M_\odot$ with $m_g \sim 18.0$ at 40 Mpc versus observed 17.5 (inside the ± 1 analytic tolerance; radiative-transfer refinement [35] queued); gap events are *brighter* at fixed distance (Table 3). The 1.1 – $2.3 M_\odot$ pulsar distribution is the low- k tail ($k \sim 10^{77}$) of the same family; the gap is where $e_{\text{int}} \sim e_{\text{ext}}$, not a boundary between kinds. If correct, stable nuclear matter ends at $\sim 10^{15}$ g/cc with no hyperon/quark branch [26] realised in nature, and r -process yields routed through AT2017gfo neutron ejecta are reinterpreted as shed-leg hadronisation — a blunt implication we state so no reader misses it, while noting that NICER radii [24, 25] and tidal deformabilities from routing stiffness are not yet derived (queued, supplement §4) and constitute the sharpest near-term test after the kilonova rate.

Table 3: Gap-kilonova brightness (g -band, analytic BC = 0; i -band not claimed). AT2017gfo anchor: predicted 18.0 vs observed 17.5 at 40 Mpc. Rubin single-visit $r = 24.5$ [34].

M_{tot}	M_{ej}	40 Mpc m_g	100 Mpc m_g	200 Mpc m_g
2.8 (GW170817-like)	0.047	18.0	20.0	21.5
5.0 (GW230529-like)	0.084	17.7	19.7	21.2
7.2 (3.6+3.6)	0.121	17.6	19.6	21.1




Figure 4: Leg-shedding ejecta vs total mass: gap mergers are kilonova-capable and brighter than GW170817 at fixed distance. Gap rate = 0 kills the proposal.

6 Falsifiers and conclusion

We state what kills the paper before concluding, with thresholds, current status, and timelines in Table 4.

Table 4: Pre-registered kill wires. “Now” = status at submission; “O5” = decisive by end of O5 unless stated.

wire	threshold	now	timeline
Gap kilonovae	10 qualifying gap mergers < 200 Mpc, < 100 deg ² , zero KN to $m < 24$	0 qualifying so far	O5 decisive
NICER radius	$R_{1.4}$ at 11–13 km, 5%, unreproducible by routing stiffness	stiffness underived	2–3 yr
Linear LIV	finite $E_{QG,1}$, or $\Delta t \propto E$, or 10^{-3} chromatic bending	Fermi already kills linear models	archival + CTA
2PN exponent	p outside 0.92 ± 0.056 at $N = 1024$ class, 80 graphs	0.913 ± 0.049 , passes $5\times$	held to $N=16000$
Lab quench	same-device grid→all:all t^* ratio < 1.3 (predicted 2–3 $\times$)	untested	when run

The gap-kilonova protocol (supplement §4, full trigger plan in `docs/observation-protocol.md`) requires multi-detector events only: single-detector non-detections such as GW230529 (24,200 deg², 7% ZTF coverage to $g = 21.1$ at ~ 197 Mpc, our $m_g \sim 21.2$ at the depth edge) are uninformative either way and do not count. Expected O5 yield is $\sim 1.05/\text{yr}$ (ours, 1.5 gap events/yr $\times$ 70% DECam-like) versus $\leq 0.3/\text{yr}$ standard. One bright gap kilonova kills neutron-star EOS models instead — the bet is symmetric and priced.

Conclusion. Starting from “all:all entanglement is zero-length connectivity” we have derived fast scrambling, the area law, the Page curve with islands, Kerr thermodynamics, Newtonian gravity with its sign, textbook redshifts, full-strength bending and Shapiro delay with $\gamma = 1$, Mercury’s 43″/cy, quadratic-only UV dispersion, and the J0737 2PN lock — all from one wiring picture with all failures and open maps kept on the ledger. The price is abandoning neutron matter above $\sim 10^{15}$ g/cc. The reward, if the sky cooperates, is the first quantum-gravity model with a kilonova prediction distinct

from astrophysics in the current run. If the compact-object unification fails, the title claim reverts with it and the surviving gravity/QI core stands alone; either way the wires above decide.

Methods summary. Full methods in the supplement. Gradient shells $p_{\text{adj}}(s) = 0.85 + 0.015s$, 8–10 shells, exact EMD on full neighbourhoods, deterministic bridge counts $n \propto r^{\beta(N)}$ with $\beta \approx 1.5/1.28/1.24$ at $N = 300/600/1020$ (log-linear to 0.74 at $N = 16000$); p from $|\kappa| \sim r^{-p}$ fits per graph plus stacked fits; 2PN via Iorio direct+total with $c_2 = p(2p - 1)$, $w = 1.953$, $R + \dot{\omega} \rightarrow M$ inversion; shedding $M_{\text{ej}}(M_{\text{tot}})$ with blue+red components, band mags, O5 yield arithmetic. Randomness seeded; tolerances recorded in-test.

Data and code availability. Code MIT (src/, scripts/, tests/, app.py); text/figures CC BY 4.0. Reproduce via:

```
pipinstall-e.
pytesttests/-q
pythonscripts/generate_figures.py
streamlitrunapp.py
```

GWOSC medians fetched live with bundled fallbacks; GW150914 posteriors cached (DOI 10.7935/82H3-HH23); PBHbounds curves vendored with attribution. No third-party figure data copied.

References

- [1] Y. Sekino and L. Susskind, JHEP 10, 065 (2008).
- [2] M. Van Raamsdonk, Gen. Rel. Grav. 42, 2323 (2010).
- [3] J. Maldacena and L. Susskind, Fortsch. Phys. 61, 781 (2013).
- [4] J. D. Bekenstein, Phys. Rev. D 7, 2333 (1973).
- [5] S. W. Hawking, Commun. Math. Phys. 43, 199 (1975).
- [6] D. N. Page, Phys. Rev. Lett. 71, 3743 (1993).
- [7] G. Penington, JHEP 09, 002 (2019).
- [8] A. Almheiri et al., JHEP 12, 063 (2019).
- [9] P. Hayden and J. Preskill, JHEP 09, 120 (2007).
- [10] C. Rovelli and L. Smolin, Nucl. Phys. B 442, 593 (1995).
- [11] V. Coffman, J. Kundu, and W. K. Wootters, Phys. Rev. A 61, 052306 (2000).
- [12] M. Kramer et al., Science 314, 97 (2006); updated $\dot{\omega} = 16.899323(13)$ deg/yr.
- [13] J. M. Weisberg and Y. Huang, ApJ 829, 55 (2016).
- [14] B. P. Abbott et al. (LVK), ApJL 848, L12 (2017); Villar et al., ApJL 851, L21 (2017).
- [15] LVK, P2300352 (2024); S. Kunnumkai et al., arXiv:2409.10651.
- [16] V. Doroshenko et al., Nature Astron. 6, 1444 (2022).

- [17] A. A. Abdo et al. (Fermi), *Nature* 462, 331 (2009); GRB 090510 $E_{QG,1} > 1.2 E_P$.
- [18] M. Gärttner et al., *Nature Phys.* 13, 781 (2017).
- [19] X. Mi et al. (Google), *Science* 374, 1479 (2021).
- [20] D. Jafferis et al., *Nature* 612, 51 (2022).
- [21] R. Abbott et al. (LVK), *ApJL* 896, L44 (2020).
- [22] R. Abbott et al. (LVK), arXiv:2111.03606 (2021).
- [23] R. Abbott et al. (LVK), arXiv:2112.06861 (2021), tests of GR with GWTC-3.
- [24] T. E. Riley et al., *ApJL* 887, L21 (2019); M. C. Miller et al., *ApJL* 887, L24 (2019).
- [25] T. E. Riley et al., *ApJL* 918, L27 (2021); M. C. Miller et al., *ApJL* 918, L28 (2021).
- [26] J. M. Lattimer and M. Prakash, *Science* 304, 536 (2004); *Phys. Rep.* 621, 127 (2016).
- [27] E. Verlinde, *JHEP* 04, 029 (2011).
- [28] T. Jacobson, *Phys. Rev. Lett.* 75, 1260 (1995).
- [29] S. Ryu and T. Takayanagi, *Phys. Rev. Lett.* 96, 181602 (2006).
- [30] C. M. Will, *Living Rev. Rel.* 17, 4 (2014), PPN and Solar-System tests.
- [31] B. Bertotti, L. Iess, and P. Tortora, *Nature* 425, 374 (2003), Cassini γ .
- [32] K. Akiyama et al. (EHT), *ApJL* 875, L1 (2019), M87* shadow.
- [33] CMS, *JHEP* 04, 047 (2023); ATLAS, *JHEP* 11, 169 (2022), BH and non-thermal searches.
- [34] Ž. Ivezić et al. (LSST), *ApJ* 873, 111 (2019).
- [35] M. Bulla, *MNRAS* 489, 5037 (2019), kilonova radiative transfer.
- [36] S. D. Mathur, *Fortsch. Phys.* 53, 793 (2005), fuzzball review.
- [37] P. O. Mazur and E. Mottola, *PNAS* 101, 9545 (2004), gravastars.
- [38] E. Witten, *Phys. Rev. D* 30, 272 (1984), strange/quark matter.
- [39] G. Amelino-Camelia, *Living Rev. Rel.* 16, 5 (2013), LIV review.
- [40] A. Kitaev, talks on SYK (2015); J. Maldacena and D. Stanford, *PRD* 94, 106002 (2016).