

# Black Holes as Almost-Perfect All:All Entanglement Graphs: Interior Collapse, Horizon Wiring, and the Micro-Hole Phase Transition

Philippe Leblond

Draft v3.10 — computational companion

`src/bh_graph` + `scripts/generate_figures.py` reproduce all figures (277 tests)

September 25, 2026

## Abstract

We formalize the intuition that a black hole interior is an almost-perfect all:all entanglement graph. Internal edges cost no exterior space; each of the  $k$  exterior legs costs  $\sim l_p^2$  of horizon area, so  $A(k) = k l_p^2$  independent of interior node count  $N$ . We show (1) the complete graph  $K_N$  destroys interior distance and, under finite-speed random circuits, derives the Sekino–Susskind  $t_* \sim \log N$  fast-scrambling bound; (2) mass enters via a derived  $k^*(N) = 16\pi(\varepsilon N/l_p)^2$  fixed point, spin/charge consume part of the exterior budget (extremal Kerr keeps half the legs), and the  $k \rightarrow 0$  monogamy limit is a baby-universe pinch-off with an explicit nonlinear CKW frontier; (3) micro-holes are pointlike until  $k l_p^2 \geq 4\pi r_{\text{point}}^2$ , then pop a horizon via a computed QES/island crossing, with an LQG-style area gap. Evaporation as leg surgery reproduces the exact Page curve (Page 1993,  $-1/2$  nat correction included) with Haar-typical fluctuations, and Hayden–Preskill mirror recovery works past  $k \sim N/2$ . Simulations in `bh_graph` accompany each claim.

## 1 Sec 1: All:all = no interior space

Interior wiring  $G_N$  compared across  $K_N$ , chain, grid, random regular under SI spread (cover time = seed eccentricity). Only  $K_N$  has diameter 1 and cover time 1 at every  $N$  (Figs. 1–2). Finite-speed random-matching circuits (Appendix A) promote this to a derived  $t_* \approx \log_2 N / \log_2(1+p)$ .

| family         | diameter         | $t_{\text{cover}}$ (SI) | $t_*$ (circuit) | gap              |
|----------------|------------------|-------------------------|-----------------|------------------|
| $K_N$          | 1                | 1                       | $\log_2 N$      | $N$              |
| chain          | $N-1$            | $\sim N/2$              | $\sim N/v$      | $\sim \pi^2/N^2$ |
| grid           | $\sim 2\sqrt{N}$ | $\sim 2\sqrt{N}$        | $O(\sqrt{N})$   | $O(1/N)$         |
| random-regular | $\sim \log N$    | $\sim \log N$           | $\sim \log N$   | $O(1)$           |

Table 1: Scrambling hierarchy; `bh_graph.scrambling`, `circuits`.

## 2 Sec 2: Horizon area counts exterior legs

Interior pairs  $N(N-1)/2$  plus  $k \ll N^2$  exterior legs:

$$A(k) = k l_p^2, \quad R(k) = \sqrt{k l_p^2 / 4\pi}, \quad (1)$$

Fig 1 — Sec 1: fast scrambling hierarchy




Figure 1: SI cover time and diameter: all:all stays 1.

so  $N$  drops out (Fig. 3). Schwarzschild consistency forces  $k(M) = 4\pi(2M)^2/l_p^2 \propto M^2$ ; Appendix B derives this from MaxEnt + self-gravity, and Appendix H extends it to Kerr–Newman  $k_{\text{eff}}(M, a, Q) = A(M, a, Q)/l_p^2$  with extremal Kerr at half the legs. Monogamy (linear toy Fig. 4; nonlinear CKW Fig. 14) sends  $e_{\text{int}} \rightarrow 1$ ,  $k \rightarrow 0$ : pinch-off into a baby universe.

### 3 Sec 3: Micro-hole phase transition

A point region of radius  $r_{\text{point}}$  embeds  $k$  legs iff  $k l_p^2 < 4\pi r_{\text{point}}^2$ . Hence

$$R_{\text{obs}}(k) = \begin{cases} r_{\text{point}} & k \leq k_{\text{crit}} \\ \sqrt{k l_p^2 / 4\pi} & k > k_{\text{crit}} \end{cases}, \quad k_{\text{crit}} = 4\pi r_{\text{point}}^2 / l_p^2, \quad (2)$$

with LQG-style quantization  $A \in \{0\} \cup [A_{\text{min}}, \infty)$  (Figs. 5–6). Appendix C shows the pop is a computed QES/island crossing at  $k_{\text{page}} = S_0 / (2s_{\text{leg}} - l_p^2/4)$ , existing iff  $s_{\text{leg}} > l_p^2/4$ .

## 4 Discussion

Evaporation = leg surgery with exact Page curve (Appendices D, I) and Hayden–Preskill mirror past  $k \sim N/2$  (Appendix F). Open: deriving  $\varepsilon$  and  $r_{\text{point}}$  coefficients from a UV embedding; Kerr Page curve with spin budget; OTOC Lyapunov check (Appendix L).

### A1 Finite-speed circuits: $t_* \sim \log N$ (Fig. 7)

One gate/qubit/step; random matching (all:all) vs dimer covering (chain). Early growth exponential on all:all; chain ballistic. Measured  $N = 8 \dots 128$ .

## A2 Deriving $k(N)$ (Fig. 8)

MaxEnt bound  $k \geq N s_{\text{node}}/s_{\text{leg}}$  plus  $R = \sqrt{k l_p^2/4\pi} = 2E$ ,  $E = \varepsilon N$ : fixed point  $k^*(N) = 16\pi(\varepsilon N/l_p)^2$ , stable from any start. Predicts  $\alpha(N) = k^*/N \propto N$ .

## A3 QES pop from entropy crossing (Fig. 9)

$S_{\text{no}} = k s_{\text{leg}}$  vs  $S_{\text{isl}} = k l_p^2/4 + \max(S_0 - k s_{\text{leg}}, 0)$ ; crossing iff  $s_{\text{leg}} > l_p^2/4$ . Explicit core+legs min-cut agrees.

## A4 Page curve from leg surgery (Fig. 10)

$S_{\text{rad}}(t) = \min(t, N_{\text{eff}} - t)$ ; area identical for wiring-only vs standard evaporation:  $A$  tracks  $k$  alone.

## A5 QEC mirror (Fig. 11)

$\text{err}(k) = \min(1/2, 2^{N/2+1-k})$ ; 99% threshold  $k \geq N/2 + 1 + \log_2 100$ ;  $k \rightarrow 0$  seals information off.

## A6 Robustness (Fig. 11)

Log slope  $1/\log_2(1+p)$  at all  $p$ ; exact quadraticity at all  $\varepsilon$ ; sharp QES boundary at  $s_{\text{leg}} = l_p^2/4$ .

## A7 Kerr–Newman (Fig. 12)

$r_+ = M + \sqrt{M^2 - a^2 - Q^2}$ ,  $A = 4\pi(r_+^2 + a^2)$ ; spin budget  $1 - A(M, a)/A(M, 0) \rightarrow 1/2$  at extremality. Area law survives.

## A8 Exact Page curve (Fig. 13)

$S(m, n) = H(mn) - H(n) - (m-1)/2n$  nats;  $-1/2$  nat dip at turnover; Haar samples ( $N = 8$ ) track the mean: typical, not tuned.

## A9 Nonlinear monogamy (Fig. 14)

Family  $\cos t|\Phi^+\rangle|0\rangle + \sin t|00\rangle|1\rangle$ : Wootters  $C_{AB}^2$  vs  $\tau_{E|AB}$  frontier lies strictly below  $x + y = 1$  (e.g. 0.70 at  $t = 0.3$ ); CKW  $\geq 0$  everywhere; baby universe at  $(1, 0)$ .

## A10 OTOC + phenomenology (Fig. 15)

$C(t) \sim e^{\lambda t}/N$  on all:all gives  $t^* = \log N/\lambda$  vs ballistic  $N/v$  on chains. PBHs evaporating today  $\sim 10^{18}\text{--}10^{19}$  Planck masses; sub-critical holes evaporate without greybody suppression; echo toy  $\Delta t \sim M \log k$ ; nearest-term test is analogue scrambling on programmable processors.

## A11 Tensor network derives $\varepsilon$ (Fig. 16)

Random star TN verifies  $S_{\text{bdy}} = \min(N \log d, k \log D)$ ; minimal wiring  $k_{\text{min}} = S_{\text{bulk}}/\log D$ . Two routes fix  $\varepsilon$ : QES coincidence  $\varepsilon = (l_p/N)\sqrt{k_{\text{page}}/16\pi}$  and crossover  $\varepsilon = l_p\sqrt{s_{\text{node}}/(16\pi \log D N_{\text{match}})}$ .

## A12 Kerr Page curve (Fig. 17)

$M \propto (1 - t/T)^{1/3}$ ,  $J$  shed faster; high spin lowers the peak and *delays* turnover (spin-shedding offsets early area loss):  $t/T = 0.73$  at  $a_0 = 0.95$  vs  $0.54$  Schwarzschild.

## A13 SYK ED vs chain (Fig. 18)

$N = 6, 8, 10$  Majorana SYK vs Ising chains, infinite- $T$  OTOC: SYK reaches threshold first and  $t^*$  stays flat while the chain's grows linearly. MSS saturation cited, hierarchy reproduced.

## A14 GWTC leg creation (Fig. 19)

All 32 GWTC-3 BBH medians satisfy  $k_f > k_1 + k_2$  (median +77% at 4% radiated): mergers create exterior legs. Spin caveat  $\sim 13\% \ll$  margin.

## A15 Hardware literature + prediction (Fig. 20)

Gärtner 2017 (all:all ions,  $m \leq 8$  OTOC coherences) and Mi 2021 (Sycamore grid, ballistic OTOC decay) both consistent. Head-to-head prediction at  $N = 53$ : 7.9 vs 14.6 steps; null result falsifies Sec 1/A.

## A16 TeV recast + echoes + EHT (Figs. 21–22)

LHC masses sit at  $k \sim 11\text{--}17$  vs  $k_{\text{crit}} \sim 50$ : thermal nulls expected. Echo spacings  $0.03\text{--}0.3\text{ s}$  land in searched windows (amplitude unmodeled). Shadow deviation  $\sim 10^{-48}$ : EHT-identical by 47 orders.

## A17 Posteriors + cosmos + Krylov (Figs. 23–25)

8350 GW150914 samples:  $P(\Delta k > 0) = 100\%$ , median +57% (Kerr both ends). Cosmic horizon owns the wiring budget ( $10^{123}$  vs  $10^{101}$ ). SYK saturates  $\sim 2\times$  the chain's Krylov spread.

## A18 Collapse + cosmic history + lunch (Figs. 26–28)

Compactness-driven local $\rightarrow$ all:all transition: diameter/gap/erasure switch together. Cosmic legs  $10^{122.3} \rightarrow 10^{123.1}$ ; patch unscrambled ( $t^* \approx 4000\text{ Gyr}$ ). Lunch  $C/k$  diverges in evaporation; remnant DM needs ultralight.

## A19 Bounds kill remnants + quench protocol (Fig. 29)

Vendored PBH bounds vs required- $\beta$ : 40+ orders apart;  $\beta > 1$  above  $10^6$  g. Same-device grid  $\rightarrow$  all:all quench predicts  $t^*$   $8 \rightarrow 1$ , gap  $0.27 \rightarrow 36$  — no cross-platform systematics.

## A20 Healing lag (Fig. 30)

$dA/dt = (kl_p^2 - A)/\tau$ ,  $\tau = 11.24M$  from ringdown (3.5 ms at  $63 M_\odot$ ). Ladder: healing, scrambling  $66\times$ , Page, evaporation; final sigh carries  $\sim 10^{-78}$  of the mass.

## A21 MSS + qualifier (Fig. 31)

Thermal SYK  $\lambda/2\pi T = 0.50 \rightarrow 0.68$  (bound held, finite-size). Uniform all:all scrambles worse (randomness load-bearing). LVK gives  $\alpha \in [9.0, 12.4]$ ; 11.24 sits inside.

## A22 Big SYK + typicality (Figs. 32–33)

Sparse  $N \leq 20$ :  $t^*$  flat vs chain  $1.8 \rightarrow 6.2$ ; MSS stable at  $N = 16$ . TN spectrum matches Marchenko-Pastur; analytic  $T(E)$  grounds  $T \equiv 1$  sub-critical emission.

## A23 Congestion + charge + bandwidth (Figs. 34–36)

Bubble iff  $\chi = kl_p^2/4\pi r_{foot}^2 > 1$ ; spread giants delocalized. Charge pins  $q = 4\pi Q^2$  legs (pinch/pointlike/extremal map). Evacuation empties babies; info-per-leg diverges  $\iff$  cloning risk.

## A24 Kill list (Figs. 37–39)

AF quench: kill below ratio 1.3 (predicted  $2\text{--}3\times$ ).  $\alpha$  monitor awaits O4/O5 rows. Min-rule violations shrink; Ising  $s_{leg} = 0.55$  vs 0.25. LHC onset  $\sim 550$  TeV; soft tail below kills.

## A25 Big pop (Fig. 40)

Hidden delocalized giants pop  $R_b(k)$  horizons on concentration:  $10^8 M_\odot$  in  $\sim 2$  Myr vs  $\sim 600$  Myr Eddington. Scenario sketch: metric, formation, and constraints still open.

## A26 Structure + scatter (Figs. 41–42)

PS halos at  $z = 8$  suffice (0.1% occupation);  $> 10^{12} M_\odot$  hosts excluded. Overmassive tail maps  $10\% \rightarrow f_w \approx 0.18$ ; null-of-300 kills the channel.

## A27 Remnant obituary + EMD window (Fig. 29, rewritten)

Formula corrected (growth + EMD): dead above  $\sim 10^6$  g and below  $\sim 10^5$  g (overclose); viable sweet spot  $\sim 4 \times 10^5$  g, currently unconstrained, poltergeist-probed in future.

## A28 Entropic Newton (Fig. 43)

Leg screens + equipartition + Bekenstein give  $F = M_1 M_2 / r^2$  exact (slope  $-2$  to  $10^{-9}$ ); leapfrog orbits close,  $T^2 \propto r^3$ . Flux  $\propto 1/r^2$  distinguished from force (temperature corrects).

## A29 Redshift (Fig. 44)

GPS  $+5.29 \times 10^{-10}$  and Pound-Rebka  $2.55 \times 10^{-15}$  derived (no new parameters). Congestion fronts give tortoise divergence;  $\alpha = 1$  matches Schwarzschild  $dr/dt$  exactly.

## A30 Einstein-Hilbert routes (Fig. 45)

Heat  $a_1$  responds to curvature; Ollivier signs exact (flat/neg/pos); Clausius fixes  $\varepsilon = \kappa/8\pi$ ,  $G = 1$ . Bridges labeled, Raychaudhuri-for-legs open.

## A31 Fission + audit + tension + GW250114 (Figs. 46–47)

$K_N^2$  two-mouth states, non-traversable; saturation 29% vs no-loss 0% locked by test. Single  $M(k)$  map audits all bounds. Tension  $E(d)$  parametrized with  $\sigma$  floors. GW250114  $\eta_A$  needs Kerr.

## A32 Overtones + tension no-go (Fig. 48)

PT barrier fit to fundamental predicts 1:3:5 vs 1:3.08:5.38 ( $< 8\%$ ). Bridge-volume tension undetermined  $\sim 10^{100}$ ;  $\sigma$  stays bounded, microphysics localized.

## A33 Big SYK to $N = 24$ (Fig. 32 ext.)

Triplet builder (verified vs ED);  $t^*$  flat 1.28–1.30 to 12 qubits vs chain 7.4; thermal  $\lambda/2\pi T$  0.66–0.86 ( $N = 10$ –24), below bound, upward within fit noise.

## A34 Lensing passes, Mercury gaps (Fig. 49)

Fermat  $n = 1/c_{eff}$  gives full  $4M/b$  (naive  $2M/b$  bet lost); 2nd order 3.36 vs 1.94. Orbits need  $g_{rr}$ : Newtonian 0 vs  $43''/\text{cy}$  gap. Lensing now chromatic via phase velocity:  $\Delta\alpha/\alpha \sim 10^{-56}$  optical (Fig. 55), achromaticity held by  $\sim 50$  orders. (Update: the orbit gap is closed by the Strain section below.)

## A35 b-crit kills isotropic (Fig. 50)

Bouguer gives  $8M$  vs GR  $3\sqrt{3}M$ : excluded  $\approx 3.6\sigma$  by EHT. Tangential must be fast (all:all); target window  $[4.42, 5.98]M$  locked.

## A36 Lattice dispersion (Fig. 51)

$v_g$  even in  $k$ : no linear term (Fermi linear bounds evaded by symmetry).  $E_{QG,2} = \sqrt{8} E_P$ ,  $10^8 \times$  above Fermi quadratic bound.

### A37 Footprint ringdown (Figs. 52–53)

Mass comb  $1/2k$ ; echoes with derived  $R \sim (\omega/\omega_P)^2 \sim 10^{-160}$ ;  $\ell(\ell+1) \leq k$  bites micro-holes; leg lines  $37.6\times$  finer than QNM.

### A38 Wave lab on defective fabric (Fig. 56)

FDTD beams: unbiased centroids, deficit  $\propto \omega^2$  ( $16\% \rightarrow 0.2\%$ ); fabric smoothness bounded to  $\varepsilon \lesssim 10^{-14}$  (TeV/Gpc).

### A39 Persistent walks (Fig. 57)

Hop-budget time dilation  $\sqrt{1-v^2}$ ; degree drift  $\sim 1/r^3$  (not Newton:  $T(r)$  load-bearing); diffusive-to-ballistic morphologies; spin effaced, Mercury MPD  $\sim 4 \times 10^{-14}$ .

### A40 Strain closes Mercury (Fig. 58)

Naive  $dl = dr/(1 - \sqrt{\chi})$  gives  $\gamma = 2$  (Cassini-excluded); tortuosity  $dl = (1 + \sqrt{\chi}/2)dr$  gives  $h = (1 + x/2)^2$ ,  $\gamma = 1$  exactly, and Mercury  $42.99''/\text{cy}$  by direct geodesic integration — the  $g_{rr}$  sector derived.

### A41 Weak-field OR profile (Fig. 59)

Grid+hub with flux stubs: radial Ollivier-Ricci negative in all configs (sign of  $h > 1$  derived from micro); scaling  $p \approx 0.9 \pm 0.5$  suggestive, not resolved — partial micro-derivation, bigger lattices needed.

### A42 PPN ledger + peel-off (Fig. 60)

Full table:  $\gamma = 1$  derived,  $\beta$  unresolved (not quoted),  $\alpha_{1,2}, \xi$  untouched; ours-vs-GR perihelion  $= -0.75 M/a$  measured.

### A43 Formal pop + Hamiltonian sketch (Fig. 61)

Packing forces the pop past  $[4\pi r^2/l_p^2]$ ;  $s_{leg} > 1/4$  boxed as postulate.  $H = H_{SYK} + H_{legs}$ :  $t^*$  survives weak legs,  $\sim 2\times$  slower at  $\lambda = 3$  (sketch,  $N = 8$ ).

### A44 Reduction: $k(M) \iff R_s(M)$

$M \mapsto k$  factors entirely through  $M \mapsto R_s$  (explicit function argument;  $3M$  input gives  $36\pi M^2$ ). Circle shrunk to one statement.

### A45 Bridge 2: constancy passes, $\eta = \ln 2$ (Fig. 62)

$S/k$  flat to  $< 1\%$  across cuts, but at  $\ln 2$ , not  $1/4$ : face-value  $G = 0.36$ . Patch =  $4 \ln 2$  or sub-maximal legs; bridges  $3/1$  gated.

## A46 Postulate B adopted

Physical legs at  $1/4$  nat (random  $\ln 2$  is kinematic max):  $\eta = 1/4$ ,  $G = 1$ ,  $S = k/4$  close with patch  $= 1$ . Mechanism unknown; bridges proceed conditionally, B load-bearing.

## A47 Walk no-go + escape (Fig. 63)

All smooth weight rules cubic ( $-2.99$ ); only absurd  $e^{2\sqrt{d-d_0}}$  escapes. Constant  $\mu$  keeps  $-3$  exactly;  $\mu(\chi)$  fluctuation walk gives  $-1.95 \approx -2$  modulo labeled  $\sqrt{\chi}$  assumption.

## A48 Review kills + Green shorting (Fig. 64)

S-wave cut (Rayleigh 11.4%, 52% speckle, interference);  $-1/4$  cut ( $n_d^{-1}$  universal). Harmonic drift fails: legs short gradients ( $h$ -steepness  $9.2 \rightarrow 4.1$ ); AS temperature triply load-bearing.

## A49 Entropy-capacity tension

Purity +  $E = \varepsilon N + S = k/4$  force  $N \leq 125$  (macro violated, any constant  $\varepsilon$ ). Holographic legs, running  $\varepsilon$ , or no interior factor; B downstream of Hilbert-space question.

## Reproducibility

```
pip install -e .; python -m pytest tests/ -q; python scripts/generate_figures.py; streamlit run app.py.
```

## References

- [1] Sekino & Susskind, JHEP 2008.
- [2] Van Raamsdonk, Gen. Rel. Grav. 2010.
- [3] Maldacena & Susskind, Fortsch. Phys. 2013.
- [4] Bekenstein, Phys. Rev. D 1973; Hawking 1975.
- [5] Rovelli & Smolin, Nucl. Phys. B 1995.
- [6] Penington; Almheiri et al., JHEP 2019–2020.
- [7] Hayden & Preskill, JHEP 2007.
- [8] Page, Phys. Rev. Lett. 1993.
- [9] Coffman, Kundu & Wootters, Phys. Rev. A 2000.