

Higher-genus GW theory of smooth CY hypersurfaces in weighted \mathbb{P}^4

Patrick Lei

March 30, 2025

Columbia University

Overview

Introduction

Mixed-Spin-P fields

Calculations

Future work

Introduction

Our goal is to understand the *Gromov-Witten theory* of a **Calabi-Yau threefold**, which in this talk will be called **Z** .

Our goal is to understand the *Gromov-Witten theory* of a **Calabi-Yau threefold**, which in this talk will be called Z .

Definition

The moduli space $\overline{M}_{g,0}(Z, d)$ of genus g , degree d unmarked stable maps to Z has a virtual cycle in degree 0 . Define the *GW invariant*

$$N_{g,d} := \deg[\overline{M}_{g,0}(Z, d)]^{\text{vir}} \in \mathbb{Q}.$$

Our goal is to understand the *Gromov-Witten theory* of a **Calabi-Yau threefold**, which in this talk will be called Z .

Definition

The moduli space $\overline{M}_{g,0}(Z, d)$ of genus g , degree d unmarked stable maps to Z has a virtual cycle in degree 0 . Define the *GW invariant*

$$N_{g,d} := \deg[\overline{M}_{g,0}(Z, d)]^{\text{vir}} \in \mathbb{Q}.$$

Goal: Calculate all $N_{g,d}$.

Our goal is to understand the *Gromov-Witten theory* of a **Calabi-Yau threefold**, which in this talk will be called Z .

Definition

The moduli space $\overline{M}_{g,0}(Z, d)$ of genus g , degree d unmarked stable maps to Z has a virtual cycle in degree 0 . Define the *GW invariant*

$$N_{g,d} := \deg[\overline{M}_{g,0}(Z, d)]^{\text{vir}} \in \mathbb{Q}.$$

Goal: Calculate all $N_{g,d}$. This is too hard!

Our goal is to understand the *Gromov-Witten theory* of a **Calabi-Yau threefold**, which in this talk will be called Z .

Definition

The moduli space $\overline{M}_{g,0}(Z, d)$ of genus g , degree d unmarked stable maps to Z has a virtual cycle in degree 0 . Define the *GW invariant*

$$N_{g,d} := \deg[\overline{M}_{g,0}(Z, d)]^{\text{vir}} \in \mathbb{Q}.$$

Goal: Say something about the behavior of the generating series of all $N_{g,d}$ when we fix the genus.

Our results

We are able to prove the following results for the threefolds $Z_6 \subset \mathbb{P}(1, 1, 1, 1, 2)$, $Z_8 \subset \mathbb{P}(1, 1, 1, 1, 4)$, and $Z_{10} \subset \mathbb{P}(1, 1, 1, 2, 5)$:

Our results

We are able to prove the following results for the threefolds $Z_6 \subset \mathbb{P}(1, 1, 1, 1, 2)$, $Z_8 \subset \mathbb{P}(1, 1, 1, 1, 4)$, and $Z_{10} \subset \mathbb{P}(1, 1, 1, 2, 5)$:

1. If $F_g = \sum_d N_{g,d} Q^d$ is the generating series of genus g invariants, we prove that a normalized version P_g is a polynomial in five generators $A_1, B = B_1, B_2, B_3, X$ defined using only genus-zero data.

Our results

We are able to prove the following results for the threefolds $Z_6 \subset \mathbb{P}(1, 1, 1, 1, 2)$, $Z_8 \subset \mathbb{P}(1, 1, 1, 1, 4)$, and $Z_{10} \subset \mathbb{P}(1, 1, 1, 2, 5)$:

2. More precisely, Coates-Corti-Lee-Tseng proved that the genus-0 GW theory of Z is controlled by the series

$$\begin{aligned} I(q, z) &:= z \sum_{d \geq 0} q^d \frac{\prod_{m=1}^{6d} (6H + mz)}{\prod_{m=1}^d (H + mz)^4 \prod_{m=1}^{2d} (2H + mz)} \\ &= I_0(q)z + I_1(q)H + I_2(q)\frac{H^2}{z} + I_3(q)\frac{H^3}{z^2}. \end{aligned}$$

Our results

We are able to prove the following results for the threefolds $Z_6 \subset \mathbb{P}(1, 1, 1, 1, 2)$, $Z_8 \subset \mathbb{P}(1, 1, 1, 1, 4)$, and $Z_{10} \subset \mathbb{P}(1, 1, 1, 2, 5)$:

3. Define $I_{11}(q) := D\left(\frac{I_1(q)}{I_0(q)}\right)$, where $D = q \frac{d}{dq}$. Then define

$$A := \frac{DI_{11}}{I_{11}}, \quad B_k := \frac{D^k I_0}{I_0}, \quad X := 1 - \frac{1}{1 - \frac{6^6}{2^2} q}.$$

Also define $Y := 1 - X$.

Our results

We are able to prove the following results for the threefolds $Z_6 \subset \mathbb{P}(1, 1, 1, 1, 2)$, $Z_8 \subset \mathbb{P}(1, 1, 1, 1, 4)$, and $Z_{10} \subset \mathbb{P}(1, 1, 1, 2, 5)$:

4. The first result, conjectured by Yamaguchi-Yau (2004), is that

$$P_{g,n} = \frac{(3Y)^{g-1} I_{11}^n}{I_0^{2g-2}} \left(Q \frac{d}{dQ} \right)^n F_g(Q) \in \mathbb{Q}[A, B, B_2, B_3, X]$$

after the substitution $Q = qe^{l_1/l_0}$.

We are able to prove the following results for the threefolds $Z_6 \subset \mathbb{P}(1, 1, 1, 1, 2)$, $Z_8 \subset \mathbb{P}(1, 1, 1, 1, 4)$, and $Z_{10} \subset \mathbb{P}(1, 1, 1, 2, 5)$:

5. The second result is the equality

$$P_{1,1} = -\frac{1}{2}A - \frac{21}{2}B - \frac{1}{12}X - \frac{7}{4}.$$

Our results

We are able to prove the following results for the threefolds $Z_6 \subset \mathbb{P}(1, 1, 1, 1, 2)$, $Z_8 \subset \mathbb{P}(1, 1, 1, 1, 4)$, and $Z_{10} \subset \mathbb{P}(1, 1, 1, 2, 5)$:

6. Consider the following Feynman rule (sum over *stable graphs*) defined by BCOV (1993) defining a power series $f_{g,m,n}$. First, introduce *propagators* $E_\psi = B$, $E_{\varphi\varphi} = A + 2B$, $E_{\varphi\psi} = -B_2$, and $E_{\psi\psi} = -B_3 + (B - X)B_2 - \frac{13}{36}BX$.

Our results

We are able to prove the following results for the threefolds $Z_6 \subset \mathbb{P}(1, 1, 1, 1, 2)$, $Z_8 \subset \mathbb{P}(1, 1, 1, 1, 4)$, and $Z_{10} \subset \mathbb{P}(1, 1, 1, 2, 5)$:

7. Place $\varphi - E_\psi \psi$ at the first m vertices and ψ at the last n vertices, place

$$E_{\varphi\varphi} \varphi \otimes \varphi + E_{\varphi\psi}(\varphi \otimes \psi + \psi \otimes \varphi) + E_{\psi\psi} \psi \otimes \psi$$

at each edge, and place

$$\varphi^{\otimes m} \otimes \psi^{\otimes n} \mapsto P_{g,m,n} := \frac{(2g - 2 + m + n - 1)!}{(2g - 2 + m - 1)!} P_{g,m}$$

at each leg. Also, set $P_{1,0,1} = -\frac{19}{2}$.

Our results

We are able to prove the following results for the threefolds $Z_6 \subset \mathbb{P}(1, 1, 1, 1, 2)$, $Z_8 \subset \mathbb{P}(1, 1, 1, 1, 4)$, and $Z_{10} \subset \mathbb{P}(1, 1, 1, 2, 5)$:

8. Our result is that the output $f_{g,m,n}$ is a polynomial of degree at most $3g - 3 + m$ in X . This implies the *modular anomaly equations*

$$-\partial_A P_g = \frac{1}{2} \left(P_{g-1,2} + \sum_{g_1+g_2=g} P_{g_1,1} P_{g_2,2} \right),$$
$$\left(-2\partial_A + \partial_B + (A + 2B)\partial_{B_2} - \left((B - X)(A + 2B) - B_2 - \frac{13}{36}X \right) \partial_{B_3} \right) P_g = 0.$$

- Our first result (polynomiality) was proved for the quintic independently by Chang-Guo-Li and Guo-Janda-Ruan in 2018.

- Our first result (polynomiality) was proved for the quintic independently by Chang-Guo-Li and Guo-Janda-Ruan in 2018.
- Our genus-one formula was first proved for the quintic by Zinger (2007) and proved for complete intersections in projective space by Popa (2010).

- Our first result (polynomiality) was proved for the quintic independently by Chang-Guo-Li and Guo-Janda-Ruan in 2018.
- Our genus-one formula was first proved for the quintic by Zinger (2007) and proved for complete intersections in projective space by Popa (2010).
- The modular anomaly equations were proved for the quintic independently by Chang-Guo-Li and Guo-Janda-Ruan in 2018.

Mixed-Spin-P fields

Our approach

We use the approach of **Mixed-Spin-P fields**, which were introduced by Chang-Li-Li-Liu (2015, 2016) and Chang-Guo-Li-Li (2018).

Our approach

We use the approach of **Mixed-Spin-P fields**, which were introduced by Chang-Li-Li-Liu (2015, 2016) and Chang-Guo-Li-Li (2018).

The original geometric intuition was to implement the master space idea of Thaddeus, but in order to perform calculations we need to introduce a parameter N (which is a positive integer) to the theory.

Let $(\mathbb{C}^*)^3$ act on \mathbb{C}^{N+7} with the following weights:

Let $(\mathbb{C}^*)^3$ act on \mathbb{C}^{N+7} with the following weights:

$$\left[\begin{array}{cccccccccc} x_1 & x_2 & x_3 & x_4 & x_5 & p & u_1 & \dots & u_N & v \\ \hline 1 & 1 & 1 & 1 & 2 & -6 & 1 & \dots & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & \dots & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & \dots & 0 & 0 \end{array} \right].$$

Let $(\mathbb{C}^\times)^3$ act on \mathbb{C}^{N+7} with the following weights:

$$\left[\begin{array}{cccccccccc} x_1 & x_2 & x_3 & x_4 & x_5 & p & u_1 & \dots & u_N & v \\ \hline 1 & 1 & 1 & 1 & 2 & -6 & 1 & \dots & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & \dots & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & \dots & 0 & 0 \end{array} \right].$$

Let $W := W_{g,n,d}$ be the stack of commutative diagrams

$$\begin{array}{ccccc} \mathcal{C} & \longrightarrow & [\mathbb{C}^{N+7}/(\mathbb{C}^\times)^3] & \longrightarrow & B(\mathbb{C}^\times)^3 \\ & \searrow \omega^{\log} & & & \downarrow \text{pr}_3 \\ & & & & B\mathbb{C}^\times \end{array}$$

subject to a stability condition.

W is not proper, but it has a cosection localized virtual cycle $[W]^{\text{vir}}$ supported on a proper closed substack W^- .

W is not proper, but it has a cosection localized virtual cycle $[W]^{\text{vir}}$ supported on a proper closed substack W^- . This virtual cycle is equivariant with respect to the action of $(\mathbb{C}^\times)^N$ on W by scaling (u_1, \dots, u_N) .

W is not proper, but it has a cosection localized virtual cycle $[W]^{\text{vir}}$ supported on a proper closed substack W^- . This virtual cycle is equivariant with respect to the action of $(\mathbb{C}^\times)^N$ on W by scaling (u_1, \dots, u_N) .

Theorem (Chang-Kiem-Li, 2015)

Indexing the fixed loci by Γ , we have

$$[W]^{\text{vir}} = \sum_{\Gamma} \frac{[W_{\Gamma}]^{\text{vir}}}{e(N_{\Gamma}^{\text{vir}})}.$$

Fixed loci in the MSP moduli space \mathcal{W} are indexed by graphs Γ . There are three kinds (denoted by levels) of vertices:

1. Level $\mathbf{0}$ vertices contribute GW invariants of \mathbf{Z} ;

Fixed loci in the MSP moduli space \mathcal{W} are indexed by graphs Γ . There are three kinds (denoted by levels) of vertices:

1. Level **0** vertices contribute GW invariants of \mathcal{Z} ;
2. Level **1** vertices contribute Hodge integrals;

Fixed loci in the MSP moduli space \mathcal{W} are indexed by graphs Γ . There are three kinds (denoted by levels) of vertices:

1. Level $\mathbf{0}$ vertices contribute GW invariants of \mathbf{Z} ;
2. Level $\mathbf{1}$ vertices contribute Hodge integrals;
3. Level ∞ vertices contribute FJRW invariants.

Fixed loci in the MSP moduli space \mathcal{W} are indexed by graphs Γ . There are three kinds (denoted by levels) of vertices:

1. Level 0 vertices contribute GW invariants of \mathbb{Z} ;
2. Level 1 vertices contribute Hodge integrals;
3. Level ∞ vertices contribute FJRW invariants.

Theorem

If Γ has an edge between a level 0 and level ∞ vertex, then $[W_\Gamma]^{\text{vir}} = 0$.

Virtual localization II

Fixed loci in the MSP moduli space \mathcal{W} are indexed by graphs Γ . There are three kinds (denoted by levels) of vertices:

1. Level 0 vertices contribute GW invariants of \mathbb{Z} ;
2. Level 1 vertices contribute Hodge integrals;
3. Level ∞ vertices contribute FJRW invariants.

Theorem

If Γ has an edge between a level 0 and level ∞ vertex, then $[W_\Gamma]^{\text{vir}} = 0$.

Definition

Define the MSP $[0, 1]$ theory by only considering the contributions of graphs without level ∞ vertices.

Calculations

Lemma

Genus zero MSP invariants equal genus zero Gromov-Witten invariants of a degree 6 hypersurface in $\mathbb{P}(1, 1, 1, 1, 2, 1, \dots, 1)$.

Genus zero MSP theory

Lemma

Genus zero MSP invariants equal genus zero Gromov-Witten invariants of a degree 6 hypersurface in $\mathbb{P}(1, 1, 1, 1, 2, 1, \dots, 1)$.

This implies that all genus zero MSP invariants are recovered from the J -function

$$z \sum_{d \geq 0} q^d \frac{\prod_{m=1}^{6d} (6p + mz)}{\prod_{m=1}^d (p + mz)^4 \prod_{m=1}^{2d} (2p + mz) \prod_{m=1}^d ((p + mz)^N - t^N)}.$$

Here, we specialize the equivariant parameters to $t_\alpha = \zeta_N^\alpha t$ for $\alpha = 1, \dots, N$.

Define the R -matrix by the Birkhoff factorization

$$S^{\text{MSP}}(z)\Delta = R(z)S^{\text{loc}}(z),$$

where Δ comes from Quantum Riemann-Roch and S^{loc} is simply the direct sum of the S -matrix of Z and N copies of the S -matrix of a point.

Define the R -matrix by the Birkhoff factorization

$$S^{\text{MSP}}(z)\Delta = R(z)S^{\text{loc}}(z),$$

where Δ comes from Quantum Riemann-Roch and S^{loc} is simply the direct sum of the S -matrix of Z and N copies of the S -matrix of a point.

Theorem

The MSP $[0, 1]$ theory is a cohomological field theory given by the action

$$R(z) \cdot \left(\Omega^Z \oplus \bigoplus_{\alpha=1}^N \Omega^{\text{pt}_\alpha} \right)$$

of $R(z)$ on the direct sum of the GW theory of Z and N copies of the GW theory of a point.

Proof of polynomiality

Lemma

*All entries of $R(z)$ lie in $\mathbb{Q}[A, B, B_2, B_3, X]$ (possibly after normalization).
There are also explicit degree bounds for the entries coming from the N points, which are polynomials in X .*

Proof of polynomiality

Lemma

All entries of $R(z)$ lie in $\mathbb{Q}[A, B, B_2, B_3, X]$ (possibly after normalization).
There are also explicit degree bounds for the entries coming from the N points, which are polynomials in X .

Lemma

The MSP $[0, 1]$ correlator $\langle p^{a_1}, \dots, p^{a_n} \rangle_{g,n}^{[0,1]}$ is a polynomial in q of degree at most $g - 1 + \frac{3g - 3 + \sum a_i}{N}$.

Proof of polynomiality

Lemma

All entries of $R(z)$ lie in $\mathbb{Q}[A, B, B_2, B_3, X]$ (possibly after normalization). There are also explicit degree bounds for the entries coming from the N points, which are polynomials in X .

Lemma

The MSP $[0, 1]$ correlator $\langle p^{a_1}, \dots, p^{a_n} \rangle_{g,n}^{[0,1]}$ is a polynomial in q of degree at most $g - 1 + \frac{3g-3+\sum a_i}{N}$.

Corollary

$$P_{g,n} = \frac{(3Y)^{g-1} I_{11}^n}{I_0^{2g-2}} \left(Q \frac{d}{dQ} \right)^n F_g(Q) \Big|_{Q=qe^{l_1/l_0}} \in \mathbb{Q}[A, B, B_2, B_3, X].$$

Edge contributions in the action of the MSP R -matrix look like the propagators $E_{\varphi\varphi}$, $E_{\varphi\psi}$, and $E_{\psi\psi}$. For example, the edge contribution between two level $\mathbf{0}$ vertices starts with

$$E_{\psi}(1 \otimes H^2 + H^2 \otimes 1) + \frac{1}{2} \left(E_{\varphi\varphi} + \frac{13}{36} X \right) H \otimes H$$

up to a prefactor.

Definition

Consider the factorization

$$R(z) = R^X(z) \begin{pmatrix} R^A(z) & \\ & I_N \end{pmatrix},$$

where

$$R^A(z)^{-1} = I - \begin{pmatrix} 0 & zE_\psi & z^2E_{\varphi\psi} & \cdots \\ & 0 & zE_{\varphi\varphi} & \cdots \\ & & 0 & zE_\psi \\ & & & 0 \end{pmatrix}.$$

Lemma

The entries of $R^X(\mathbf{z})$ are polynomials in X with an explicit degree bound.

Feynman rule II

Lemma

The entries of $R^X(z)$ are polynomials in X with an explicit degree bound.

Corollary (MSP Feynman rule)

Let $f_{g,m,n}^A$ be the generating functions of the cohomological field theory

$$R^A(z) \cdot \Omega^Z.$$

Then $f_{g,m,n}^A$ is a polynomial in X of degree at most $3g - 3 + m$.

Feynman rule III

Recall that we let $f_{g,m,n}$ be the output of the physics Feynman rule.

Feynman rule III

Recall that we let $f_{g,m,n}$ be the output of the physics Feynman rule.

Theorem

We have

$$f_{g,m,n} = f_{g,m,n}^A - \delta_{g,1} \delta_{m,0} (n-1)!.$$

Feynman rule III

Recall that we let $f_{g,m,n}$ be the output of the physics Feynman rule.

Theorem

We have

$$f_{g,m,n} = f_{g,m,n}^A - \delta_{g,1} \delta_{m,0} (n-1)!.$$

As a corollary, we prove the physics Feynman rule.

Feynman rule III

Recall that we let $f_{g,m,n}$ be the output of the physics Feynman rule.

Theorem

We have

$$f_{g,m,n} = f_{g,m,n}^A - \delta_{g,1} \delta_{m,0} (n-1)!.$$

As a corollary, we prove the physics Feynman rule.

Corollary

We have the modular anomaly equations

$$\begin{aligned} -\partial_A P_g &= \frac{1}{2} \left(P_{g-1,2} + \sum_{g_1+g_2=g} P_{g_1,1} P_{g_2,2} \right), \\ \left(-2\partial_A + \partial_B + (A+2B)\partial_{B_2} - \left((B-X)(A+2B) - B_2 - \frac{13}{36}X \right) \partial_{B_3} \right) P_g &= 0. \end{aligned}$$

Feynman rule III

Recall that we let $f_{g,m,n}$ be the output of the physics Feynman rule.

Theorem

We have

$$f_{g,m,n} = f_{g,m,n}^A - \delta_{g,1} \delta_{m,0} (n-1)!.$$

As a corollary, we prove the physics Feynman rule.

Corollary

We have the modular anomaly equations

$$\begin{aligned} -\partial_A P_g &= \frac{1}{2} \left(P_{g-1,2} + \sum_{g_1+g_2=g} P_{g_1,1} P_{g_2,2} \right), \\ \left(-2\partial_A + \partial_B + (A+2B)\partial_{B_2} - \left((B-X)(A+2B) - B_2 - \frac{13}{36}X \right) \partial_{B_3} \right) P_g &= 0. \end{aligned}$$

Proof.

Differentiate the physics Feynman rule.



Future work

- We have proved results for Calabi-Yau threefolds with $h^{1,1} = 1$. A natural question is to extend these results to the case when $h^{1,1} > 1$.

Multi-parameter models

- We have proved results for Calabi-Yau threefolds with $h^{1,1} = 1$. A natural question is to extend these results to the case when $h^{1,1} > 1$.
- A natural direction is to study hypersurfaces in toric Fano orbifolds whose corresponding reflexive polytope is a simplex. This includes weighted projective space examples and the mirror quintic.

Multi-parameter models

- We have proved results for Calabi-Yau threefolds with $h^{1,1} = 1$. A natural question is to extend these results to the case when $h^{1,1} > 1$.
- A natural direction is to study hypersurfaces in toric Fano orbifolds whose corresponding reflexive polytope is a simplex. This includes weighted projective space examples and the mirror quintic.
- Stay tuned in the future!