

DEFORMATIONS OF SINGULARITIES

PATRICK LEI

ABSTRACT. We will study explicitly the embedded deformations of singular affine schemes via explicit lifting of equations and relations. We prove that embedded deformations of codimension 2 Cohen-Macaulay closed subschemes are unobstructed. As a corollary, Hilbert schemes of smooth surfaces are smooth. Finally, we give an example of an obstructed deformation.

We begin by fixing some notation. Let k be a field and $R = P/I$, where $P = k[x_1, \dots, x_n]$ and $I = (f_1, \dots, f_r)$ is an ideal. Throughout this lecture, we will denote local Artinian rings with residue field k by A, B, C, \dots and rings by R, S, T, \dots . Finally, denote $Z = \text{Spec } R$.

1. EXPLICIT CRITERIA FOR FLATNESS

We will study (embedded) deformations of singular affine schemes embedded in \mathbb{A}^n . The first thing we want to understand is to explicitly understand flatness of some R_A over A , where $R_A \otimes_A k = R$. We will write $R_A = P_A/I_A$, where $P_A = A[x_1, \dots, x_n] = A \otimes_k P$. Recall that over a Noetherian local ring S with residue field k , a module M is flat if and only if it is free, and this is equivalent to $\text{Tor}_1^S(M, k) = 0$ by standard results in commutative algebra.

Now consider the exact sequence

$$0 \rightarrow I_A \rightarrow P_A \rightarrow R_A \rightarrow 0.$$

After tensoring with k , we have

$$0 \rightarrow \text{Tor}_1(R_A, k) \rightarrow I_A \otimes_A k \rightarrow P \rightarrow R \rightarrow 0.$$

Therefore, we know that R_A is flat over A if and only if $I_A \otimes_A k = I$. We would like to understand this statement.

Consider a presentation

$$P_A^s \rightarrow P_A^r \rightarrow I_A \rightarrow 0$$

of I_A . Then we know R_A is flat over A if and only if after tensoring with k , we obtain an exact sequence

$$P^s \rightarrow P^r \rightarrow I \rightarrow 0.$$

Note that to give this presentation $P^s \rightarrow P^r \rightarrow I \rightarrow 0$ is the same as giving a complete set of relations among the generators of I .

Proposition 1.1. *Suppose that*

$$(1) \quad P^s \rightarrow P^r \rightarrow P \rightarrow R \rightarrow 0$$

is exact and

$$(2) \quad P_A^s \rightarrow P_A^r \rightarrow P_A \rightarrow R_A \rightarrow 0$$

is a complex such that $P_A^r \rightarrow R_A \rightarrow R_A \rightarrow 0$ is exact and tensoring (2) with k gives (1). Then R_A is flat over A .

Proof. Note that the hypotheses are equivalent to the fact that all relations in I can be lifted to I_A . Now given $g'_1, \dots, g'_r \in P_A$ such that

$$\sum_{i=1}^r g'_i f'_i = 0,$$

this clearly descends to a relation in I by killing the maximal ideal of A . But now if we choose a complete set of relations for I_A , this descends to a complete set of relations in I , so we may in fact assume that (2) is exact.

In this case, there exists some L_A such that the sequence splits as

$$P_A^s \rightarrow L_A \rightarrow 0 \quad 0 \rightarrow L_A \rightarrow P_A^r \rightarrow I_A \rightarrow 0 \quad 0 \rightarrow I_A \rightarrow P_A \rightarrow R_A \rightarrow 0.$$

By right exactness of the tensor product, we know $P_A^s \otimes k \rightarrow L_A \otimes k \rightarrow 0$ is exact. We also know that

$$L_A \otimes k \rightarrow P_A^r \otimes k \rightarrow I_A \otimes k \rightarrow 0$$

is exact, again by right exactness. But this means that $I_A \otimes k$ is the cokernel of $P^s \rightarrow P^r$, and therefore $I_A \otimes k = I$. This means that R_A is flat. \square

Corollary 1.2. *Let $R = P/I$ and $R_A = P_A/I_A$, where $I = (f_1, \dots, f_r)$ and $I_A = (f'_1, \dots, f'_r)$ such that f'_i is a lift of f_i . Then R_A is flat over A if and only if every relation among the f_i lifts to a relation among the f'_i .*

Remark 1.3. This result essentially gives us that first-order embedded deformations of $\text{Spec } R \subset \mathbb{A}^n$ are given by $\text{Hom}(I, R)$. The first-order (not embedded) deformations of Z are given by the cokernel of

$$0 \rightarrow T_X \rightarrow T_{\mathbb{A}^n}|_X \rightarrow N_{X/\mathbb{A}^n},$$

which arises from the exact sequence

$$I/I^2 \rightarrow \Omega_{\mathbb{A}^n}^1|_X \rightarrow \Omega_X^1 \rightarrow 0,$$

and this is supported on the singular points of X , so when X has isolated singularities, this is finite-dimensional.

Note that if $\text{Spec } R \subset \mathbb{A}^n$ is a complete intersection, then I is generated by a regular sequence, so in particular the Koszul complex is a free resolution of R and therefore there are only trivial relations among the f_i (this means the relations are generated by $f_i f_j - f_j f_i = 0$). Clearly, because we are only considering commutative rings (after all, this is normal algebraic geometry), this means that all deformations of $\text{Spec } R$ are unobstructed.

2. HILBERT SCHEMES OF SMOOTH SURFACES

We will prove that deformations of finite length closed subschemes of \mathbb{A}^2 are unobstructed. In particular, this will imply that the Hilbert scheme $\text{Hilb}(\mathbb{A}^2, n)$ is smooth.

Let $Z \subset \mathbb{A}^2$ be a closed subscheme of dimension 0. Then because $P = k[x, y]$ has dimension 2, there exists a free resolution

$$0 \rightarrow P^s \xrightarrow{(g_{ij})} P^r \rightarrow P \rightarrow R \rightarrow 0$$

of R . In this case it is possible to understand the matrix (g_{ij}) , and in fact this is the special case of a more general result. First, when we study the local behavior, we have the following result.

Theorem 2.1 (Hilbert, Burch). *Let P be a regular local ring of dimension n and $R = P/I$ be a Cohen-Macaulay quotient of codimension 2. Then there exists an $(r-1) \times r$ matrix $G = (g_{ij})$ whose maximal minors f_1, \dots, f_r minimally generate I , and there is a free resolution*

$$0 \rightarrow P^{r-1} \xrightarrow{(g_{ij})} P^r \xrightarrow{(f_i)} P \rightarrow R \rightarrow 0.$$

Proof. Note that the fact that the free resolution has this length is a corollary of the Auslander-Buchsbaum formula, which says that for a ring R and module M , we have

$$\text{depth } M + \text{proj. dim } M = \text{depth } R$$

and the fact that depth equals dimension for Cohen-Macaulay things. Thus we have a free resolution

$$0 \rightarrow P^{r-1} \xrightarrow{(g_{ij})} P^r \xrightarrow{(a_i)} P \rightarrow R \rightarrow 0,$$

where a_1, \dots, a_r are a minimal set of generators for I . Let f_i is $(-1)^i$ times the determinant of the i -th minor of g_{ij} . We will prove that the map (f_i) is the same as the map (a_i) ; clearly

$$0 \rightarrow P^{r-1} \xrightarrow{(g_{ij})} P^r \xrightarrow{(f_i)} P \rightarrow R \rightarrow 0.$$

is a resolution. This is because at the generic point of P , we know (g_{ij}) is injective, so at least one f_i is nonzero. But then we know $\text{coker}(g_{ij})$ is torsion-free (because I is torsion-free), and so it in fact must vanish by rank reasons. Thus (a_1, \dots, a_r) and (f_1, \dots, f_r) are isomorphic as P -modules.

At a codimension 1 point in $\text{Spec } P$, note that $0 \rightarrow P^{r-1} \rightarrow P^r \xrightarrow{(a_i)} P \rightarrow B \rightarrow 0$ is split exact (because I has codimension 2). This implies that at least one f_i is a unit, and thus (f_1, \dots, f_r) has codimension at least 2. But then the isomorphism $I \cong (f_1, \dots, f_r)$ is given by multiplication by some nonzero element of P which is a unit away from codimension 2. But this means it is a unit everywhere. \square

Considering the global picture in \mathbb{A}^n , we obtain the following result.

Theorem 2.2 (Hilbert, Schaps). *Let $Z = \text{Spec } R \subset \mathbb{A}^n$ be a Cohen-Macaulay closed subscheme of codimension 2. Then $R = P/I$ has a free resolution of the form*

$$0 \rightarrow P^{r-1} \xrightarrow{(g_{ij})} P^r \xrightarrow{(f_i)} P \rightarrow R \rightarrow 0$$

where the f_i are the maximal minors of the matrix (g_{ij}) .

This result in fact holds over any Artinian local ring A , which we will use later.

Next, we want to understand what happens if we choose some Artinian local ring with residue field k and lift the g_{ij} to g'_{ij} , where $g'_{ij} \in P_A$.

Theorem 2.3 (Schaps). *If A is a square zero extension of k , then the sequence*

$$0 \rightarrow P_A^{r-1} \xrightarrow{(g'_{ij})} P_A^r \xrightarrow{(f'_i)} P_A \rightarrow R_A \rightarrow 0$$

is exact. Moreover, any lifting of R over A arises by lifting the matrix (g_{ij}) .

Proof. We know that

$$L_A^\bullet := P_A^{r-1} \rightarrow P_A^r \rightarrow P_A$$

is a complex. This is because composing the two maps amounts to evaluating determinants with a repeated column. Because P_A is free (and therefore flat), we can tensor with the exact sequence

$$0 \rightarrow \mathfrak{m}_A \rightarrow A \rightarrow k \rightarrow 0$$

to obtain an exact sequence of complexes

$$0 \rightarrow L_A^\bullet \otimes_A \mathfrak{m}_A \rightarrow L_A^\bullet \rightarrow L_A^\bullet \otimes_A k \rightarrow 0.$$

Note that

$$L_A^\bullet \otimes_A k = P^{r-1} \xrightarrow{(g_{ij})} P^r \xrightarrow{(f_i)} P =: L^\bullet.$$

In particular, this term is exact by Hilbert-Schaps. In addition, clearly $L_A^\bullet \otimes_A \mathfrak{m}_A = L^\bullet \otimes_k \mathfrak{m}_A$ because $A \rightarrow k$ is a square zero extension, so the complex $L_A^\bullet \otimes_A \mathfrak{m}_A$ is exact. By the long exact sequence in homology, we know that L_A^\bullet is exact. Note that L^\bullet extends to an exact sequence

$$0 \rightarrow P^{r-1} \rightarrow P^r \rightarrow P \rightarrow R \rightarrow 0,$$

and L_A^\bullet extends to an exact sequence

$$0 \rightarrow P_A^{r-1} \rightarrow P_A^r \rightarrow P_A \rightarrow R_A \rightarrow 0.$$

However, by the homology long exact sequence, we have an exact sequence

$$0 \rightarrow R \otimes_k \mathfrak{m}_A \rightarrow R_A \rightarrow R \rightarrow 0.$$

But this implies that $R_A \otimes_A k = R$. Finally, by the local criterion for flatness, we see that R_A is flat over A .

Let $R_A = P_A/I_A$ be a lifting of R over A . Lift $f_i \in I$ to $h_i \in I_A$. By Nakayama, these generate I_A , so we obtain a free resolution

$$0 \rightarrow P_A^{r-1} \xrightarrow{(g'_{ij})} P_A^r \xrightarrow{(h_i)} P_A \rightarrow R_A \rightarrow 0,$$

where g'_{ij} lift the g_{ij} . However, we already have a lift

$$0 \rightarrow P_{\Lambda}^{r-1} \xrightarrow{(g'_{ij})} P_{\Lambda}^r \xrightarrow{(f'_i)} P_{\Lambda} \rightarrow R'_{\Lambda} \rightarrow 0,$$

and so we must show $R_{\Lambda} = R'_{\Lambda}$. But we know that the ideals $I_{\Lambda} = (h_1, \dots, h_r)$ and $I'_{\Lambda} = (f'_1, \dots, f'_r)$ are isomorphic as P_{Λ} -modules. But then if we restrict this isomorphism to $\mathbb{A}_{\Lambda}^n \setminus \text{supp } B$, we obtain a unit in $H^0(\mathbb{A}_{\Lambda}^n \setminus \text{supp } B, \mathcal{O}_{\mathbb{A}_{\Lambda}^n})$. Because functions extend over codimension 2, we have $H^0(\mathbb{A}_{\Lambda}^n \setminus \text{supp } B, \mathcal{O}_{\mathbb{A}_{\Lambda}^n}) = P_{\Lambda}$, so this is a global unit. This gives the desired result. \square

This result holds if we replace $A \rightarrow k$ with any square-zero extension of Artinian local rings $B \rightarrow A$ and P, P_{Λ} with flat things, and so we see that (embedded) deformations of codimension 2 Cohen-Macaulay subschemes of \mathbb{A}^n are unobstructed. In particular, any dimension 0 closed subscheme $Z \subset \mathbb{A}^2$ is automatically Cohen-Macaulay (because it is dimension 0), so its embedded deformations are unobstructed. By some cohomological argument, the tangent space to $\text{Hilb}(\mathbb{A}^2, n)$ is isomorphic to $\text{Hom}(R, R)$ and has dimension $2n$, so

3. AN OBSTRUCTED DEFORMATION

Let $R = k[x, y, z]/(z^2, xy, xz, yz)$. Note that this scheme has an embedded point at the origin, so in particular it is **not** Cohen-Macaulay.




FIGURE 1. Drawing of $\text{Spec } R$

We will study embedded deformations of $\text{Spec } R$ and see that they are obstructed. In particular, we will choose two deformations of R over $k[\epsilon]$ that cannot be simultaneously lifted. We claim that a complete set of relations (using the ordering (xy, xz, yz, z^2) for the generators of I) is given by the matrix

$$G = \begin{pmatrix} z & -y & 0 & 0 \\ z & 0 & -x & 0 \\ 0 & z & 0 & -x \\ 0 & 0 & z & -y \end{pmatrix}.$$

Now a first-order deformation of $\text{Spec } R$ is given by lifting (xy, xz, yz, z^2) over $k[\varepsilon]$, and the first candidate is to consider $I_{\varepsilon_1} = (xy + \varepsilon_1 y, xz, yz, z^2)$. Then we note that

$$G \begin{pmatrix} xy + \varepsilon_1 y \\ xz \\ yz \\ z^2 \end{pmatrix} = \varepsilon_1 \begin{pmatrix} yz \\ yz \\ 0 \\ 0 \end{pmatrix},$$

and we can lift G to kill this vector with the matrix

$$G_{\varepsilon_1} = \begin{pmatrix} z & -y & -\varepsilon_1 & 0 \\ z & 0 & -x - \varepsilon_1 & 0 \\ 0 & z & 0 & -x \\ 0 & 0 & z & -y \end{pmatrix} = G + \begin{pmatrix} 0 & 0 & -\varepsilon_1 & 0 \\ 0 & 0 & -\varepsilon_1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} =: G + G_1.$$

Next consider the deformation given by $I_{\varepsilon_2} = (xy, xz, yz + \varepsilon_2 z, z^2)$. We note that

$$G \begin{pmatrix} xy \\ xz \\ yz + \varepsilon_2 z \\ z^2 \end{pmatrix} = \varepsilon_2 \begin{pmatrix} 0 \\ -xz \\ 0 \\ z^2 \end{pmatrix},$$

and we can lift G to kill this vector with the matrix

$$G_{\varepsilon_2} = \begin{pmatrix} z & -y & 0 & 0 \\ z & \varepsilon_2 & -x & 0 \\ 0 & z & 0 & -x \\ 0 & 0 & z & -y - \varepsilon_2 \end{pmatrix} = G + \begin{pmatrix} 0 & \varepsilon_2 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\varepsilon_2 \\ 0 & 0 & 0 & 0 \end{pmatrix} =: G + G_2.$$

Now we consider $I_{\varepsilon_1^2, \varepsilon_2^2, \varepsilon_1 \varepsilon_2} = (xy + \varepsilon_1 y, xz, yz + \varepsilon_2 z, z^2)$ and attempt to lift this deformation to $k[\varepsilon_1, \varepsilon_2]/(\varepsilon_1^2, \varepsilon_2^2)$. Note that

$$\begin{aligned} (G + G_1 + G_2) \begin{pmatrix} xy + \varepsilon_1 y \\ xz \\ yz + \varepsilon_2 z \\ z^2 \end{pmatrix} &= \begin{pmatrix} z & -y & -\varepsilon_1 & 0 \\ z & \varepsilon_2 & -x - \varepsilon_1 & 0 \\ 0 & z & 0 & -x \\ 0 & 0 & z & -y - \varepsilon_2 \end{pmatrix} \begin{pmatrix} xy + \varepsilon_1 y \\ xz \\ yz + \varepsilon_2 z \\ z^2 \end{pmatrix} \\ &= \varepsilon_1 \varepsilon_2 \begin{pmatrix} -z \\ -z \\ 0 \\ 0 \end{pmatrix}, \end{aligned}$$

and clearly $z \notin I$, so in fact we cannot lift this deformation to $k[\varepsilon_1, \varepsilon_2]/(\varepsilon_1^2, \varepsilon_2^2)$. This proves obstructedness.