NOTES ON THE B-MODEL

LECTURES ON BILIBILI BY SI LI
NOTES BY PATRICK LEI

Abstract. These are notes taken based on the Bilibili minicourse on the B-model given by Si Li in Spring 2025 at the Chinese University of Hong Kong. The minicourse consisted of 12 lectures.

Contents

1. Introduction 1
2. Calabi-Yau geometry 5
3. Period map 8
4. Extended period map 11
5. Classical BCOV action 14
6. Hodge structure and primitive forms 18
7. Classical master equation 22
8. Quantum master equation 25
9. Homotopic renormalization 27
10. Quantum BCOV theory 27
11. Landau-Ginzburg B-model 27
12. Open-closed B-model 27

1. Introduction

The B-model has its origins in *mirror symmetry*, which is a duality of two dimensional superconformal field theories (SCFTs) in physics. These describe the geometry of a mapping space from a surface Σ to a target space X (see Figure 1). The duality gives mathematical predictions which relate symplectic geometry (the A-model) to complex geometry (the B-model).

There is a duality

$$\begin{array}{ccc} \int_{\mathrm{Map}(\Sigma,X)} (\text{A-model}) & \overset{\text{Fourier T-duality}}{\rightsquigarrow} & \int_{\mathrm{Map}(\Sigma,X)} (\text{B-model}) \\ \Big\downarrow \text{localize} & & \Big\downarrow \text{localize} \\ \int_{\mathrm{Hol}(\Sigma,X)} \cdots & & \int_{\mathrm{Const}(\Sigma,X)} \cdots \\ \Big\Downarrow & & \Big\Downarrow \\ \text{Gromov-Witten type} & & \text{Hodge-type} \end{array}$$

Date: February 2026.




FIGURE 1. Map from a surface Σ to a target space X.

where both sides involve some kind of infinite-dimensional geometry. However, we can reduce this to a finite-dimensional geometry by a form of supersymmetric localization, which we will not explain. The interested reader can refer to the speaker's lectures at the Perimeter Institute on supersymmetry and TQFTs, which are also on Bilibili.

There is a string description via a worldsheet theory, where we consider maps as in Figure 2. This theory is called *string field theory*. This describes string theory using the language of quantum field theory. The correspondence goes as in Table 1.






FIGURE 2. Map in the worldsheet theory X.

TABLE 1. Idea of string field theory

String Fock space	String field theory
$\lvert\psi\rangle$	ψ
Dynamics	$S[\psi]$ String field action
	

We outline some historical developments in both mathematics and physics related to the B-model.

- In 1992, Zwiebach described a general structure for closed string field theory and discovered that it is described by a string action which contains infinitely many vertices and satisfies the *BV master equation*, something we will describe later in the language of the B-model. In particular, if we consider the genus 0 closed string field, we will discover a homotopy Lie algebra or an L_∞-algebra.
- In 1997, Zwiebach extended this theory to the open string field, which is described by an A_∞-algebra. When coupled with the closed string field, it satisfies the *open-closed master equation*, which is described via the moduli of bordered Riemann surfaces.

We will consider the B-model from the perspective of string field theory. This is usually referred to as the B-twisted topological field theory, and when coupled to gravity, it becomes the topological string theory. It is localized around constant maps, and near each constant map, we see the "local" string field action. The correspondence goes as in Figure 3.




FIGURE 3. Localization of the B-model.

Here are some developments in the history of describing this new action:

- Witten described the A-model open string field using Chern-Simons and instanton theories in 1992. As for the B-model open string field, he described it as holomorphic Chern-Simons theory. Finally, he speculated that the theory is actually UV-finite, which we know now is true by the work of M. Wang and J. Yan.
- The famous work of Bershadsky-Cecotti-Ooguri-Vafa in 1994 described the B-model closed string field theory on a Calabi-Yau threefold as a gauge theory, which they call *Kodaira-Spencer gravity*. This has some interesting features:
  - At the classical level, the equation of motion describes deformation of complex structures on the Calabi-Yau threefold. By Yau's proof of the Calabi conjecture, this leads to a family of solutions which are Ricci-flat metrics.
  - The leading cubic vertex of Zwiebach's string action in the topological B-model admits a full description.

- There is a construction by Barannikov-Kontsevich which constructs the Frobenius manifold (which encodes classical genus-zero data) of this theory via the geometry of polyvector fields, which will be discussed in the next few lectures.
- In 2012, the speaker and Kevin Costello gave a mathematical definition of the full string action for the topological B-model on a Calabi-Yau manifold of any dimension and proved that it satisfies the full BV master equation. They refer to this as *BCOV theory*.
- In 2015/16, the same authors determined the B-model open-closed quantum coupling in the large N limit and from this perspective they worked out a version of twisted supergravity. Most of the lectures will explain the speaker's work with Costello.

Convention 1.1. We will now fix some conventions. We will work with the category of $\mathbb{Z}$-graded vector spaces

$$V = \bigoplus_{m\in\mathbb{Z}} V_m.$$

For any $a \in V_m$, we will denote its degree by $\deg(a) = |a| = m$. We will denote degree shifts by

$$(V[n])_m = V_{m+n},$$

or namely, shifting to the left. Finally, we will discuss some conventions on tensors. We will denote the symmetric power by

$$\mathrm{Sym}^n(V) = V^{\otimes n}/(a\otimes b \sim (-1)^{|a||b|} b\otimes a)$$

and the exterior powers by

$$\Lambda^n(V) = V^{\otimes n}/(a\otimes b \sim -(-1)^{|a||b|} b\otimes a).$$

These are sometimes referred to as the *graded symmetric product* and the *graded skew-symmetric product*. Finally, we will consider the symmetric algebra (or polynomial algebra)

$$\mathrm{Sym}(V) = \bigoplus_m \mathrm{Sym}^m(V)$$

and its completion (a power series ring)

$$\widehat{\mathrm{Sym}}(V) = \prod_m \mathrm{Sym}^m(V).$$

For a graded algebra A, we will always use $[-,-]$ to refer to the graded commutator, which is defined by

$$[a,b] = ab - (-1)^{|a||b|} ba.$$

Finally, if X is a smooth manifold X, we will write

$$\Omega^\bullet(X) = \bigoplus_k \Omega^k(X)$$

to denote smooth differential forms on X and denote the de Rham differential by d. If X is a complex manifold, we can decompose

$$\Omega^k(X) = \bigoplus_{p+q=k} \Omega^{p,q}(X)$$

and decompose the differential as $\mathrm{d} = \bar\partial + \partial$.

Exercise 1.2. Show that

$$\mathrm{Sym}^n(V[1]) \simeq \Lambda^n(V)[n].$$

2. Calabi-Yau geometry

In this section, we will build some foundations for Calabi-Yau geometry. Let X be a complex manifold of dimension $\dim_{\mathbb{C}} = d$. We will denote the holomorphic tangent bundle by T_X. We will denote polyvector fields by

$$\mathrm{PV}^{i,j}(X) := \Omega^{0,j}(X, \Lambda^i T_X)$$

and the space of all polyvector fields by

$$\mathrm{PV}(X) = \bigoplus_{i,j} \mathrm{PV}^{i,j}(X).$$

Locally, we can write $\mu \in \mathrm{PV}^{i,j}(X)$ can be written as

$$\mu = \sum_{\substack{|I|=i \\ |J|=j}} \mu_J^I \, \mathrm{d}\bar{z}^I \otimes \partial_{z^I},$$

where $I = \{k_1, \dots, k_i\}$ and $J = \{s_1, \dots, s_j\}$ are multi-indices. Here, the index notation looks like

$$\mathrm{d}\bar{z}^J = \mathrm{d}\bar{z}^{s_1} \wedge \cdots \wedge \mathrm{d}\bar{z}^{s_j} \qquad \text{and} \qquad \partial_{z^I} = \partial_{z^{k_1}} \wedge \cdots \wedge \partial_{z^{k_i}}.$$

These look like differential forms, but are in fact subtly different. All μ_J^I are smooth and skew-symmetric for indices in both I and in J.

There is a natural (wedge) product

$$\mathrm{PV}^{i_1,j_1} \otimes \mathrm{PV}^{i_2,j_2} \xrightarrow{\wedge} \mathrm{PV}^{i_1+i_2,j_1+j_2}$$

which makes $\mathrm{PV}(X)$ a bi-graded algebra. We will define the total degree by $\left|\mu\right| = i + j$ for $\mu \in \mathrm{PV}^{i,j}(X)$. This makes $(\mathrm{PV}(X), \wedge)$ a graded-commutative algebra, namely that $\alpha \wedge \beta = (-1)^{|\alpha||\beta|} \beta \wedge \alpha$.

We can also extend the $\bar{\partial}$-operator to polyvector fields, which has degree 1 and goes as

$$\bar{\partial} \colon \mathrm{PV}^{i,j}(X) \to \mathrm{PV}^{i,j+1}(X).$$

This is in fact a derivation which satisfies the graded Leibniz rule

$$\bar{\partial}(\alpha \wedge \beta) = (\bar{\partial}\alpha) \wedge \beta + (-1)^{|\alpha|} \alpha \wedge \bar{\partial}\beta.$$

There are many definitions for Calabi-Yau manifolds, so we will choose one.

Definition 2.1. A *Calabi-Yau manifold* is a pair (X, Ω_X) where X is a complex manifold and Ω_X is a nowhere-vanishing holomorphic volume form on X.

In local holomorphic coordinates $\left\{z^1, \dots, z^d\right\}$ on X, we can write

$$\Omega_X = \varphi(z) \, \mathrm{d}z^1 \wedge \cdots \wedge \mathrm{d}z^d,$$

where $\varphi(z)$ is a nowhere-vanishing holomorphic function. The Calabi-Yau form induces a linear isomorphism

$$\mathrm{PV}^{i,j} \xrightarrow{\lrcorner \Omega_X} \Omega^{d-i,j} \qquad \mu \mapsto \mu \lrcorner \Omega_X.$$

We now consider the *divergence operator*

$$\partial_\Omega \colon \mathrm{PV}^{i,j} \to \mathrm{PV}^{i-1,j}$$

with respect to the volume form Ω_X, which is defined using the commutative diagram

$$\begin{array}{ccc} \mathrm{PV}^{i,j} & \xrightarrow{\partial_\Omega} & \mathrm{PV}^{i-1,j} \\ {\scriptstyle \Omega_X}\downarrow{\scriptstyle \sim} & & {\scriptstyle \sim}\uparrow{\scriptstyle \Omega_X} \\ \Omega^{d-i,j} & \xrightarrow{\partial} & \Omega^{d-i+1,j}. \end{array}$$

Note that this depends on the choice of volume form. Unfortunately, we will abuse notation and write ∂ for ∂_Ω and whether we mean the divergence operator or the holomorphic de Rham operator will be clear from context. Unfortunately, the divergence operator is **not** a derivation. The failure is described by the tensor

$$\{\alpha, \beta\} = \partial(\alpha \wedge \beta) - (\partial\alpha) \wedge \beta - (-1)^{|\alpha|}\alpha \wedge (\partial\beta).$$

This bracket is known in geometry and is related to the Schouten-Nijenhuis bracket $[-,-]_{\mathrm{SN}}$ by the sign

$$[\alpha, \beta]_{\mathrm{SN}} = -(-1)^{|\alpha|}\{\alpha, \beta\}.$$

In mathematics, this is sometimes called the Bogomolov-Tian-Todorov lemma and in physics it is related to the BV algebra. In fact, the difference between the Schouten-Nijenhuis bracket and the BV bracket is related to shifting the degree by 1. The BV bracket has degree given by

$$\{-,-\} \colon \mathrm{PV}^{i_1,j_1} \otimes \mathrm{PV}^{i_2,j_2} \to \mathrm{PV}^{i_1+i_2-1,j_1+j_2}.$$

It is in fact intrinsically defined independent of the choice of volume form (or divergence operator).

Remark 2.2. The lifting of $\{-,-\}$ to the volume form Ω_X or the divergence operator ∂ is related to *BV quantization*.

Exercise 2.3. Show that the BV bracket satisfies graded symmetry

$$\{\alpha, \beta\} = -(-1)^{(|\alpha|)(|\beta|)}\{\beta, \alpha\},$$

the Leibniz relation

$$\{\alpha, \beta \wedge \gamma\} = \{\alpha, \beta\} \wedge \gamma + (-1)^{|\beta||\gamma|}\{\alpha, \gamma\} \wedge \beta,$$

and the graded Jacobi identity

$$\{\{\alpha, \beta\}, \gamma\} = (-1)^{|\alpha|}\{\alpha, \{\beta, \gamma\}\} - (-1)^{(|\alpha|+1)|\beta|}\{\beta, \{\alpha, \gamma\}\}.$$

Remark 2.4. The algebra $(\mathrm{PV}, \bar{\partial}, \partial, \{-,-\})$ is a fundamental example of a *differential graded Gerstenhaber BV algebra* and exists in a large class of examples.

We can now define a *trace map*

$$\mathrm{Tr} \colon \mathrm{PV}(X) \to \mathbb{C} \qquad \mathrm{Tr}(\mu) \coloneqq \int_X (\mu \lrcorner \Omega_X) \wedge \Omega_X.$$

This trace is only nontrivial for $\mathrm{PV}^{d,d}$ and will be used later to discuss higher residue pairings.

Exercise 2.5. Show that the trace map is compatible with the the two differentials via the identities

$$\mathrm{Tr}((\bar\partial\alpha)\wedge\beta) = -(-1)^{|\alpha|}\,\mathrm{Tr}(\alpha\wedge\bar\partial\beta)$$

and

$$\mathrm{Tr}((\partial\alpha)\wedge\beta) = (-1)^{|\alpha|}\,\mathrm{Tr}(\alpha\wedge\partial\beta).$$

We will now let X be a compact Kähler manifold. The deformation space of complex structures on X is locally given by

$$\mathscr{M}^{\mathrm{cx}} = \left\{\mu\in \mathrm{PV}^{1,1}(X)\middle|\|\mu\|<\varepsilon, \bar\partial\mu+\frac{1}{2}\{\mu,\mu\}=0, \bar\partial^*\mu=0\right\}.$$

Here, the final condition is a gauge-fixing condition and $\bar\partial\mu+\frac{1}{2}\{\mu,\mu\}=0$ is known as the *Maurer-Cartan equation*.

Theorem 2.6 (Bogomolov-Tian-Todorov)**.** *Let X be a compact Kähler Calabi-Yau manifold. Then the deformation space $\mathscr{M}^{\mathrm{cx}}$ is smooth and has dimension*

$$\dim_{\mathbb{C}}\mathscr{M}^{\mathrm{cx}} = h^1(X,T_X).$$

In general, we can consider the *extended moduli space* $\mathscr{M}$ by solving

$$\bar\partial\mu+\frac{1}{2}\{\mu,\mu\}=0$$

modulo gauge for all types of polyvector fields. We can view the original moduli space as a submanifold of the extended one.

Theorem 2.7 (Barannikov-Kontsevich)**.** *The extended moduli space $\mathscr{M}$ is smooth.*

Proof. Consider the divergence-free polyvector fields $\ker\partial\subset\mathrm{PV}$. Given $\alpha,\beta\in\ker\partial$, we consider their BV bracket

$$\begin{aligned}\{\alpha,\beta\} &= \partial(\alpha\wedge\beta)-(\partial\alpha)\wedge\beta-(-1)^{|\alpha|}\alpha\wedge(\partial\beta)\\ &= \partial(\alpha\wedge\beta),\end{aligned}$$

which is ∂-exact. Therefore,

$$\{-,-\}\colon\ \ker\partial\otimes\ker\partial\to\mathrm{Im}\partial.$$

This fact will be crucial.

The complex $(\mathrm{PV},\bar\partial,\{-,-\})$ is actually a (shifted) dg Lie algbera, and we consider the inclusion

$$j\colon(\ker\partial,\bar\partial,\{-,-\})\hookrightarrow(\mathrm{PV},\bar\partial,\{-,-\}).$$

This is actually a morphism of dg Lie algebras and is in fact a quasi-isomorphism. If we allow $\mathbb{H}\subset\mathrm{PV}(X)$ be the harmonic elements, then extending Hodge theory gives us

$$\mathbb{H}\simeq\ker\partial/\mathrm{Im}\partial\simeq\ker\bar\partial/\mathrm{Im}\bar\partial$$

by the Kähler condition. Now define the projection

$$\Pi\colon\ \ker\partial\to\mathbb{H}.$$

By the $\partial\bar\partial$-lemma, we see that $\bar\partial(\ker\partial)\subset\mathrm{Im}\partial$, so we can consider the map

$$(\ker\partial,\bar\partial,\{-,-\})\xrightarrow{\Pi}(\mathbb{H},0,0),$$

which is again a morphism of dg Lie algebras. Now we have the zig-zag

$$\begin{array}{ccccc} & & (\ker\partial,\bar\partial,\{-,-\}) & & \\ & \swarrow^{j} & & \searrow^{\Pi} & \\ (\mathrm{PV},\bar\partial,\{-,-\}) & & & & (\mathbb{H},0,0), \end{array}$$

where both maps are quasi-isomorphisms. By some abstract nonsense, solving the Maurer-Cartan up to gauge is invariant under quasi-isomorphisms, and solving the Maurer-Cartan equation in $(\mathbb{H},0,0)$ is trivial. Therefore we have $\mathscr{M} \simeq \mathbb{H}$. □

3. Period map

Before we move to the B-model, we need to discuss the period map, which will allow us to write down action functionals. Let X, Ω_X be a compact Kähler Calabi-Yau manifold. Given $\mu \in \mathrm{PV}^{1,1}(X)$ satisfying the Maurer-Cartan equation

$$\bar\partial\mu + \frac{1}{2}\{\mu,\mu\} = 0,$$

we will use $[X_\mu] \in \mathscr{M}^{\mathrm{cx}}$ to denote the complex manifold specified by the deformation μ. Assume that $\{z^i\}$ are local holomorphic coordinates on X. We can write

$$\mu = \sum_{i,j} \mu^i_{\bar\jmath}\, \mathrm{d}\bar z^j \otimes \partial_{z^i}.$$

Then we can define new holomorphic 1-forms by

$$\begin{aligned}(\mathrm{d}z^i + \mu \lrcorner\, \mathrm{d}z^i) &= \mathrm{d}z^i + \sum_j \mu^i_{\bar\jmath}\, \mathrm{d}\bar z^j \\ &= e^{\mu} \lrcorner\, \mathrm{d}z^i,\end{aligned}$$

and these form a local basis of $(1,0)$-forms on X_μ.

Let $\mathscr{H}^d$ denote the complex vector bundle over $\mathscr{M}^{\mathrm{cx}}$ whose fiber at $[X_\mu]$ is the de Rham cohomology $H^d(X_\mu)$. This bundle admits a flat connection, called the *Gauss-Manin connection*, and denoted by ∇^{GM}, which identifies the nearby fibers to the central one as

$$H^d(X_\mu) \simeq H^d(X).$$

There is some global issue with monodromy, but locally the identification is canonical. We now define the holomorphic subbundles (the Hodge filtration) $\mathscr{F}^p\mathscr{H}^d \subset \mathscr{H}^d$ with fibers

$$\mathscr{F}^p\mathscr{H}^d|_{[X_\mu]} = \bigoplus_{k\geq p} H^{k,d-k}(X_\mu).$$

In particular, there is the line bundle $L = \mathscr{F}^d\mathscr{H}^d$ which picks up the holomorphic volume form on X_μ. In physics, this is known as the *vacuum line bundle*. Later, we will see that invariants of the B-model are actually sections of powers of this line bundle.

The period map is locally given by the map

$$\mathscr{M}^{\mathrm{cx}} \hookrightarrow \mathbb{P}(H^d(X,\mathbb{C})) \qquad [X_\mu] \mapsto [\Omega_{X_\mu}].$$

Here, we need to use the Gauss-Manin connection to identify the cohomology of X_μ with that of X. The period map is actually a local embedding (the local Torelli

theorem). Sometimes it is helpful to remember the actual volume form, and if we consider the $\mathbb{C}^\times$-bundle

$$\hat{\mathscr{M}}^{\mathrm{cx}} = \mathrm{Tot}(L \smallsetminus \{0\})$$

over $\mathscr{M}^{\mathrm{cx}}$, then a point here represents a pair (X, Ω_X), so we have defined the local moduli of deformations of a pair (X, Ω_X). This allows us to lift the period map to an embedding

$$\hat{\mathscr{M}}^{\mathrm{cx}} \hookrightarrow H^d(X, \mathbb{C}) \qquad (X_\mu, \Omega_{X_\mu}) \mapsto [\Omega_{X_\mu}].$$

Our goal is to understand the geometry of the period map, which will allow us to understand the action functional of the B-model. We will begin with the case of Calabi-Yau threefolds, which is classical, and then move to a more modern perspective. Let (X, Ω_X) be a Calabi-Yau threefold. Then $H^3(X, \mathbb{C})$ has a symplectic pairing given by

$$\omega(\alpha, \beta) = \int_X \alpha \wedge \beta.$$

Considering the Hodge filtration, the subspace

$$F^2H^3 = H^{3,0} \oplus H^{2,1} \subset H^3$$

is a Lagrangian subspace. Over the moduli space $\mathscr{M}^{\mathrm{cx}}$ ω defines a fiberwise symplectic pairing on $\mathscr{H}^3$ which is flat with respect to the Gauss-Manin connection. Then we can lift the Hodge filtration to a subbundle

$$\mathscr{F}^2\mathscr{H}^3 \subset \mathscr{H}^3,$$

which gives a holomorphic family of Lagrangian subspaces.

Proposition 3.1. *The embedding $\hat{\mathscr{M}}^{\mathrm{cx}} \hookrightarrow H^3$ defines a Lagrangian submanifold.*

Proof. Let Ω_s be a holomorphic family of Calabi-Yau forms parameterized by s. By Griffiths transversality, we see that

$$\nabla^{\mathrm{GM}}\Omega_s \subset \mathscr{F}^2\mathscr{H}^3.$$

Every fiber is a linear Lagrangian subspace, so we are done. □

From this data, we can cook up B-model invariants as follows. Choose a Lagrangian splitting $\mathscr{L} \subset H^3(X, \mathbb{C})$ be a transversal linear Lagrangian subspace such that

$$H^3 = F^2H^3 \oplus \mathscr{L}$$

is an isotropic splitting. This allows us to identify

$$H^3 \cong T^*F^2,$$

where we use F^2 to refer to F^2H^3. Then $\hat{\mathscr{M}}^{\mathrm{cx}}$ is a Lagrangian submanifold as in Figure 4. Therefore, there is some function $\hat{F}_0 \in \mathscr{O}(F^2)$ such that $\hat{\mathscr{M}}^{\mathrm{cx}}$ is the graph of $\mathrm{d}\hat{F}_0$.

Example 3.2. Choose $\mathscr{L} = H^{1,2} \oplus H^{0,3}$ (the complex conjugate splitting). Physicists would now write down some special geometry and prepotentials. We parametrize $\hat{\mathscr{M}}^{\mathrm{cx}}$ by

$$\hat{\mathscr{M}}^{\mathrm{cx}} = \left\{(t^0, t^i\partial_{t^i}\hat{F}_0, \partial_{t^0}\hat{F}_0)\right\}.$$




FIGURE 4. The moduli space $\hat{\mathscr{M}}^{\mathrm{cx}}$ as a Lagrangian submanifold in $H^3 \simeq T^*F^2$.

Because $\mathscr{M}^{\mathrm{cx}} = \hat{\mathscr{M}}^{\mathrm{cx}}/\mathbb{C}^\times$, we see that $\hat{F}_0$ has homogeneous degree 2. The local coordinates on $\mathscr{M}^{\mathrm{cx}}$ are given by $\tau^i = \frac{t^i}{t^0}$. Therefore, we can write

$$\hat{F}_0 = (t_0)^2 \mathscr{F}_0(\tau^i).$$

Geometrically, $\hat{F}_0$ actually defines a local section of $L^{\otimes -2}$, where L is the vacuum line bundle. We can now reparaterize the extended moduli space by

$$\hat{\mathscr{M}}^{\mathrm{cx}} = \left\{ t^0 (1, \tau^i \partial_{\tau^i} \mathscr{F}_0, 2\mathscr{F}_0 - \tau^i \partial_{\tau^i} \mathscr{F}_0) \right\}.$$

The factor inside the parentheses is called the *primitive period map*. We now obtain a particular choice of holomorphic volume form on the moduli space given by

$$[\Omega_\tau] = (1, \tau^i, \partial_{\tau^i} \mathscr{F}_0, 2\mathscr{F}_0 - \tau^i \partial_{\tau^i} \mathscr{F}_0).$$

We are now finally able to calculate the *Yukawa coupling* for the moduli space by

$$\begin{aligned} C_{ijk}(\tau) &= \int \nabla^{\mathrm{GM}}_{\partial_{\tau^i}} [\Omega_\tau] \wedge \nabla^{\mathrm{GM}}_{\partial_{\tau^j}} \nabla^{\mathrm{GM}}_{\partial_{\tau^k}} [\Omega_\tau] \\ &= \partial_{\tau^i} \partial_{\tau^j} \partial_{\tau^k} \mathscr{F}_0(\tau). \end{aligned}$$

In the physics literature, $\mathscr{F}_0(\tau)$ is called the *prepotential*.

Exercise 3.3. Prove the formula for the Yukawa coupling in the previous example.

Before we proceed, we need to describe the deformation of the pair (X, Ω_X). We know that deformations of X are given by $\mu \in \mathrm{PV}^{1,1}(X)$ satisfying the Maurer-Cartan

equation

$$\bar{\partial}\mu + \frac{1}{2}\{\mu, \mu\} = 0.$$

In the complex structure X_μ specified by μ, the elements

$$e^{\mu} \lrcorner \, \mathrm{d}z^i = \mathrm{d}z^i + \mu \lrcorner \, \mathrm{d}z^i$$

form a new basis of $(1,0)$-forms on X_μ.

Lemma 3.4. *We have*

$$e^{\mu} \lrcorner (\alpha \wedge \beta) = (e^{\mu} \lrcorner \alpha) \wedge (e^{\mu} \lrcorner \beta)$$

for any differential forms α, β.

The lemma now implies that

$$e^{\mu} \lrcorner (\mathrm{d}z^1 \wedge \cdots \wedge \mathrm{d}z^d)$$

is of type $(d,0)$ with respect to X_μ. Applying this globally, the form

$$e^{\mu} \lrcorner \Omega_X$$

is of type $(d,0)$ in X_μ. Therefore, there exists a smooth function ρ such that

$$e^{\rho} e^{\mu} \lrcorner \Omega_X$$

is a holomorphic volume form on X_μ. Because the type is $(d,0)$, then the form is holomorphic if and only if it is closed, or in other words that $\mathrm{d}(e^{\rho} e^{\mu} \lrcorner \Omega_X) = 0$. This gives us the two equations

$$\begin{cases} \bar{\partial}\mu + \frac{1}{2}\{\mu, \mu\} = 0 & \rightsquigarrow \text{deform to } X_\mu \\ \bar{\partial}\rho + \partial\mu + \{\mu, \rho\} = 0 & \rightsquigarrow \text{deform to } \Omega_{X_\mu} \end{cases}$$

which govern deformations of the pair (X, Ω_X). These equations were studied by Barannikov in his work on variations of semi-infinite Hodge structures.

4. Extended period map

Introduce a formal parameter t of degree 2, which will play the role of gravitational descendents and consider the combined differential

$$Q = \bar{\partial} + t\partial.$$

We can rewrite our two equations as the single equation

$$Q\hat{\mu} + \frac{1}{2}\{\hat{\mu}, \hat{\mu}\} = 0,$$

where we have defined

$$\hat{\mu} = \mu + t\rho.$$

Introduce a new (shifted) dg Lie algebra

$$(\mathrm{PV}(X)[\![t]\!], Q = \bar{\partial} + t\partial, \{-,-\}).$$

Remark 4.1. The data of $(\mathrm{PV}(X)[\![t]\!], Q)$ can be viewed as a homological replacement of volume-preserving (divergence-free) vector fields $(\ker \partial, \bar{\partial})$. We may now write the complex

$$\ker \partial \to \mathrm{PV} \xrightarrow{t\partial} t\mathrm{PV} \xrightarrow{t\partial} t^2\mathrm{PV} \to \cdots$$

which behaves badly, so we replace it with $(\mathrm{PV}[\![t]\!], Q)$.

Remark 4.2. The variable t plays the role of gravitational descendants (namely ψ-classes on the A-side) in the B-model.

Remark 4.3. If $(\mathrm{PV}, \bar{\partial}, \{-.-\})$ gives extended moduli for X, then $(\mathrm{PV}[\![t]\!], Q, \{-,-\})$ gives extended moduli for the Calabi-Yau geometry.

Remark 4.4. The variable t can be viewed as an equivariant parameter for rotation on the worldsheet in the worldsheet B-model.

Question 4.5. *Can we express the above equations as the variational critical locus of some action functional?*

One reason we would be interested in this question is because this is the usual approach to classical physics, and then we can quantize these using path integrals. This is unfortunately highly nontrivial, and on a Calabi-Yau threefold, BCOV work with some $\mu \in \ker \partial$. They found that the Maurer-Cartan equation

$$\bar{\partial}\mu + \frac{1}{2}\{\mu, \mu\} = 0$$

is equivalent to $\frac{\delta \mathrm{KS}}{\delta \mu} = 0$ for the Kodaira-Spencer action KS.

Warning 4.6. This approach unfortunately violates locality for various reasons:

- $\ker \partial$ is not a module over $C^\infty(X)$;
- One of the terms in KS is nonlocal.

Consider the Calabi-Yau geometry and let $\mu \in \mathrm{PV}(X)[\![t]\!]$. We can ask for an action functional whose critical points are solutions of

$$Q\mu + \frac{1}{2}\{\mu, \mu\} = 0.$$

Unfortunately, the speaker was unable to find this, but there is some magic involved. If we perform an L_∞ transformation, this equation becomes

$$Q\mu + \sum_{k \geq 2} \frac{1}{k!} \ell_k(\mu^{\otimes k}) = 0,$$

and for this equation there is a BCOV action which was found by Costello and the speaker. The L_∞ transformation is produced directly from the period map.

Definition 4.7. Define

$$\mathrm{S}(X) := \mathrm{PV}(X)(\!(t)\!)[2],$$

where the shift makes $\mu + t\rho$ have degree 0. Then define

$$\mathrm{S}_+(X) := \mathrm{PV}(X)[\![t]\!][2] \qquad \text{and} \qquad \mathrm{S}_-(X) := t^{-1}\mathrm{PV}(X)[t^{-1}][2].$$

This makes $\mathrm{S}(X) = \mathrm{S}_+(X) \oplus \mathrm{S}_-(X)$, and once we define a symplectic structure it will be a splitting by isotropic subspaces. However, Q does **not** preserve the splitting, but this will lead to the classical master equation.

We define a t-sesquilinear pairing on $\mathrm{S}(X)$ (inspired by the higher residue pairing of Kyoji Saito)

$$\omega \colon \mathrm{S}(X) \otimes \mathrm{S}(X) \to \mathbb{C}$$

by the formula

$$\omega(f(t)\alpha, g(t)\beta) := \mathrm{Tr}\, \alpha\beta \, \mathrm{Res}_{t=0} f(t)g(-t)\, \mathrm{d}t.$$

Observe that

$$\begin{aligned}\omega(g(t)\beta, f(t)\alpha) &= \operatorname{Tr}\beta\alpha \operatorname{Res}_{t=0} g(t)f(-t)\,\mathrm{d}t \\ &= -(-1)^{|\alpha||\beta|}\operatorname{Tr}\alpha\beta \operatorname{Res}_{t=0} f(t)g(-t)\,\mathrm{d}t \\ &= -(-1)^{|\alpha||\beta|}\omega(f(t)\alpha, g(t)\beta).\end{aligned}$$

If we calculate the degrees, then ω is only nontrivial when

$$\begin{aligned}\deg(f(t)\alpha) + \deg(g(t)\beta) &= \deg(\alpha\wedge\beta) + \deg f(t) + \deg g(t) \\ &= 2d-2,\end{aligned}$$

so if we take the original shift by 2 into account, we have proved the following result.

Proposition 4.8. *The space* $(\mathrm{S}(X), \omega)$ *has a* $(6-2d)$*-shifted symplectic structure.*

Remark 4.9. For a Calabi-Yau threefold X, we have the splitting

$$\Omega^3 = \Omega^{3,0}\oplus\Omega^{2,1}\oplus\Omega^{1,2}\oplus\Omega^{0,3},$$

and the we will be able to identify this with

$$t\mathrm{PV}^{0,0}\oplus\mathrm{PV}^{1,1}\oplus t^{-1}\mathrm{PV}^{2,2}\oplus t^{-2}\mathrm{PV}^{3,3}$$

via contraction with Ω_X, which we will call Γ. The higher residue pairing will be identified with the usual pairing on differential forms. We should note however that this only works for threefolds.

Proposition 4.10. *The operator* Q *is* (*graded*) *skew-symmetric with respect to* ω*.*

Proof. The proof is left as an exercise. The key point is to study the operators $\bar\partial$ and ∂. □

We now consider the (cochain) period map

$$\mathscr{P}\colon \mathrm{S}_+(X)\to\mathrm{S}(X)\qquad \mu\mapsto t\Big(1-e^{\frac{\mu}{t}}\Big).$$

Remark 4.11. This $\mathscr{P}$ is understood as a map on formal schemes. For any Artinian ring $(R,\mathfrak{m})$, we send

$$\mu\in\mathrm{S}_+(X)\otimes\mathfrak{m}\mapsto t\Big(1-e^{\frac{\mu}{t}}\Big)\in\mathrm{S}(X)\otimes\mathfrak{m}.$$

This truncates at finite order, so the map is well-defined.

We compute the tangent map at $\mu=0$, and we see

$$\delta\mu\mapsto t\left(1-e^{\frac{\delta\mu}{t}}\right) = -\delta\mu+\cdots,$$

so we see that

$$\mathrm{d}\mathscr{P}_*\colon T_0\mathrm{S}_+(X) = \mathrm{S}_+(X)\to T_0\mathrm{S}(X) = \mathrm{S}(X)$$

is simply the inclusion with a sign of -1. See Figure 5 for a visualization. This implies that $\mathscr{P}$ is a formal embedding.

This map is actually an L_∞-morphism

$$(\mathrm{S}_+(X), Q, \{-,-\})\xrightarrow{\mathscr{P}}(\mathrm{S}(X), Q, 0).$$




FIGURE 5. The cochain period map

One way to see this is to check that the Maurer-Cartan equation is preserved. We can check that

$$Q\left(t\left(1-e^{\frac{\mu}{t}}\right)\right) = -\left(Q\mu + \frac{1}{2}\{\mu,\mu\}\right)e^{\frac{\mu}{t}},$$

and therefore we see that μ satisfies the Maurer-Cartan equation if and only if $\mathscr{P}(\mu)$ does. Therefore, if we view Q as a vector field on $S(X)$, then Q is tangent to $\mathrm{Im}(\mathscr{P})$ and is thus the pushforward of a vector field on $S_+(X)$ which defines the L_∞ structure.

The image $\mathrm{Im}(\mathscr{P}) \subset S(X)$ of the period map is a formal Lagrangian submanifold. If we calculate the variation of the period map, we see that

$$\begin{aligned}\delta\left(t\left(1-e^{\frac{\mu}{t}}\right)\right) &= -\delta\mu e^{\frac{\mu}{t}} \\ &\subset S_+(X)e^{\frac{\mu}{t}}.\end{aligned}$$

Proposition 4.12. *For all $\alpha, \beta \in S_+(X)$, we have*

$$\omega\left(\alpha e^{\frac{\mu}{t}}, \beta e^{\frac{\mu}{t}}\right) = 0.$$

Proof. The proof is left as an exercise. □

5. Classical BCOV action

Some references for this include the surveys Some classical/quantum aspects of Calabi-Yau moduli and Renormalization method and mirror symmetry by the lecturer in addition to his work with Costello.

We will begin with a toy model to describe the geometry of gravitational descendants. Suppose the case when X is a point. In this case, $S = \mathbb{C}(\!(t)\!)$, $S_+ = \mathbb{C}[\![t]\!]$, and $S_- = t^{-1}\mathbb{C}[t^{-1}]$. The symplectic pairing is given by

$$\omega(f(t), g(t)) = \mathrm{Res}_{t=0} f(t)g(-t)\,\mathrm{d}t.$$

The period map now looks like Figure 6. The image of the period map is a formal Lagrangian submanifold.




FIGURE 6. The cochain period map for $X = \mathrm{pt}$

Under the identification $S = T^*S_+$, we write the image of the period map as the graph of dI_0 for some $I_0 \in \mathscr{O}(S_+)$. We will parameterize $\tau \in S_+$ as

$$\tau = \sum_{k=0}^{\infty} \tau_k t^k.$$

Then we can write $I_0 = I_0(\tau_k)$, and our goal is to compute this. We see that

$$\mathrm{Im}\mathscr{P} = \sum_{k=0}^{\infty} \tau_k t^k + \sum_{k=0}^{\infty} \partial_{\tau_k} I_0 (-t)^{-(k+1)}.$$

Denoting this by $\varphi(\tau_k)$, we will try to find its geometry. This actually becomes a combinatorial problem, but this actually becomes too complicated. Instead, it can be charcterized by various properties, which correspond to properties of the Lagrangian cone in Gromov-Witten theory.

The first property is the *string equation*. Let $\hat{t}\colon S \to S$ be the operator which sends $t^k \mapsto t^{k+1}$. Now define

$$L_{-1} = \hat{t}^{-1},$$

which is essentially multiplying by t^{-1}. We can view L_{-1} as a vector field on S, which is actually symplectic, namely satisfying the equation

$$\omega(L_{-1}f, g) = -\omega(f, L_{-1}g).$$

Writing the symplectic form as

$$\begin{aligned}\omega &= \sum_{k\geq 0} d\tau_k \wedge d\tau_{-k-1}(-1)^{k+1} \\ &= \frac{1}{2} \sum_{i+j=-1} (-1)^j \, d\tau_i \wedge d\tau_j,\end{aligned}$$

we see that

$$L_{-1} = \sum_i \tau_{i+1} \partial_{\tau_i}.$$

This implies that

$$\begin{aligned}\iota_{L_{-1}}\omega &= \sum_{i+j=-1} (-1)^j \tau_{i+1} \, d\tau_j \\ &= \frac{1}{2} d\left(\sum_{i+j=0} (-1)^j \tau_i \tau_j \right).\end{aligned}$$

Therefore, we will write the Hamiltonian as

$$h_{-1} = \frac{1}{2}\tau_0^2 + \sum_{k\geq 1}(-1)^k \tau_k \tau_{-k}.$$

The following result is one of the basic results in the Givental formalism.

Lemma 5.1. *The vector field L_{-1} is tangent to the object $\mathrm{Im}\mathscr{P} - t$, which is known as the dilaton shift.*

Proof. Let $\alpha = \mathscr{P}(f) - t = -te^{\frac{f}{t}}$. We calculate the tangent space

$$\begin{aligned} T_\alpha(\mathrm{Im}\mathscr{P} - t) &= \left\{-\delta f e^{\frac{f}{t}}\right\} \\ &= \mathrm{S}_+ e^{\frac{f}{t}}. \end{aligned}$$

We conclude by calculating $L_{-1}|_\alpha = L_{-1}(\alpha) = -e^{\frac{f}{t}} \subset T_\alpha(\mathrm{Im}\mathscr{P} - t)$. □

We can now write down a Hamilton-Jacobi equation, which says that $h_{-1}|_{\mathrm{Im}\mathscr{P}-t}$ is constant. In this example, the constant will be zero, we see that

$$\frac{1}{2}\tau_0^2 + \sum_{k=0}^{\infty} \tau_{k+1}\frac{\partial}{\partial \tau_k} I_0 - \frac{\partial}{\partial \tau_0} I_0 = 0.$$

This is equivalent to the PDE

$$\frac{\partial}{\partial \tau_0} I_0 = \frac{1}{2}\tau_0^2 + \sum_{k=0}^{\infty} \tau_{k+1}\frac{\partial}{\partial \tau_k} I_0,$$

which is exactly the string equation from Gromov-Witten theory.

The second property we will consider is the *dilaton equation*. Consider the operator

$$L_0 \colon \mathrm{S} \to \mathrm{S} \qquad t^k \mapsto \left(k + \frac{1}{2}\right) t^k.$$

This is also a symplectic operator satisfying

$$\omega(L_0 f, g) + \omega(f, L_0 g) = 0.$$

The Hamiltonian function is now given by

$$h_0 = \sum_{k=0}^{\infty}\left(k + \frac{1}{2}\right)(-1)^{-k-1}\tau_k \tau_{-k-1}.$$

Note that all of our operators are linear, so the Hamiltonians are quadratic.

Lemma 5.2. *The vector field L_0 is tangent to $\mathrm{Im}\mathscr{P} - t$.*

Proof. The proof is similar to the string equation, so it is left as an exercise. □

We can once again write the Hamilton-Jacobi equation, which says that

$$\sum_{k=0}^{\infty}\left(k + \frac{1}{2}\right)(-1)^{-k-1}\tau_k \tau_{-k-1}\Big|_{\mathrm{Im}\mathscr{P}-t} = 0.$$

This implies that

$$\frac{3}{2}\frac{\partial}{\partial \tau_1} I_0 = \sum_{k=0}^{\infty}\left(k + \frac{1}{2}\right)\tau_k \frac{\partial}{\partial \tau_k} I_0,$$

which is exactly the dilaton equation from Gromov-Witten theory. We may continue this and write down the classical Virasoro constraints, but we will stop here.

There is a third equation, called the *grading equation*. Consider the *degree operator*

$$G\colon \mathrm{S} \to \mathrm{S} \qquad t^k \mapsto (2k-2)t^k.$$

Here, we are remembering that there is actually a shift by 2 that we have been suppressing and that t has degree 2.

Lemma 5.3. *G is tangent to the Lagrangian* $\operatorname{Im}\mathscr{P}$ *(with no dilaton shift).*

Unfortunately, this is not a Hamiltonian vector field (in particular the symplectic form has a nonzero degree). The lemma is equivalent to the statement that $\operatorname{Im}\mathscr{P}$ is preserved by the $\mathbb{C}^\times$-action taking $t^k \mapsto \lambda^{2k-2}t^k$. In particular, we see that $I_0(\tau_k)$ is homogeneous of degree 6 if we assign $\deg \tau_k = 2-2k$. Combining this with the string and dilaton equations, it is easy to show the following.

Proposition 5.4. *For each $k_1, \dots, k_n$, we have*

$$\frac{\partial}{\partial\tau_{k_1}}\cdots\frac{\partial}{\partial\tau_{k_n}}I_0\bigg|_{\tau=0} = \int_{\bar{\mathscr{M}}_{0,n}} \psi_1^{k_1}\cdots\psi_n^{k_n} = \binom{n-3}{k_1,\dots,k_n}.$$

The general case follows easily from the point case, and in particular we can determine the *classical BCOV action*. Denote the image of the cochain period map (see Figure 5) by $\mathscr{L}_X$, which is a formal Lagrangian submanifold of $\mathrm{S}(X) = \mathrm{S}_+(X)\oplus \mathrm{S}_-(X) = T^*\mathrm{S}_+(X)$. Writing $\mathscr{L}_X$ as the graph of $\mathrm{d}I_0^X$, we can determine I_0^X as a function on $\mathrm{S}_+(X) = \mathrm{PV}(X)[\![t]\!][2]$. Denote

$$\left\langle t^{k_0}\otimes\cdots\otimes t^{k_n}\right\rangle_0 \coloneqq \int_{\bar{\mathscr{M}}_{0,n}} \psi_1^{k_1}\cdots\psi_n^{k_n} = \binom{n-3}{k_1,\dots,k_n}.$$

This extends $\mathrm{PV}(X)$-linearly to a map

$$\langle -\rangle_0\colon \operatorname{Sym}^\bullet(\mathrm{S}_+(X)) \to \mathrm{PV}(X).$$

Theorem 5.5 (Costello-Li)**.** *We have the formula*

$$\begin{aligned} I_0^X(\mu) &= \operatorname{Tr}\langle e^\mu\rangle_0 \\ &= \sum_{n\geq 3}\frac{1}{n!}\operatorname{Tr}\langle\mu^{\otimes n}\rangle_0. \end{aligned}$$

This is known as the classical BCOV action.

If we write $\mu = \mu_0 + \mu_1 t + \mu_2 t^2 + \cdots$ for $\mu_k \in \mathrm{PV}(X)$, we see that

$$I_0^X = \frac{1}{3!}\operatorname{Tr}\mu_0^3 + \cdots$$

In the Kodaira-Spencer gravity of BCOV, the fields are $\mu_0 \in \ker\partial \subset \mathrm{PV}(X)$, and their action is given by

$$\mathrm{KS}[\mu_0] = \frac{1}{2}\operatorname{Tr}\mu_0\frac{\bar\partial}{\partial}\mu_0 + \frac{1}{3!}\operatorname{Tr}\mu_0^3.$$

This looks a bit unusual, because we can find the Maurer-Cartan equation, but the original theory is nonlocal, which poses problems for quantization (which is why we replace $\ker\partial$ with the complex $[\mathrm{PV}\xrightarrow{t\partial} t\mathrm{PV}\xrightarrow{t\partial} t^2\mathrm{PV}\to\cdots]$).

6. Hodge structure and primitive forms

We can pass our picture from the cochan level to cohomology (or really the Maurer-Cartan functor). We will let

$$\mathscr{M}_X = \left\{\mu \in \mathrm{S}_+(X) \mid Q\mu + \frac{1}{2}\{\mu,\mu\} = 0\right\}/\text{gauge equivalence.}$$

The extended period map

$$\mathscr{P}\colon \mathscr{M}_X \hookrightarrow H^\bullet(\mathrm{S}(X), Q)$$

Because Q is compatible with ω, there is an induced symplectic form (shifted by $6-2d$) on $H^\bullet(\mathrm{S}(X),Q)$, which we also call ω. If we consider the subcomplex

$$(\mathrm{S}_+(X), Q) \subset (\mathrm{S}(X), Q),$$

then we see an isotropic subspace

$$H^\bullet(\mathrm{S}_+(X), Q) \subset H^\bullet(\mathrm{S}(X), Q).$$

However, S_- is not preserved by Q, so we do not have a preferred choice of splitting at this level.

Definition 6.1. An *opposite filtration* of $H^\bullet(\mathrm{S}(X),Q)$ is a choice of linear isotropic subspace $\mathscr{L} \subset H^\bullet(\mathrm{S}(X),Q)$ such that

(1) There is an isotropic splitting $H^\bullet(\mathrm{S}(X),Q) = H^\bullet(\mathrm{S}_+(X),Q) \oplus \mathscr{L}$;
(2) $\mathscr{L}$ is preserved by t^{-1}.

A choice of opposite filtration splits the map

$$\begin{array}{ccc} H^\bullet(\mathrm{S}_+(X),Q) & \longrightarrow & H^\bullet(\mathrm{S}_+(X),Q)/tH^\bullet(\mathrm{S}_+(X),Q) \\ & \overset{\mathscr{L}}{\nwarrow} & \| \\ & & H^\bullet(\mathrm{PV}(X),\bar\partial) \end{array}$$

by the natural isomorphism

$$\begin{aligned} B^{\mathscr{L}} &:= H^\bullet(\mathrm{S}_+(X),Q) \cap t\mathscr{L}H^\bullet(\mathrm{PV}(X),\bar\partial) \\ &\simeq H^\bullet(\mathrm{S}_+(X),Q)/tH^\bullet(\mathrm{S}_+(X),Q) \\ &\simeq H^\bullet(\mathrm{PV},\bar\partial). \end{aligned}$$

If we write $\mathscr{H} := H^\bullet(\mathrm{S}(X),Q)$ and $\mathscr{H}_+ = H^\bullet(\mathrm{S}_+(X),Q)$, then $\mathscr{H} = B^{\mathscr{L}}(\!(t)\!)$ and $\mathscr{H}_+ = B^{\mathscr{L}}[\![t]\!]$, where we choose $\mathscr{L} = t^{-1}B^{\mathscr{L}}[t^{-1}]$. This allows us to write

$$\mathscr{H} = T^*\mathscr{H}_+.$$

Using this, the period map looks like Figure 7. We then see that $\mathscr{P}(\mathscr{M}_X)$ is the graph of $\mathrm{d}\mathscr{F}_0^{X,\mathscr{L}}$, where $\mathscr{F}_0^{X,\mathscr{L}}$ is a function on $\mathscr{H}_+ = H^\bullet(\mathrm{PV},\bar\partial)[\![t]\!]$, which is called the *B-model state space* (and $\mathscr{F}_0^{X,\mathscr{L}}$ is the *genus-zero B-model potential*).

We now consider the geometry of the opposite filtration. Define the map

$$\Gamma_\Omega\colon \mathrm{PV}(X)(\!(t)\!) \to \Omega(X)[\![t]\!] \qquad t^k\mu \mapsto t^{k+i-1}\mu \lrcorner \Omega$$

for $\mu \in \mathrm{PV}^{i,j}$. This intertwines the operator Q with the standard de Rham differential $\mathrm{d} = \bar\partial + \partial$. Observe that

$$\Gamma_\Omega(\mathrm{S}_+(X)) = \prod_{p\in\mathbb{Z}} t^{d-p-1}\mathscr{F}^p\Omega(X),$$




FIGURE 7. The period map at the cohomology level

where $\mathscr{F}^p\Omega(X) = \Omega^{\geq p,*}(X)$ is the Hodge filtration. Here, we can see that

$$\begin{aligned}\Gamma_\Omega(t^k \mathrm{PV}^{i,*}) &= t^{k+i-1}\Omega^{d-i,*} \\ &= t^{d-p-1}\Omega^{p+k,*},\end{aligned}$$

where we identify $p = d - i - k$. In particular, we see that

$$\Gamma_\Omega \colon \mathrm{S}_+(X)/t\mathrm{S}_+(X) \simeq \prod_p t^{d-p-1}\,\mathrm{Gr}^p\,\Omega(X)$$

gives us a sum of graded pieces of the Hodge filtration. In particular, we see that $\mathscr{L}$ is equivalent to a splitting of the Hodge filtration. Some examples include the following:

(1) If we let $\mathscr{L} = \overline{\mathscr{F}}^p$ is the complex conjugate splitting (the harmonic splitting), we obtain the usual Hodge decomposition

$$H^n(X) = \bigoplus_{p+q=n} H^{p,q}(X).$$

However, this splitting does not give the correct B-model invariants.

(2) We can let $\mathscr{L}$ be the monodromy splitting, which arises near the large complex structure limit. This is in fact the correct splitting in the B-model and gives the mirror to Gromov-Witten theory. If we move to a different point (maybe the Gepner point), we get a different monodromy splitting, which is mirror to FJRW theory.

Remark 6.2. For a Calabi-Yau threefold, we have a symplectic subspace

$$t\mathrm{PV}^{0,0} \oplus \mathrm{PV}^{1,1} \oplus t^{-1}\mathrm{PV}^{2,2} \oplus t^{-2}\mathrm{PV}^{3,3}$$

and Γ_Ω identifies this with the space

$$\Omega^{3,0} \oplus \Omega^{2,1} \oplus \Omega^{1,2} \oplus \Omega^{0,3}.$$

We will now introduce primitive forms (due to Kyoji Saito), or equivalently, J-functions (due to Givental). Fix a splitting $\mathscr{L}$ and a subspace

$$B^{\mathscr{L}} := H^\bullet(\mathrm{S}_+(X), Q) \cap t\mathscr{L} \simeq H^\bullet(\mathrm{PV}, \bar{\partial})$$

with basis $\{\varphi_\alpha\}$.

Definition 6.3 (Higher residue pairing)**.** Define the pairing

$$K\colon \mathscr{H}\otimes\mathscr{H}\to\mathbb{C}((t)) \qquad K(\alpha f(t),\beta g(t)) \coloneqq \mathrm{Tr}(\alpha\beta)f(t)g(-t).$$

When we come to the Landau-Ginzburg B-model, this is literally Saito's higher residue pairing. In addition, note that

$$\omega(\mu,\nu)=\oint K(\mu,\nu)\,\mathrm{d}t.$$

If we reduce modulo t, this simply becomes the trace pairing on $H^\bullet(X,\Lambda^*TX)=H^\bullet(\mathsf{PV},\bar\partial)$.

There are some peculiar properties here. First, the isotropic splitting

$$\mathscr{H}=\mathscr{H}_+\oplus\mathscr{L}$$

implies that

$$K(\mathscr{H}_+,\mathscr{H}_+)\subset\mathbb{C}[\![t]\!], \qquad K(\mathscr{L},\mathscr{L})\subset t^{-2}\mathbb{C}[t^{-1}].$$

Together, these tell us that

$$K\colon B^{\mathscr{L}}\otimes B^{\mathscr{L}}\to\mathbb{C}$$

sends $B^{\mathscr{L}}$ to constants. In our basis, we will write

$$K(\varphi_\alpha,\varphi_\beta)=\eta_{\alpha\beta}.$$

Saito calls this choice a *good basis*. With our choice of splitting, write

$$\pi_+^{\mathscr{L}}\colon\mathscr{H}\to\mathscr{H}_+$$

for the corresponding projection. Under the choice of good basis $B^{\mathscr{L}}$, we see that

$$\pi_+^{\mathscr{L}}\colon\mathscr{H}=B^{\mathscr{L}}((t))\to\mathscr{H}_+=B^{\mathscr{L}}[\![t]\!]$$

simply eliminates the terms of negative degree in t.

Now consider the fiber product

$$\begin{array}{ccc}\mathscr{M}_X^0 & \hookrightarrow & \mathscr{M}_X\\ \simeq\downarrow & & \simeq\downarrow{\scriptstyle \pi_+^{\mathscr{L}}\circ\mathscr{P}}\\ B^{\mathscr{L}} & \hookrightarrow & \mathscr{H}_+.\end{array}$$

Let's write coordinates $\left\{\tau_k^\alpha\right\}_{k\geq 0}$ for $\mathscr{H}$ and coordinates $\left\{\tau_0^\alpha,\tau_1^\alpha=0,\ldots\right\}$ for $B^{\mathscr{L}}$. This identifies a family

$$\mu^{\mathscr{L}}(\tau_0^\alpha)\in\mathsf{PV}(X)[\![t]\!]$$

solving the Maurer-Cartan equation

$$Q\mu^{\mathscr{L}}(\tau_0^\alpha)+\frac{1}{2}\left\{\mu^{\mathscr{L}}(\tau_0^\alpha),\mu^{\mathscr{L}}(\tau_0^\alpha)\right\}=0.$$

This will play the role of the *primitive form* in Saito's theory. The solution also satisfies

$$\pi_+^{\mathscr{L}}\left[t\left(1-e^{\frac{\mu^{\mathscr{L}}}{t}}\right)\right]_Q=\sum_\alpha\tau_0^\alpha\varphi_\alpha.$$

Here, the notation $[-]_Q$ means taking the cohomology class with respect to Q. This can be rewritten as the equation

$$\left[t\left(1-e^{\frac{\mu^{\mathscr{L}}}{t}}\right)\right]_Q = \sum_\alpha \tau_0^\alpha \varphi_\alpha + \sum_{k=0}^\infty (-1)^{-k-1} t^{-k-1} \eta^{\alpha\beta} \frac{\partial \mathscr{F}_0^{\mathscr{L}}}{\partial \tau_k^\beta} \varphi_\beta \Bigg|_{\tau_{>0}=0}$$

using the fact that the period map has image the graph of $\mathrm{d}\mathscr{F}_0^{\mathscr{L}}$. We can see that the right-hand side of this equation is the J-function $J(\tau_0^\alpha; t)$, which is the analogue of the J-function defined by Givental in Gromov-Witten theory. We can again rewrite this as

$$\left[e^{\frac{\mu^{\mathscr{L}}}{t}}\right]_Q = 1 - t^{-1} J,$$

and the left-hand side here is the *primitive form* defined by Saito.

Proposition 6.4. *The quantity*

$$K(\partial_{\tau_0^\alpha} J, \partial_{\tau_0^\beta} J) = \eta_{\alpha\beta}$$

is independent of t.

Proof. By the expression of J, we see that

$$\begin{aligned} K(\partial_{\tau_0^\alpha} J, \partial_{\tau_0^\beta} J) &= K(\varphi_\alpha + O(t^{-1}), \varphi_\beta + O(t^{-1})) \\ &= \eta_{\alpha\beta} + O(t^{-1}). \end{aligned}$$

On the other hand, we can calculate

$$\begin{aligned} K(\partial_{\tau_0^\alpha} J, \partial_{\tau_0^\beta} J) &= K\left(\partial_{\tau_0^\alpha}\left[t - te^{\frac{\mu^{\mathscr{L}}}{t}}\right]_Q, \partial_{\tau_0^\beta}\left[t - te^{\frac{\mu^{\mathscr{L}}}{t}}\right]_Q\right) \\ &= K\left((\partial_{\tau_0^\alpha}\mu^{\mathscr{L}})e^{\frac{\mu^{\mathscr{L}}}{t}}, (\partial_{\tau_0^\beta}\mu^{\mathscr{L}})e^{\frac{\mu^{\mathscr{L}}}{t}}\right) \in \mathbb{C}[\![t]\!], \end{aligned}$$

where we use the fact that this last expression lies in the tangent space to the Lagrangian cone. The conclusion follows by comparing our two calculations. □

We see now that $\left\{\partial_{\tau_0^\alpha} J\right\}$ defines a family of *good bases* parameterized by τ_0^α. Geometrically, we see that

$$\mathrm{Span}\left\{\partial_{\tau_0^\alpha} J\right\}_{\tau_0^\alpha} = T_{\tau_0^\alpha} \mathscr{M}_X \cap t\mathscr{L}.$$

At $\tau_0^\alpha = 0$, we see that $T_0\mathscr{M} \cap t\mathscr{L} = \mathscr{H}_+ \cap t\mathscr{L} = B^{\mathscr{L}}$ is our original choice of good basis. Because we have the splitting

$$\mathscr{H} = T_0\mathscr{M} \oplus \mathscr{L},$$

we see that the splitting

$$\mathscr{H} = T_{\tau_0^\alpha}\mathscr{M} \oplus \mathscr{L}$$

holds for nearby τ_0^α. The idea (due to Givental) is that the set $\left\{\partial_{\tau_0^\alpha} J\right\}$ forms a basis of $T_{\tau_0^\alpha}\mathscr{M} \cap t\mathscr{L}$.

We can now compute the second derivatives $\partial_{\tau_0^\alpha}\partial_{\tau_0^\beta}J$ in two ways as we did with the higher residue pairing. On one hand, the expression of J tells us that

$$\partial_{\tau_0^\alpha}\partial_{\tau_0^\beta}J \in \mathscr{L}.$$

On the other hand, we calculate

$$\begin{aligned}\partial_{\tau_0^\alpha}\partial_{\tau_0^\beta}J &= \partial_{\tau_0^\alpha}\partial_{\tau_0^\beta}\left[t - te^{\frac{\mu^{\mathscr{L}}}{t}}\right]_Q \\ &= \left[-\left(\frac{1}{t}\partial_{\tau_0^\alpha}\mu^{\mathscr{L}}\partial_{\tau_0^\beta}\mu^{\mathscr{L}} + \partial_{\tau_0^\alpha}\partial_{\tau_0^\beta}\mu^{\mathscr{L}}\right)e^{\frac{\mu^{\mathscr{L}}}{t}}\right]_Q \in \frac{1}{t}T_{\tau_0^\alpha}\mathscr{M}.\end{aligned}$$

In particular, the two calculations imply that

$$\partial_{\tau_0^\alpha}\partial_{\tau_0^\beta}J \in \frac{1}{t}T_{\tau_0^\alpha}\mathscr{M} \cap \mathscr{L} = \frac{1}{t}\left(T_{\tau_0^\alpha}\mathscr{M} \cap t\mathscr{L}\right).$$

This now implies that there exists $A^\gamma_{\alpha\beta}(\tau_0^\bullet)$ such that

$$\partial_{\tau_0^\alpha}\partial_{\tau_0^\beta}J + \frac{1}{t}A^\gamma_{\alpha\beta}\partial_{\tau_0^\gamma}J = 0.$$

Passing to the primitive form, this is now equivalent to the statement that

$$\left(\partial_{\tau_0^\alpha}\partial_{\tau_0^\beta} + \frac{1}{t}A^\gamma_{\alpha\beta}\partial_{\tau_0^\gamma}\right)\left[e^{\frac{\mu^{\mathscr{L}}}{t}}\right] = 0.$$

Remark 6.5. There is a perturbative algorithm for the Landau-Ginzburg B-model due to Li-Li-Saito, which allows us to compute the primitive form order by order using a method similar to Birkhoff factorization.

7. Classical master equation

Recall the period map as in Figure 5. Using the string and dilaton equations, we determined that $\mathrm{Im}\mathscr{P}$ was the graph of $\mathrm{d}I_0^X$, where

$$I_0^X(\mu) = \mathrm{Tr}\langle e^\mu\rangle_0.$$

Our goal now will be to study some structure relating to Q, which will allow us to pass to a finite-dimensional model. Q is odd (whereas the Virasoro constraints are even), so we will write down an odd version of the Hamilton-Jacobi equation, which will be the *classical master equation*.

Because Q preserves ω, we will write down the quadratic Hamiltonian h_Q, which will satisfy $h_Q|_{\mathrm{Im}\mathscr{P}} = 0$. The differential $Q = \bar\partial + t\partial$ satisfies $Q^2 = 0$, and is therefore a *cohomological vector field*. Here are some properties of the operator Q:

- It preserves $\mathrm{S}_+(X)$;
- It does not preserve $\mathrm{S}_-(X)$.

To give an idea of the general picture and how to quantize the theory, we will consider a finite-dimensional toy model. Let

$$(V, \omega, Q)$$

be a dg (even) symplectic vector space (and for simplicity, we will assume the degree is 0), so in particular we can consider one of

$$\omega \in \Lambda^2 V^* \qquad \text{or} \qquad \omega^{-1} \in \Lambda^2 V.$$

By definition, the differential Q is skew-symmetric with respect to ω. Assume that we have a splitting

$$V = V_+ \oplus V_-,$$

where both V_+ and V_- are Lagrangian subspaces. We will require that Q preserves V_+ but not necessarily V_-. The splitting will allow us to make the identification $V = T^*V_+$ and consider the projection

$$\pi_+ \colon V \to V_+.$$

We know that $\omega^{-1} \in \Lambda^2 V \subset (V_+ \otimes V_-) \oplus (V_- \otimes V_+)$. To write down the Hamiltonian function, consider

$$(Q \otimes 1)\omega^{-1} \in (V_+ \otimes V_-) \oplus (V_- \otimes V_+) \oplus (V_+ \otimes V_+).$$

Denote the component in $V_+ \otimes V_+$ by

$$P = (\pi_+ \otimes \pi_+)[(Q \otimes 1)\omega^{-1}] \in V_+ \otimes V_+,$$

which plays the role of the BV kernel and measures the failure of Q to preserve V_-.

Proposition 7.1. *P defines a degree 1 element of $\mathrm{Sym}^2(V_+)$ and is Q-compatible, namely that*

$$(Q \otimes 1 + 1 \otimes Q)P = 0.$$

Proof. Using the fact that Q is skew-symmetric with respect to ω, we see that

$$(Q \otimes 1 + 1 \otimes Q)\omega^{-1} = 0.$$

This implies that

$$(Q \otimes 1)\omega^{-1} \in \mathrm{Sym}^2(V)$$

is symmetric, and projecting to $V_+ \otimes V_+$ concludes the proof. □

This is actually a dual description of the Hamiltonian function h_Q. Geometrically, we can consider

$$(Q \otimes 1)\omega^{-1} \in \mathrm{Sym}^2(V) \xrightarrow{\omega} \mathrm{Sym}^2(V^*) \ni h_Q,$$

which exactly gives the quadratic Hamiltonian associated to Q. Using the fact that

$$(Q \otimes 1)\omega^{-1} \in (V_+ \otimes V_-) \oplus (V_- \otimes V_+) \oplus \mathrm{Sym}^2(V_+),$$

we see that

$$h_Q \in (V_-^* \otimes V_+^*) \oplus (V_+^* \otimes V_-^*) \oplus \mathrm{Sym}^2(V_-^*).$$

We see that the Hamilton-Jacobi equation will have a linear term coming from the first two components and a quadratic term coming from the last component.

Definition 7.2. Denote the differential operator

$$\partial_P \colon \mathrm{Sym}^n(V_+^*) \to \mathrm{Sym}^{n-2}(V_+^*)$$

by contracting with $P \in \mathrm{Sym}^2(V_+)$. More explicitly, we define

$$(\partial_P \varphi)(a_1 \otimes \cdots \otimes a_{n-2}) \coloneqq (-1)^{|\varphi|}\varphi(P \otimes a_1 \otimes \cdots \otimes a_{n-2}).$$

Also define the *BV bracket*

$$\{f, g\}_P \coloneqq \partial_P(fg) - (\partial_P f)g - (-1)^{|f|} f \partial_P g.$$

Proposition 7.3. *The data $(\mathscr{O}(V_+) = \mathrm{Sym}^\bullet(V_+^*), Q, \partial_P, \{-,-\}_P)$ defines a dg Gerstenhaber-Batalin-Vilkovisky (dGBV) algebra.*

Proof. Because $(Q\otimes 1+1\otimes Q)P=0$, we see that $[Q,\partial_P]=0$. The rest of the properties are straightforward to check. □

Now assume that we have a Lagrangian submanifold

$$\mathscr{L}\subset V=T^*V_+$$

which is the graph of $\mathrm{d}I_0$ for some function $I_0\in\mathscr{O}(V_+)$.

Proposition 7.4. *Q is tangent to $\mathscr{L}$ if and only if*

$$QI_0+\frac{1}{2}\{I_0,I_0\}_P=0.$$

This is called the classical master equation.

Proof. The odd Hamilton-Jacobi equation says that $h_Q|_{\mathscr{L}}=0$. The mixed part of h_Q corresponds to Q and the purely momentum part corresponds to the BV bracket. □

Now denote by

$$\delta=Q+\{I_0,-\}_P$$

the corresponding cohomological vector field. The classical master equation can now be rewritten as $\delta^2=0$. In fact, with the picture as in Figure 8, we see that

$$\delta=(\pi_+)_*[Q|_{\mathscr{L}}].$$

Because $Q^2=0$, we see that $\delta^2=0$ as well.




FIGURE 8. Q tangent to $\mathscr{L}$ and the map π_+.

We will now return to our original infinite-dimensional setting of the B-model on a Calabi-Yau manifold X. The structure is a bit more complicated but is still similar. Recall that

$$Q\colon \mathrm{S}_-(X)\to \mathrm{S}_+(X)\oplus \mathrm{S}_-(X).$$

Also recall that

$$\omega(-,-)=\mathrm{Tr}(-,-)\,\mathrm{Res}_{t=0}(-\cdot -).$$

The problem comes from the trace pairing, which causes the Poisson tensor ω^{-1} to involve distributions as

$$\omega^{-1}=\sum_{k\in\mathbb{Z}}\delta_\Delta t^k\otimes(-t)^{-k-1},$$

where δ_Δ is a distributional element of $\mathrm{PV}(X)\otimes\mathrm{PV}(X)$ and is the integral kernel for the identity operator

$$1\colon \mathrm{PV}(X)\to\mathrm{PV}(X).$$

Note that this is fine classically but will presernt problems for quantizing the theory. We can still write down the BV kernel

$$\begin{aligned} P &= (\pi_+ \otimes \pi_+)[(Q \otimes 1)\omega^{-1}] \\ &= (\partial \otimes 1)\delta_\Delta \end{aligned}$$

because Q only fails to preserve $S_-(X)$ on $t^{-1}\mathrm{PV}$ via the $t\partial$ term. Given a local functional I on $S_+(X) = \mathrm{PV}(X)[\![t]\!]$ of the form $I = I(\mu_k) = \int \cdots$, its variation can be computed as

$$\delta I = \sum_k \mathrm{Tr}\left(\delta\mu_k \frac{\delta I}{\delta \mu_k}\right).$$

We define the BV bracket for two local functionals I_1 and I_2 by

$$\{I_1, I_2\}_0 = \mathrm{Tr}\left(\frac{\delta I_1}{\delta \mu_0}\right)\partial\left(\frac{\delta I_2}{\delta \mu_0}\right),$$

which produces another local functional. Copying the finite-dimensional case, we have the following theorem.

Theorem 7.5 (Costello-Li)**.** *The BCOV action $I_0^X = \mathrm{Tr}\langle e^\mu \rangle_0$ satisfies the classical master equation*

$$QI_0^X + \frac{1}{2}\left\{I_0^X, I_0^X\right\}_0 = 0.$$

Remark 7.6. The theorem is proved directly in the lecturer's PhD thesis, where he finds that the classical master equation is equivalent to the genus-zero topological recursion relations.

8. Quantum master equation

Note that the map $\pi_+ \circ \mathscr{P} \colon S_+(X) \to S_+(X)$ is a diffeomorphism. However, we gain the classical interaction I_0^X and the Maurer-Cartan equation with respect to δ becomes

$$Q\mu + \sum_{k\geq 2} \frac{1}{k!}\ell_k(\mu^{\otimes k}) = 0,$$

where $\ell_2, \ell_3, \ldots$ emerge from the $\left\{I_0^X, -\right\}$ term.

We will now return to our toy model (V, Q, ω) of a dg symplectic vector space. We also continue to assume the Poisson kernel ω^{-1} and the isotropic splitting $V = V_+ \oplus V_-$. Q preserves V_+ but not V_-, and we wrote down the tensor

$$P = \pi_+ \otimes \pi_+[(Q \otimes 1)\omega^{-1}] \in \mathrm{Sym}^2(V_+).$$

This produced a dGBV algebra

$$(\mathscr{O}(V_+), Q, \partial_P, \{-,-\}_P)$$

Definition 8.1. We define the (formal) *Weyl algebra*

$$\mathscr{W}(V) \coloneqq \prod_{n\geq 0} (V^*)^{\otimes n}[\![\hbar]\!] \Big/ \sim,$$

where $\hbar$ is the formal (quantum) parameter and the relation $\sim$ is generated by

$$[a, b] \coloneqq a \otimes b - (-1)^{|a||b|} b \otimes a \sim \hbar\omega^{-1}(a \otimes b).$$

Note that the classical limit

$$\mathscr{W}(V)/\hbar \simeq \hat{\mathscr{O}}(V)$$

is the formal functions on V. We denote the annihilator $\mathrm{Ann}(V_+) \subset V^*$ to be

$$\mathrm{Ann}(V_+) = \left\{\varphi \in V^* \mid \varphi(V_+) = 0\right\}.$$

Because the differential Q preserves V_+, the induced differential $Q\colon V^* \to V^*$ preserves $\mathrm{Ann}(V_+)$ as well.

Proposition 8.2. *The annihilator* $\mathrm{Ann}(V_+) \subset V^*$ *is a subcomplex.*

Definition 8.3. The (formal) *Fock space* is defined by

$$\mathscr{F}\mathrm{ock}(V_+) \coloneqq \mathscr{W}(V)/\mathscr{W}(V)\,\mathrm{Ann}(V_+).$$

Remark 8.4. We treat $\mathrm{Ann}(V_+)$ as annihilation operators.

Because Q is compatible with ω, we see that $(\mathscr{W}(V), Q)$ defines a cochain complex with subcomplex $\mathscr{W}(V)\,\mathrm{Ann}(V_+) \subset \mathscr{W}(V)$. This induces a differential

$$Q\colon \mathscr{F}\mathrm{ock}(V_+) \to \mathscr{F}\mathrm{ock}(V_+).$$

The splitting $V = V_+ \oplus V_- \simeq T^*V_+$ induces a natural embedding $V_+^* \subset V^*$ given by extension by zero on V_-. Let

$$\hat{\mathscr{O}}(V_+) \coloneqq \prod_{n\geq 0} \mathrm{Sym}^n(V_+^*)$$

be the formal functions on V_+. This induces

$$\begin{array}{ccc} \hat{\mathscr{O}}(V_+)[\![\hbar]\!] & \hookrightarrow & \mathscr{W}(v) \\ & \overset{\simeq}{\searrow}^{\Phi} & \downarrow \\ & & \mathscr{F}\mathrm{ock}(V_+) \end{array}$$

and realizes $\hat{\mathscr{O}}(V_+)[\![\hbar]\!]$ as creation operators (which for physicists look like expressions of the form $\cdots a^\dagger|0\rangle$). Under this isomorphism, there is a differential $\hat{Q}$ on $\hat{\mathscr{O}}(V_+)[\![\hbar]\!]$, namely the unique one making the diagram

$$\begin{array}{ccc} \hat{\mathscr{O}}(V_+)[\![\hbar]\!] & \xrightarrow{\Phi} & \mathscr{F}\mathrm{ock}(V_+) \\ \downarrow{\scriptstyle \hat{Q}} & & \downarrow{\scriptstyle Q} \\ \hat{\mathscr{O}}(V_+)[\![\hbar]\!] & \xrightarrow{\Phi} & \mathscr{F}\mathrm{ock}(V_+) \end{array}$$

commute. However, this is not the same as the one induced from $Q\colon V_+ \to V_+$ (which extends to $Q\colon \hat{O}(V_+) \to \hat{\mathscr{O}}(V_+)$).

Proposition 8.5. *The difference between* $\hat{Q}$ *and* Q *is precisely the failure of* Q *to preserve the splitting, namely*

$$\hat{Q} = Q + \hbar\partial_P.$$

Proof. This is standard quantization of quadratic Hamiltonians and is left as an exercise. □

Definition 8.6. A formal function

$$F = \sum_{g \geq 0} F_g \hbar^g \in \hat{\mathscr{O}}(V_+)[\![\hbar]\!]$$

satisfies the *quantum BV master equation* (*QME*) if

$$(Q + \hbar\partial_P)e^{\frac{F}{\hbar}} = 0.$$

Equivalently, we can rewrite this as

$$(Q + \hbar\partial_P)F + \frac{1}{2}\{F, F\}_P = 0.$$

Note that in the classical limit $\hbar \to 0$, we recover the classical master equation

$$QF_0 + \frac{1}{2}\{F_0, F_0\}_P = 0.$$

We now discuss what is going on in a physical sense. Consider the state

$$|F\rangle \coloneqq \Phi\left(e^{\frac{F}{\hbar}}\right) = e^{\frac{F}{\hbar}}|0\rangle.$$

Then the quantum master equation is equivalent to the statement that $Q|F\rangle = 0$, which in physics is called the *quantum gauge consistency*. Passing to Q-cohomology (what physicists call *physical states*), we pass to

$$[|F\rangle] \in H^\bullet(\mathscr{F}\mathrm{ock}(V_+), Q),$$

which usually an element of a finite-dimensional space. If we let $\mathscr{H} = H^\bullet(V, Q)$ and $\mathscr{H}_+ = H^\bullet(V_+, Q)$, then we have a symplectic vector space $(\mathscr{H}, \omega)$ with $\mathscr{H}_+ \subset \mathscr{H}$ as an isotropic subspace. If we choose a polarization $\mathscr{H} = \mathscr{H}_+ \oplus \mathscr{L}$, the surjection $\mathscr{W}(V) \twoheadrightarrow \mathscr{F}\mathrm{ock}(V_+)$ reduces to a surjection

$$\mathscr{W}\mathrm{eyl}(\mathscr{H}) \twoheadrightarrow \mathscr{F}\mathrm{ock}(\mathscr{H}_+),$$

and the target is once again isomorphic to $\hat{\mathscr{O}}(\mathscr{H}_+)[\![\hbar]\!]$ because the polarization gives an embedding $\mathscr{H}_+^* \subset \mathscr{H}^*$. Because $Q|F\rangle = 0$, we obtain a state

$$[|F\rangle] \in \mathscr{F}\mathrm{ock}(\mathscr{H}_+) \cong H^\bullet(\mathscr{F}\mathrm{ock}(V_+), Q).$$

The choice of splitting gives us a formal function

$$F^{\mathscr{L}} = \sum_{g \geq 0} F_g^{\mathscr{L}} \hbar^g \in \hat{\mathscr{O}}(\mathscr{H}_+)[\![\hbar]\!].$$

Unfortunately, this is extremely hard to compute.

Warning 8.7. If we try to redo everything in the actual B-model, the "quantum master equation"

$$(Q + \hbar\partial_P)e^{\frac{F}{\hbar}} = 0$$

is ill-defined because P is singular. This will require choosing a Kähler metric, doing renormalization, and defining everything in a homotopical way.

9. Homotopic renormalization

10. Quantum BCOV theory

11. Landau-Ginzburg B-model

12. Open-closed B-model