Idea 1: Varying θ_{Frogs}

Frogs

November 30, 2024

1 Modelling and Data

1.1 Modelling

- $\theta_{Frogs}^{(0)} \sim \text{Po}(\lambda)$: initial number of frogs
- $\theta_{Frogs}^{(t)} = \theta_{Frogs}^{(t-1)} + R^{(t)}$: number of frogs at time t, $R^{(t)} \sim F_r$ is a random variable, can be Geometric, Possion, etc, with parameter r
- θ_{Love} , where $\theta_{Love} + 1 \sim \text{Geo}(p_{Love})$ (Can change dist): true love, (0 means don't want love, 1 means congwen, rest means frog number $\theta_{Love} 1$), assume first send letter = more attractive
- $\theta_p \sim U(0,1)$: Pr(reply|is true love)
- Each Frogs have probability p_{send} to send letter, and they will only send one letter
- Congwen will send a letter in sats and suns

1.2 Data x_t

- $t = 0, \dots, T$: time t
- $m_t \ge 0$: number of letters from congwen at time t
- $n_t \geq 0$: number of letters from other frogs at time t
- $Z_1, Z_2, \ldots, Z_{m_t} \in \{0, 1\}$: whether the congwen's letter is replied
- $X_1, X_2, \dots, X_{n_t} \in \{0, 1\}$: whether the other frog's letter is replied
- $\bullet \ Z_t = X_t = 0$
- Truely random variables are n_t , X_t , Z_t

2 Computations

$$\begin{split} f(Z_1 = \ldots = Z_{m_t} = 0 | m_t, \theta_{Frogs}^{(t)}, \theta_{Love}, \theta_p) &= (1 - \theta_p)^{m_t \mathbb{1}_{\{\theta_{Love} = 1\}}} \\ f(X_1 = \ldots = X_{n_t} = 0 | n_t, \theta_{Frogs}^{(t)}, \theta_{Love}, \theta_p) &= (1 - \theta_p)^{\mathbb{1}_{\{2 \le \theta_{Love} \le n_t + 1\}}} \\ f(n_t | \theta_{Frogs}^{(t)}, \theta_{Love}, \theta_p) &= \begin{pmatrix} \theta_{Frogs}^{(t)} \\ \theta_{Frogs}^{(t)} \end{pmatrix} p_{send}^{n_t} (1 - p_{send})^{\theta_{Frogs}^{(t)} - n_t} \mathbb{1}_{\{\theta_{Frogs}^{(t)} \ge n_t\}} \\ f(x_t | \theta_{Frogs}^{(t)}, \theta_{Love}, \theta_p) &= \frac{\theta_{Frogs}^{(t)} !}{n_t ! (\theta_{Frogs}^{(t)} - n_t)!} p_{send}^{n_t} (1 - p_{send})^{\theta_{Frogs}^{(t)} - n_t} \\ &\quad \cdot (1 - \theta_p)^{\mathbb{1}_{\{2 \le \theta_{Love} \le n_t + 1\}} + m_t \mathbb{1}_{\{\theta_{Love} = 1\}}} \mathbb{1}_{\{\theta_{Frogs}^{(t)} \ge n_t\}} \\ \theta_{Frogs}^{(t)} &= \theta_{Frogs}^{(0)} + \sum_{i=1}^{t} R^{(t)} \sim \text{Po}(\lambda + tr) \\ \pi(\theta_{Frogs}^{(0)}, \theta_{Love}, \theta_p) &= \frac{e^{-\lambda} \lambda_{Frogs}^{(0)}}{\theta_{Frogs}^{(0)}} p_{Love} (1 - p_{Love})^{\theta_{Love}} \mathbb{1}_{\{\theta_{Frogs}^{(t)}, \theta_{Love} \ge 0, 0 \le \theta_p \le 1\}} \\ \pi(\theta_{Frogs}^{(t)}, \theta_{Love}, \theta_p) &= \frac{e^{-(\lambda + tr)} (\lambda + tr)^{\theta_{Frogs}^{(t)}}}{\theta_{Frogs}^{(t)}} p_{Love} (1 - p_{Love})^{\theta_{Love}} \mathbb{1}_{\{\theta_{Frogs}^{(t)}, \theta_{Love} \ge 0, 0 \le \theta_p \le 1\}} \end{split}$$

$$\begin{split} & \chi([g_{Progs}^{(0)}, \theta_{Love}, \theta_{p}]x_{1}) \\ & \propto f(x_{1}|\theta) \pi(\theta) \\ & = \frac{\theta_{Progs}^{(1)}}{n_{1}!(\theta_{Progs}^{(0)}, -1)!} P_{send}^{send}(1 - p_{send})^{\theta_{Progs}^{(0)}} - n_{1} \\ & \cdot (1 - \theta_{p})^{1/2} \varepsilon^{s}_{trave} \varepsilon^{s}_{trave} \varepsilon^{s}_{trave})^{s}_{trave}^{s}_{trave} - 1 \\ & \cdot \frac{e^{-(\lambda+tr)}(\lambda + tr)^{\theta_{Progs}^{(0)}}}{\theta_{Progs}^{(1)}} p_{Love}(1 - p_{Love})^{\theta_{Love}} - 1 \\ & \cdot \frac{e^{-(\lambda+tr)}(\lambda + tr)^{\theta_{Progs}^{(0)}}}{\theta_{Progs}^{(1)}} p_{Love}^{s}(1 - p_{Love})^{\theta_{Love}^{s}}_{trave}^{s}_{$$

$$\begin{split} &\pi(\theta_{Frogs}^{(0)}, \theta_{Love}, \theta_{p} | x_{t}) \\ &= \sum_{\theta_{Frogs}^{(t)} = 0}^{\infty} \pi(\theta_{Frogs}^{(0)} | \theta_{Frogs}^{(t)}) \pi(\theta_{Frogs}^{(t)}, \theta_{Love}, \theta_{p} | x_{t}) \\ &= \frac{(\lambda/tr)^{\theta_{Frogs}^{(0)}}}{C(x_{t})\theta_{Frogs}^{(0)}!} (1 - p_{Love})^{\theta_{Love}} (1 - \theta_{p})^{\mathbb{I}_{\{2 \le \theta_{Love} \le n_{t}+1\}} + m_{t}\mathbb{I}_{\{\theta_{Love} = 1\}}} \mathbb{1}_{\{\theta_{Love} \ge 0, 0 \le \theta_{p} \le 1\}} \\ &\times \sum_{\theta_{Frogs}^{(t)} = \max\{n_{t}, \theta_{Frogs}^{(0)}\}} \frac{\theta_{Frogs}^{(t)}! ((1 - p_{send})tr)^{\theta_{Frogs}^{(t)}}}{(\theta_{Frogs}^{(t)} - \theta_{Frogs}^{(0)})! (\theta_{Frogs}^{(t)} - n_{t})!} \mathbb{1}_{\{\theta_{Frogs}^{(0)} \ge 0\}} \\ &\approx \frac{(\lambda/tr)^{\theta_{Frog}^{(0)}}}{C(x_{t})\theta_{Frogs}^{(0)}!} (1 - p_{Love})^{\theta_{Love}} (1 - \theta_{p})^{\mathbb{I}_{\{2 \le \theta_{Love} \le n_{t}+1\}} + m_{t}\mathbb{I}_{\{\theta_{Love} = 1\}}} \mathbb{1}_{\{\theta_{Frogs}^{(0)} \ge 0, \theta_{Love} \ge 0, 0 \le \theta_{p} \le 1\}} \\ &\times \sum_{i=1}^{n} \frac{(P_{i} + \max\{n_{t}, \theta_{Frogs}^{(0)}\})!}{(P_{i} + |n_{t} - \theta_{Frogs}^{(0)}|)!} ((1 - p_{send})tr)^{\max\{n_{t}, \theta_{Frogs}^{(0)}\}} e^{(1 - p_{send})tr} \qquad P_{1:n} \sim \text{Po}((1 - p_{seed})tr) \end{split}$$

Interesting:

$$(\theta_{Frogs}^{(t)} - n_t)|x_t \sim \text{Po}((1 - p_{send})(\lambda + tr))$$

$$\theta_{Frogs}^{(0)}|\theta_{Frogs}^{(t)} \sim \text{Bin}\left(\theta_{Frogs}^{(t)}, \frac{\lambda}{\lambda + tr}\right)$$