

# **ENGR 0021 Recitation 4 (Exam 1 Review)**

**February 6th, 2026**

Rie Huntington

Department of Industrial Engineering  
University of Pittsburgh

# **Descriptive Statistics**

# Measures of Central Tendency

---

**Key Concept:** Robustness

- **Mean:** Highly sensitive to outliers.
- **Median:** Robust (resistant) to outliers.
- **Mode:** Most frequent value.

**Example:** Consider the dataset  $\{1, 2, 3, 4, 100\}$ .

# Measures of Central Tendency

---

**Key Concept:** Robustness

- **Mean:** Highly sensitive to outliers.
- **Median:** Robust (resistant) to outliers.
- **Mode:** Most frequent value.

**Example:** Consider the dataset  $\{1, 2, 3, 4, 100\}$ .

- The mean is  $\frac{110}{5} = 22$ . The median is 3.
- If 100 changes to 1000, the mean jumps massively. The median stays 3.

# Variance and Standard Deviation

---

**Question:** You have a dataset of 50 numbers. The standard deviation, when calculated, is very small compared to the mean. What does this imply?

# Variance and Standard Deviation

---

**Question:** You have a dataset of 50 numbers. The standard deviation, when calculated, is very small compared to the mean. What does this imply?

→ It implies there is **very little variability**. Every single number in the dataset must be close to the mean for the standard deviation to be small.

# Variance and Standard Deviation

---

**Question:** You have a dataset of 50 numbers. The standard deviation, when calculated, is very small compared to the mean. What does this imply?

→ It implies there is **very little variability**. Every single number in the dataset must be close to the mean for the standard deviation to be small.

**Question:** If you add +5 to every number in a dataset, what happens to the Standard Deviation?

# Variance and Standard Deviation

---

**Question:** You have a dataset of 50 numbers. The standard deviation, when calculated, is very small compared to the mean. What does this imply?

→ It implies there is **very little variability**. Every single number in the dataset must be close to the mean for the standard deviation to be small.

**Question:** If you add +5 to every number in a dataset, what happens to the Standard Deviation?

→ It stays the **same**. Shifting data does not change the "spread" or width of the distribution.



# Probability Basics

# Conditional Probability Practice (Coin Toss)

---

**Problem:** A fair coin is tossed 3 times. Consider the following events:

- $A =$  "Exactly two heads occur"
- $B =$  "The first toss is Heads"

**Part 1:** List the outcomes and find  $P(A)$  and  $P(B)$ .

# Conditional Probability Practice (Coin Toss)

---

**Problem:** A fair coin is tossed 3 times. Consider the following events:

- $A =$  "Exactly two heads occur"
- $B =$  "The first toss is Heads"

**Part 1:** List the outcomes and find  $P(A)$  and  $P(B)$ .

- Total outcomes ( $2^3 = 8$ ):  $\{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$

# Conditional Probability Practice (Coin Toss)

---

**Problem:** A fair coin is tossed 3 times. Consider the following events:

- $A =$  "Exactly two heads occur"
- $B =$  "The first toss is Heads"

**Part 1:** List the outcomes and find  $P(A)$  and  $P(B)$ .

- Total outcomes ( $2^3 = 8$ ):  $\{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$
- Event  $A$  (Exactly 2 Heads):  $\{HHT, HTH, THH\}$ .  
 $\rightarrow |A| = 3$ , so  $P(A) = 3/8$

# Conditional Probability Practice (Coin Toss)

---

**Problem:** A fair coin is tossed 3 times. Consider the following events:

- $A =$  "Exactly two heads occur"
- $B =$  "The first toss is Heads"

**Part 1:** List the outcomes and find  $P(A)$  and  $P(B)$ .

- Total outcomes ( $2^3 = 8$ ):  $\{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$
- Event  $A$  (Exactly 2 Heads):  $\{HHT, HTH, THH\}$ .  
→  $|A| = 3$ , so  $P(A) = 3/8$
- Event  $B$  (First is Heads):  $\{HHH, HHT, HTH, HTT\}$ .  
→  $|B| = 4$ , so  $P(B) = 4/8 = 1/2$

## Conditional Probability Practice (Continued)

---

**Part 2:** Find  $P(A|B)$  and  $P(B|A)$ .

## Conditional Probability Practice (Continued)

---

**Part 2:** Find  $P(A|B)$  and  $P(B|A)$ .

**Step 1: Find the Intersection  $A \cap B$**

- Outcomes in  $A$  (Exactly 2 H) **AND** in  $B$  (Start with H):
- $A \cap B = \{HHT, HTH\}$
- $P(A \cap B) = 2/8 = 1/4$

# Conditional Probability Practice (Continued)

---

**Part 2:** Find  $P(A|B)$  and  $P(B|A)$ .

**Step 1: Find the Intersection  $A \cap B$**

- Outcomes in  $A$  (Exactly 2 H) **AND** in  $B$  (Start with H):
- $A \cap B = \{HHT, HTH\}$
- $P(A \cap B) = 2/8 = 1/4$

**Step 2: Apply Formula**

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{2/8}{4/8} = \frac{2}{4} = \frac{1}{2}$$

$$P(B|A) = \frac{P(A \cap B)}{P(A)} = \frac{2/8}{3/8} = \frac{2}{3}$$



# Unions and Independence

---

**Union Rule:**  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ .

**Question:** Let  $P(A) = 0.4$  and  $P(B) = 0.5$ .

1. What is the maximum possible value of  $P(A \cup B)$ ?
2. If  $A$  and  $B$  are independent, what is  $P(A \cup B)$ ?

# Unions and Independence

---

**Union Rule:**  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ .

**Question:** Let  $P(A) = 0.4$  and  $P(B) = 0.5$ .

1. What is the maximum possible value of  $P(A \cup B)$ ?
2. If  $A$  and  $B$  are independent, what is  $P(A \cup B)$ ?

1. Max value occurs if  $A$  and  $B$  are disjoint (mutually exclusive).

$P(A \cup B) = 0.4 + 0.5 = 0.9$ . (Note: It cannot exceed 1 since it is a probability).

# Unions and Independence

---

**Union Rule:**  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ .

**Question:** Let  $P(A) = 0.4$  and  $P(B) = 0.5$ .

1. What is the maximum possible value of  $P(A \cup B)$ ?
2. If  $A$  and  $B$  are independent, what is  $P(A \cup B)$ ?

1. Max value occurs if  $A$  and  $B$  are disjoint (mutually exclusive).

$P(A \cup B) = 0.4 + 0.5 = 0.9$ . (Note: It cannot exceed 1 since it is a probability).

2. If independent,  $P(A \cap B) = P(A)P(B) = 0.4 \times 0.5 = 0.2$ .

$P(A \cup B) = 0.4 + 0.5 - 0.2 = 0.7$ .

# Independence vs Correlation

---

**True or False:** If the correlation coefficient  $\rho_{XY} = 0$ , then  $X$  and  $Y$  are independent.

## Independence vs Correlation

---

**True or False:** If the correlation coefficient  $\rho_{XY} = 0$ , then  $X$  and  $Y$  are independent.

**Answer: False.**

# Independence vs Correlation

---

**True or False:** If the correlation coefficient  $\rho_{XY} = 0$ , then  $X$  and  $Y$  are independent.

**Answer: False.**

**Practical Reasoning:**

- **Correlation** ( $\rho$ ) is a linearity score. It only tests if a *straight line* fits the data.
- **Independence** means there is *no relationship whatsoever*.

**What about a Parabola?** Consider  $Y = X^2$  (a perfect U-shape).

- Are they related? **Yes**, perfectly.
- Is the correlation zero? **Yes**, because a straight line fits a U-shape poorly.

# Combinatorics

# Counting Rules

---

**Permutations (Order Matters):**  $P_k^n = \frac{n!}{(n-k)!}$

**Combinations (Order Doesn't Matter):**  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$

**Practice:**

1. How many ways can you arrange 4 distinct books on a shelf?
2. How many ways can you choose 3 committee members from 10 people?



# Counting Rules

---

**Permutations (Order Matters):**  $P_k^n = \frac{n!}{(n-k)!}$

**Combinations (Order Doesn't Matter):**  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$

**Practice:**

1. How many ways can you arrange 4 distinct books on a shelf?
2. How many ways can you choose 3 committee members from 10 people?

1.  $4! = 4 \times 3 \times 2 \times 1 = 24.$

# Counting Rules

---

**Permutations (Order Matters):**  $P_k^n = \frac{n!}{(n-k)!}$

**Combinations (Order Doesn't Matter):**  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$

## Practice:

1. How many ways can you arrange 4 distinct books on a shelf?
2. How many ways can you choose 3 committee members from 10 people?

1.  $4! = 4 \times 3 \times 2 \times 1 = 24.$

2. Order doesn't matter (committee members are equal). Use Combinations.  
 $\binom{10}{3} = \frac{10 \times 9 \times 8}{3 \times 2 \times 1} = 120.$

# The Urn Problem

---

You have a box with 6 red balls and 4 green balls. You pick 3 balls at random (no replacement).

**Question:** What is the probability of getting exactly 2 red balls?

# The Urn Problem

---

You have a box with 6 red balls and 4 green balls. You pick 3 balls at random (no replacement).

**Question:** What is the probability of getting exactly 2 red balls?

- Total ways to pick 3 balls from 10:  $\binom{10}{3}$ .
- We need 2 red from the 6 available:  $\binom{6}{2}$ .
- We need 1 green from the 4 available (to make 3 total):  $\binom{4}{1}$ .

# The Urn Problem

---

You have a box with 6 red balls and 4 green balls. You pick 3 balls at random (no replacement).

**Question:** What is the probability of getting exactly 2 red balls?

- Total ways to pick 3 balls from 10:  $\binom{10}{3}$ .
- We need 2 red from the 6 available:  $\binom{6}{2}$ .
- We need 1 green from the 4 available (to make 3 total):  $\binom{4}{1}$ .

$$\begin{aligned} P(2R, 1G) &= \frac{\binom{6}{2} \cdot \binom{4}{1}}{\binom{10}{3}} \\ &= \frac{15 \cdot 4}{120} = \frac{60}{120} = 0.5 \end{aligned}$$

# Random Variables

## Expectation and Variance Rules

---

For any random variable  $X$  and constants  $a, b$ :

$$E[aX + b] = aE[X] + b$$

$$\text{Var}(aX + b) = a^2 \text{Var}(X) \quad (\text{Shift } b \text{ does nothing to spread!})$$

**Practice:** Suppose  $E[X] = 2$  and  $\text{Var}(X) = 3$ . Find  $E[X^2]$  and  $\text{Var}(3X + 5)$ .

## Expectation and Variance Rules

---

For any random variable  $X$  and constants  $a, b$ :

$$E[aX + b] = aE[X] + b$$

$$\text{Var}(aX + b) = a^2 \text{Var}(X) \quad (\text{Shift } b \text{ does nothing to spread!})$$

**Practice:** Suppose  $E[X] = 2$  and  $\text{Var}(X) = 3$ . Find  $E[X^2]$  and  $\text{Var}(3X + 5)$ .

**1. Find  $\text{Var}(3X + 5)$ :**

$$\text{Var}(3X + 5) = 3^2 \text{Var}(X) = 9(3) = 27.$$



## Expectation and Variance Rules

---

For any random variable  $X$  and constants  $a, b$ :

$$E[aX + b] = aE[X] + b$$

$$\text{Var}(aX + b) = a^2 \text{Var}(X) \quad (\text{Shift } b \text{ does nothing to spread!})$$

**Practice:** Suppose  $E[X] = 2$  and  $\text{Var}(X) = 3$ . Find  $E[X^2]$  and  $\text{Var}(3X + 5)$ .

**1. Find  $\text{Var}(3X + 5)$ :**

$$\text{Var}(3X + 5) = 3^2 \text{Var}(X) = 9(3) = 27.$$

**2. Find  $E[X^2]$ :**

$$\text{Recall } \text{Var}(X) = E[X^2] - (E[X])^2.$$

$$3 = E[X^2] - (2)^2 \implies E[X^2] = 3 + 4 = 7.$$

# Discrete Random Variables

---

Consider the following PMF for random variable  $X$ :

| $x$        | 0   | 1   | 2 |
|------------|-----|-----|---|
| $P(X = x)$ | 0.2 | 0.5 | ? |

**Part A:** What is  $P(X = 2)$ ?

# Discrete Random Variables

---

Consider the following PMF for random variable  $X$ :

|            |     |     |   |
|------------|-----|-----|---|
| $x$        | 0   | 1   | 2 |
| $P(X = x)$ | 0.2 | 0.5 | ? |

**Part A:** What is  $P(X = 2)$ ? Probabilities must sum to 1.  $1 - (0.2 + 0.5) = 0.3$ .

**Part B:** What is the CDF value  $F(1)$ ?

# Discrete Random Variables

---

Consider the following PMF for random variable  $X$ :

| $x$        | 0   | 1   | 2 |
|------------|-----|-----|---|
| $P(X = x)$ | 0.2 | 0.5 | ? |

**Part A:** What is  $P(X = 2)$ ? Probabilities must sum to 1.  $1 - (0.2 + 0.5) = 0.3$ .

**Part B:** What is the CDF value  $F(1)$ ?

$$F(1) = P(X \leq 1) = P(X = 0) + P(X = 1) = 0.2 + 0.5 = 0.7.$$

# Joint Probabilities & Bayes

# Joint Distributions

---

**Marginal Distributions:** To find the distribution of  $X$  from a joint table, sum across the rows (or columns) of  $Y$ .

**Independence Check:**  $X$  and  $Y$  are independent **only if**  
 $P(X = x, Y = y) = P(X = x)P(Y = y)$  for **every** cell in the table.

# Joint Distributions

---

**Marginal Distributions:** To find the distribution of  $X$  from a joint table, sum across the rows (or columns) of  $Y$ .

**Independence Check:**  $X$  and  $Y$  are independent **only if**  
 $P(X = x, Y = y) = P(X = x)P(Y = y)$  for **every** cell in the table.

**Conditional Probabilities:** If you are asked to find  $P(X = 1|Y = 2)$ , for example, read the intersection from the table and divide by the marginal of  $Y$ .

$$P(X = 1|Y = 2) = \frac{P(X = 1 \cap Y = 2)}{P(Y = 2)}$$

# Bayes' Theorem Practice

---

**Scenario:** A company has two machines.

- Machine 1 produces 60% of items, with a 5% defect rate.
- Machine 2 produces 40% of items, with a 2% defect rate.

You pick an item at random and it is **defective**. What is the probability it came from Machine 1?



# Bayes' Theorem Practice

---

**Scenario:** A company has two machines.

- Machine 1 produces 60% of items, with a 5% defect rate.
- Machine 2 produces 40% of items, with a 2% defect rate.

You pick an item at random and it is **defective**. What is the probability it came from Machine 1?

- Let  $M_1$  be Machine 1,  $D$  be Defective.
- We want  $P(M_1|D)$ .

# Bayes' Theorem Practice

---

**Scenario:** A company has two machines.

- Machine 1 produces 60% of items, with a 5% defect rate.
- Machine 2 produces 40% of items, with a 2% defect rate.

You pick an item at random and it is **defective**. What is the probability it came from Machine 1?

- Let  $M_1$  be Machine 1,  $D$  be Defective.
- We want  $P(M_1|D)$ .

$$\begin{aligned} P(M_1|D) &= \frac{P(D|M_1)P(M_1)}{P(D|M_1)P(M_1) + P(D|M_2)P(M_2)} \\ &= \frac{0.05(0.60)}{0.05(0.60) + 0.02(0.40)} = \frac{0.030}{0.030 + 0.008} = \frac{0.030}{0.038} \approx 0.789 \end{aligned}$$

# Reminders

---

## Tips:

1. Write the general formula (e.g., Bayes, Total Probability) before plugging in numbers. You cannot get partial credit without showing your work.
2. If you are stuck on a word problem with probabilities, try:
  - Defining the events which are occurring
  - Writing given probabilities in terms of events
  - Checking equations to determine what you might be able to do with what you are given
3. For continuous variables,  $\int_{-\infty}^{\infty} f(x)dx = 1$ .
4. If you get a negative probability or variance, you made a mistake.

## Other:

1. You are allowed one  $9.5'' \times 11''$  note sheet (front and back)

**Good Luck on Tuesday!**