

Mann §3.1.4 — Multiplication Tables

A Plain-Language Walkthrough (with diagrams)

Reading companion — Mann, *An Introduction to Particle Physics and the Standard Model*, Ch. 3

2026-05-20

Where we are. In §3.1.1–§3.1.3 we built up the abstract machinery: a *group* is a closed, associative set with identity and inverses; a *representation* is a concrete realization of those abstract elements as matrices; and an *irreducible* representation is one of the indivisible atomic pieces from which all representations are built. Section 3.1.4 takes a small but important detour: how do we actually *write down* all the information in a (finite) group? The answer is the **multiplication table**, and it leads naturally to the idea of the **order of an element** and **cyclic subgroups**.

The one-line idea

A multiplication table is a grid that lists $g \bullet h$ for every pair of group elements g (row) and h (column). It is a *complete* description of a finite group — every piece of structure is in there.

1. The familiar analogy: numbers

Everyone knows the grade-school times table. Pick a row x , pick a column y , the cell at the intersection is $x \times y$:

$\times$	1	2	3
1	1	2	3
2	2	4	6
3	3	6	9

A *group* multiplication table is the exact same idea — except the operation $\times$ is replaced by the group’s combining operation $\bullet$, and the entries are group elements rather than numbers.

2. A real group: $\mathbb{Z}_3$ (addition mod 3)

The simplest non-trivial example is $\mathbb{Z}_3 = \{0, 1, 2\}$ with operation “+ mod 3” (clock arithmetic, 3 hours). Identity = 0; inverses $1 \leftrightarrow 2$, $0 \leftrightarrow 0$. Its multiplication table:

+	0	1	2
0	0	1	2
1	1	2	0
2	2	0	1

row 2 + col 1 = 0

Figure 1: The multiplication table of $\mathbb{Z}_3 = \{0, 1, 2\}$ under addition mod 3. The highlighted cell shows $2 + 1 \equiv 0 \pmod{3}$.

What you can read off the table by inspection

- (a) **Closure.** Every cell is in $\{0, 1, 2\}$ — nothing escapes the group.
- (b) **Identity.** Row 0 (and column 0) is a verbatim copy of the header. That is the fingerprint of the identity.
- (c) **Inverses.** Find the identity (here 0) in each row. Row 1 has it in column 2, so $1^{-1} = 2$. Row 2 has it in column 1, so $2^{-1} = 1$.
- (d) **Abelian?** The table is *symmetric across the main diagonal* iff the group is commutative. $\mathbb{Z}_3$ is abelian.
- (e) **Latin-square property.** Every group element appears *exactly once* in each row and each column. This is true for *every* group table and is a great sanity check when constructing one.

3. Why Mann bothers to introduce tables here

There are two reasons, and the second one is the punchline:

1. Tables give us an *honest, complete* description of a finite group. They contain everything.
2. Tables are *only practical for small finite groups*. For huge or infinite groups (e.g. the rotation group $SO(3)$, or any of the Lie groups that govern particle physics), you would need an infinite table. So tables motivate a question that drives the rest of the chapter: *how do we describe groups when the table is too big to write down?* The answer, coming in §3.2–§3.3, is **generators** and **Lie algebras**.

4. Powers of a single element

Now Mann pivots to a different question: what happens if you take one element $x \in G$ and keep combining it with itself?

$$x, \quad x^2 = x \bullet x, \quad x^3 = x \bullet x \bullet x, \quad x^4, \quad \dots$$

Claim (Mann)

In a finite group this sequence eventually *cycles*, and the first time it repeats, it repeats by returning to x itself. Equivalently: there exists a smallest integer $n \geq 1$ such that $x^n = e$. This n is called the **order** of x .

The powers of x trace out a closed loop that always passes through the identity:




Figure 2: The powers of a single element x of order n form a **cyclic subgroup** $\{e, x, x^2, \dots, x^{n-1}\}$. The cycle *always* includes the identity — it cannot “lasso” partway around.

5. Mann’s tiny proof, decoded

Mann compresses the argument into about four lines. Here it is unpacked.

The hypothesis

Suppose the sequence $x, x^2, x^3, \dots$ first repeats at position p , and the repetition occurs after $q + 1$ steps. In symbols:

$$x^{q+1} = x^p, \quad \text{with } p \leq q + 1. \quad (*)$$

We want to prove $p = 1$.

Step 1. Multiply both sides of $(*)$ on the left by x^{-1} :

$$x^{-1} \bullet x^{q+1} = x^{-1} \bullet x^p \implies x^q = x^{p-1}.$$

Step 2. Look at what this new equation says. The left side x^q appears *earlier* in the sequence than x^{q+1} . The right side x^{p-1} appears *earlier* than x^p . So $x^q = x^{p-1}$ is itself a repeat *occurring before position p* .

Step 3. But p was supposed to be the *first* place a repeat happens. The only way to avoid contradicting that is for $p - 1 = 0$, i.e.

$$\boxed{p = 1.}$$

Plugging back: $x^{q+1} = x^1 = x$, hence $x^q = e$. □

Moral

The powers of x cannot fall into a cycle that skips the identity — the cycle *must* loop through

e . Equivalently, every element of a finite group generates a clean cyclic subgroup

$$\langle x \rangle = \{e, x, x^2, \dots, x^{n-1}\}.$$

Geometric picture: think of the powers of x as beads on a string. This result says the string can't form a lasso — it must form a closed loop that comes back to the starting bead e .

Concrete check in $\mathbb{Z}_3$

Take $x = 1 \in \mathbb{Z}_3$:

$$1, \quad 1 + 1 = 2, \quad 1 + 1 + 1 = 0 = e.$$

So $1^3 = e$ — the order of 1 is 3. The cycle $e \rightarrow 1 \rightarrow 2 \rightarrow e$ in fact sweeps through *every* element of the group. An element whose powers generate the entire group is called a **generator**; a group possessing such an element is called **cyclic**. $\mathbb{Z}_3$ is cyclic, and 1 is a generator.

6. Looking ahead

The picture “element x + its powers + a rule that loops back to e ” is the finite-group baby version of what becomes, in continuous (Lie) groups, the picture

$$\text{generator } T \longrightarrow \text{one-parameter subgroup } e^{i\theta T},$$

where instead of stepping through discrete powers $x, x^2, x^3, \dots$ we slide a continuous parameter θ . Sections 3.2 and 3.3, which we turn to next, upgrade the cyclic-subgroup idea from finite loops to smooth curves — and that is the door to Lie algebras and to the symmetry groups that organize particle physics.

Five-bullet summary

- A multiplication table is a complete encoding of a finite group.
- The table makes closure, identity, inverses, and commutativity visible at a glance; every row and every column is a permutation of the group's elements (Latin-square property).
- Tables are impractical for large or infinite groups — this motivates generators and Lie algebras later.
- Powers of any element x in a finite group cycle back to the identity after a smallest number n of steps; n is the **order** of x .
- The cycle *always* contains the identity (Mann's short proof); the powers form a cyclic subgroup $\{e, x, x^2, \dots, x^{n-1}\}$.