

Local Gauge Invariance, Its Grand Consequences, and the Higgs Mechanism

Teaching Notes for PHYS 7502

Covering: Griffiths Ch. 10 (end), Thomson Ch. 10 & 17, Mann Ch. 12 & 23, Lancaster & Blundell

Motivating Question (F4)

What are the two biggest benefits of being able to show that QED and QCD can be generated by the principle of local gauge invariance? How does the Higgs mechanism use the Higgs field to give masses to the weak gauge bosons? Why is the fact that ϕ^4 theory is renormalizable relevant?

Contents

1	The Big Picture: Why Local Gauge Invariance Matters	2
2	Constructing QED from Local U(1) Gauge Invariance	2
2.1	The Free Dirac Equation	2
2.2	The Local Phase Transformation	2
2.3	Why the Free Lagrangian Fails	3
2.4	The Fix: Introduce a Gauge Field	3
3	From QED to QCD: Local SU(3) Gauge Invariance	4
3.1	The Local SU(3) Transformation	4
3.2	The Non-Abelian Twist: Gluon Self-Interactions	4
4	The Two Grand Benefits of Local Gauge Invariance	6
5	The Problem: Gauge Bosons Must Be Massless	8
6	Spontaneous Symmetry Breaking: Warm-Up	9
6.1	A Scalar Field with a “Mexican Hat” Potential	9
6.2	Expanding Around the Vacuum	9
7	The Higgs Mechanism: U(1) Prototype	11
7.1	The Setup	11
7.2	Breaking the Symmetry	11
7.3	The Magic: What Happens to the Lagrangian	11

8	The Standard Model Higgs: Breaking $SU(2)_L \times U(1)_Y$	13
8.1	The Higgs Doublet	13
8.2	Symmetry Breaking and the Vacuum	13
8.3	The Covariant Derivative and Mass Generation	13
9	Fermion Masses via the Higgs	15
10	Why ϕ^4 and Renormalizability Matter	16
10.1	The Power-Counting Argument	16
10.2	Connection to the Higgs Mechanism	17
11	Summary: The Higgs Mechanism at a Glance	19

1 The Big Picture: Why Local Gauge Invariance Matters

Before we get into the math, let's set up the story. Imagine you have the Dirac equation for a free electron. It describes a fermion moving through space, minding its own business, not interacting with anything. Now you ask: *what if I demand that the physics not change when I multiply the wavefunction by a phase that can be different at every point in spacetime?* This sounds like an innocent mathematical requirement, but it turns out to have enormous physical consequences. That's the principle of **local gauge invariance**.

Analogy: The Currency Exchange

Think of a “global” gauge transformation as converting all prices in a country from dollars to euros using a single exchange rate. Nothing physical changes. A “local” gauge transformation is like allowing *every city* to use its own exchange rate. Suddenly you need a network of currency exchange offices (= gauge fields) to keep trade consistent. The *existence* and *rules* of those exchange offices are forced on you by requiring economic consistency.

2 Constructing QED from Local U(1) Gauge Invariance

2.1 The Free Dirac Equation

The Lagrangian for a free spin- $\frac{1}{2}$ particle is:

$$\mathcal{L}_{\text{free}} = \bar{\psi}(i\gamma^\mu\partial_\mu - m)\psi \quad (1)$$

Term Refresher: Lagrangian Density

In quantum field theory, the **Lagrangian density** $\mathcal{L}$ is a function of the fields and their derivatives. It plays the same role as the Lagrangian $L = T - V$ in classical mechanics: the equations of motion (here, the field equations like the Dirac equation) follow from the Euler–Lagrange equations applied to $\mathcal{L}$. The term $\bar{\psi}(i\gamma^\mu\partial_\mu)\psi$ is the kinetic piece, and $-m\bar{\psi}\psi$ is the mass term.

2.2 The Local Phase Transformation

We now *demand* that $\mathcal{L}$ be invariant under a **local** $U(1)$ phase transformation:

$$\psi(x) \longrightarrow \psi'(x) = e^{iq\chi(x)}\psi(x), \quad (2)$$

where q is the charge, and $\chi(x)$ is an *arbitrary function of spacetime*.

Global vs. Local

If χ were a constant (same everywhere), this would be a *global* transformation and $\mathcal{L}_{\text{free}}$ is trivially invariant. Making χ depend on x is what makes it *local*, and this is where the trouble (and the magic) begins.

2.3 Why the Free Lagrangian Fails

Under the transformation (2), the derivative term picks up an extra piece:

$$\partial_\mu \psi \longrightarrow e^{iq\chi} (\partial_\mu \psi + iq(\partial_\mu \chi)\psi). \quad (3)$$

The $iq(\partial_\mu \chi)\psi$ term does *not* cancel. Therefore:

$$\mathcal{L}_{\text{free}} \longrightarrow \mathcal{L}_{\text{free}} - q\bar{\psi}\gamma^\mu(\partial_\mu \chi)\psi \neq \mathcal{L}_{\text{free}}. \quad (4)$$

Key Conclusion

A free-particle theory (no interactions) is *never* invariant under a local gauge transformation. To restore invariance, we *must* introduce interactions.

2.4 The Fix: Introduce a Gauge Field

We replace ordinary derivatives with **covariant derivatives**:

$$\partial_\mu \longrightarrow D_\mu \equiv \partial_\mu + iqA_\mu(x), \quad (5)$$

where $A_\mu(x)$ is a new vector field. The modified Lagrangian is:

$$\mathcal{L} = \bar{\psi}(i\gamma^\mu D_\mu - m)\psi = \bar{\psi}(i\gamma^\mu \partial_\mu - m)\psi - q\bar{\psi}\gamma^\mu A_\mu \psi. \quad (6)$$

The unwanted $(\partial_\mu \chi)$ term is cancelled *provided* A_μ transforms as:

$$A_\mu \longrightarrow A'_\mu = A_\mu - \partial_\mu \chi. \quad (7)$$

This is exactly the gauge transformation of classical electromagnetism!

QED Emerges

The full QED Lagrangian, including the kinetic term for the photon field, is:

$$\mathcal{L}_{\text{QED}} = \underbrace{\bar{\psi}(i\gamma^\mu \partial_\mu - m_e)\psi}_{\text{free electron}} + \underbrace{e\bar{\psi}\gamma^\mu \psi A_\mu}_{\text{interaction}} - \underbrace{\frac{1}{4}F_{\mu\nu}F^{\mu\nu}}_{\text{free photon}}, \quad (8)$$

where $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the electromagnetic field-strength tensor. The photon kinetic term is *already* gauge invariant on its own.

Physical Insight

The interaction term $e\bar{\psi}\gamma^\mu \psi A_\mu$ was *not* put in by hand. It was *forced* on us by demanding local $U(1)$ invariance. Furthermore, the Euler–Lagrange equations for A_μ give $\partial_\mu F^{\mu\nu} = j^\nu$, which is the covariant form of **Maxwell’s equations**. So the entirety of electromagnetism — including how charges interact and how electromagnetic waves propagate — is a *consequence* of requiring local $U(1)$ gauge symmetry.

3 From QED to QCD: Local SU(3) Gauge Invariance

The extension to QCD follows the same logic, but with a bigger symmetry group.

Term Refresher: SU(3) vs. U(1)

$U(1)$ transformations are just phase rotations $e^{i\alpha}$ — they commute ($e^{i\alpha}e^{i\beta} = e^{i\beta}e^{i\alpha}$), making $U(1)$ an **Abelian** group. $SU(3)$ transformations act on three-component “colour” vectors using 3×3 unitary matrices with unit determinant. Their generators (the eight Gell-Mann matrices λ_a , $a = 1, \dots, 8$) do *not* commute: $[\lambda_i, \lambda_j] = 2if_{ijk}\lambda_k$. This makes $SU(3)$ a **non-Abelian** group.

3.1 The Local SU(3) Transformation

We demand invariance under:

$$\psi(x) \longrightarrow \psi'(x) = \exp(ig_s \boldsymbol{\alpha}(x) \cdot \mathbf{T}) \psi(x), \quad (9)$$

where $\mathbf{T} = \{T_a\} = \{\frac{1}{2}\lambda_a\}$ are the eight generators of $SU(3)$, and $\boldsymbol{\alpha}(x) = (\alpha_1(x), \dots, \alpha_8(x))$ are eight *spacetime-dependent* functions.

The covariant derivative now requires *eight* gauge fields G_μ^a (one for each generator):

$$D_\mu = \partial_\mu + ig_s T_a G_\mu^a(x). \quad (10)$$

QCD Emerges

The locally gauge-invariant Dirac equation for a quark becomes:

$$i\gamma^\mu (\partial_\mu + ig_s T_a G_\mu^a) \psi - m\psi = 0. \quad (11)$$



The eight fields G_μ^a are the **gluons**. The form of the quark–gluon interaction vertex ($g_s T_a \gamma^\mu$) is entirely dictated by the gauge symmetry.

3.2 The Non-Abelian Twist: Gluon Self-Interactions

Because the $SU(3)$ generators do not commute, the gauge fields must transform as:

$$G_\mu^k \longrightarrow G_\mu^{k'} = G_\mu^k - \partial_\mu \alpha_k - g_s f_{ijk} \alpha_i G_\mu^j. \quad (12)$$

The last term (absent in QED!) means that the gauge fields themselves carry colour charge. This leads to **triple and quartic gluon self-interaction vertices**, whose forms are completely fixed by the $SU(3)$ gauge symmetry.



Why Gluon Self-Interaction Is Physical

In QED, photons carry no electric charge, so they don't interact with each other directly. In QCD, gluons carry colour charge, so they *do* interact with each other. This is the origin of colour confinement and asymptotic freedom — the defining features of the strong interaction. None of this was put in by hand; it all follows from the non-Abelian structure of the gauge group.

4 The Two Grand Benefits of Local Gauge Invariance

We can now clearly state the two biggest payoffs of the gauge principle.

Benefit #1: A Machine for Determining Interactions

Local gauge invariance is not merely a symmetry — it is a **constructive principle**. Given only:

- (a) the gauge group ($U(1)$ for QED, $SU(3)$ for QCD, $SU(2)_L$ for weak), and
- (b) the representations of the matter fields (how quarks/leptons transform),

the principle *uniquely determines* the form of all interaction vertices, including the coupling of matter to gauge bosons, and (for non-Abelian theories) the self-interactions of the gauge bosons themselves.

Before the gauge principle, constructing $\mathcal{L}_{\text{int}}$ was an *ad hoc* guess. Now it is derived from a symmetry requirement. As Griffiths says (end of Ch. 10): the principle gives us a “machine for determining the couplings.”

Benefit #2: Gauge Theories Are Renormalizable

In the early 1970s, Gerard 't Hooft proved that **all gauge theories are renormalizable**.

A **renormalizable** theory is one in which all infinities that appear in loop (higher-order) Feynman diagrams can be absorbed into a redefinition of a *finite* number of physical parameters (masses and couplings). This means:

- (a) Predictions beyond lowest order are *finite and unambiguous*.
- (b) The theory has genuine **predictive power**: once you measure a finite set of inputs, everything else is calculable.

A non-renormalizable theory, by contrast, generates *new* types of infinities at every order of perturbation theory, requiring infinitely many parameters — effectively making it unpredictable at high energies. The classic example is Fermi’s original four-fermion theory of the weak interaction, whose coupling G_F has dimensions of $[\text{mass}]^{-2}$.

Why These Two Benefits Are “The Biggest”

Together, these two facts mean that if you can show a theory is a gauge theory, you simultaneously know:

1. *What* the interactions look like (Benefit #1), and
2. That the theory is *mathematically consistent* to all orders of perturbation theory (Benefit #2).

This is why the Standard Model — $SU(3)_C \times SU(2)_L \times U(1)_Y$ — is built entirely on gauge principles. Anything less would either be arbitrary or inconsistent.

5 The Problem: Gauge Bosons Must Be Massless

There is a catch. The gauge principle works beautifully, but it requires the gauge fields to be **massless**. Here's why.

A mass term for a vector boson looks like:

$$\mathcal{L}_{\text{mass}} = \frac{1}{2}m^2 A_\mu A^\mu. \quad (13)$$

Under $A_\mu \rightarrow A_\mu - \partial_\mu \chi$:

$$\frac{1}{2}m^2 A_\mu A^\mu \longrightarrow \frac{1}{2}m^2 (A_\mu - \partial_\mu \chi)(A^\mu - \partial^\mu \chi) \neq \frac{1}{2}m^2 A_\mu A^\mu. \quad (14)$$

The Mass–Gauge Dilemma

A boson mass term *breaks* gauge invariance. So gauge invariance requires massless gauge bosons. This is perfectly fine for:

- QED: the photon is massless. ✓
- QCD: gluons are massless. ✓

But the weak interaction has massive gauge bosons: $m_W \approx 80 \text{ GeV}$, $m_Z \approx 91 \text{ GeV}$. ✗

If we just force a mass term into the weak-interaction Lagrangian, the gauge invariance is broken, and by 't Hooft's theorem the theory loses renormalizability (and therefore predictive power). This is exactly what happens with Fermi's theory.

The Higgs mechanism resolves this by generating gauge boson masses in a way that *preserves* the underlying gauge symmetry.

6 Spontaneous Symmetry Breaking: Warm-Up

6.1 A Scalar Field with a “Mexican Hat” Potential



Term Refresher: Scalar Field

A **scalar field** $\phi(x)$ has spin-0. Its Lagrangian involves $(\partial_\mu \phi)(\partial^\mu \phi)$ (kinetic energy) and $V(\phi)$ (potential energy). The Higgs boson is a scalar particle.

Consider a complex scalar field $\phi = \frac{1}{\sqrt{2}}(\phi_1 + i\phi_2)$ with the potential:

$$V(\phi) = \mu^2(\phi^* \phi) + \lambda(\phi^* \phi)^2, \quad \lambda > 0. \quad (15)$$

There are two cases:



When $\mu^2 < 0$, the minimum is *not* at $\phi = 0$ but at:

$$|\phi_0|^2 = \frac{-\mu^2}{2\lambda} \equiv \frac{v^2}{2}, \quad v = \sqrt{\frac{-\mu^2}{\lambda}}. \quad (16)$$

The quantity v is called the **vacuum expectation value** (vev) of the field.

6.2 Expanding Around the Vacuum

We choose the vacuum to lie along the real direction: $\langle \phi \rangle = v/\sqrt{2}$. Then we expand:

$$\phi(x) = \frac{1}{\sqrt{2}}(v + \eta(x) + i\xi(x)), \quad (17)$$

where $\eta(x)$ represents oscillations in the “radial” direction (where the potential is curved) and $\xi(x)$ represents oscillations in the “angular” direction (where the potential is flat).

Substituting into $\mathcal{L}$ and collecting terms:

$$\mathcal{L} = \underbrace{\frac{1}{2}(\partial_\mu \eta)(\partial^\mu \eta) - \lambda v^2 \eta^2}_{\text{massive } \eta \text{ (mass } m_\eta = \sqrt{2\lambda} v)} + \underbrace{\frac{1}{2}(\partial_\mu \xi)(\partial^\mu \xi)}_{\text{massless } \xi} - V_{\text{int}}(\eta, \xi). \quad (18)$$

Goldstone's Theorem

Whenever a *continuous global* symmetry is spontaneously broken, a massless scalar particle (a **Goldstone boson**) appears. Here, that's ξ . It corresponds to oscillations along the flat direction of the potential.

The Problem with Goldstone Bosons

No massless scalar particles have ever been observed in nature. If we stopped here, the theory would predict something that doesn't exist. This is resolved beautifully when we combine spontaneous symmetry breaking with *local* gauge invariance — the Higgs mechanism.

7 The Higgs Mechanism: U(1) Prototype

Now we combine the two ideas: a complex scalar field with $\mu^2 < 0$ and local $U(1)$ gauge invariance.

7.1 The Setup

Start with the Lagrangian:

$$\mathcal{L} = (D_\mu \phi)^*(D^\mu \phi) - \mu^2 \phi^* \phi - \lambda(\phi^* \phi)^2 - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}, \quad (19)$$

where $D_\mu = \partial_\mu + igB_\mu$ is the covariant derivative, and B_μ is a (initially massless) gauge field.

7.2 Breaking the Symmetry

With $\mu^2 < 0$, the vacuum sits at $|\phi| = v/\sqrt{2}$. Choose the vacuum along the real axis, and work in the **unitary gauge** (this is a specific gauge choice that eliminates the Goldstone boson explicitly):

$$\phi(x) = \frac{1}{\sqrt{2}}(v + h(x)), \quad (20)$$

where $h(x)$ is the physical **Higgs field** describing oscillations about the vacuum.

7.3 The Magic: What Happens to the Lagrangian

Substituting into $\mathcal{L}$ and expanding (the calculation is detailed in Thomson §17.5.3):

The Higgs Mechanism Result (U(1) case)

$$\mathcal{L} = \underbrace{\frac{1}{2}(\partial_\mu h)(\partial^\mu h) - \lambda v^2 h^2}_{\text{massive Higgs boson}} - \underbrace{\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \frac{1}{2}g^2 v^2 B_\mu B^\mu}_{\text{massive gauge boson!}} + \text{interactions} \quad (21)$$

- The Higgs boson has mass $m_h = \sqrt{2\lambda} v$.
- The gauge boson has mass $m_B = gv$.
- The Goldstone boson ξ has *disappeared*.

Where Did the Goldstone Boson Go?

A massless vector boson has **2 polarization states** (transverse). A massive vector boson has **3 polarization states** (two transverse + one longitudinal). The Goldstone boson has been “eaten” by the gauge field, providing the extra longitudinal degree of freedom needed for a massive vector boson.

Degree-of-freedom counting:

Before SSB:

Massless gauge boson (2 d.o.f.)

Complex scalar (2 real d.o.f.)

Total: 4 d.o.f.

After SSB:

Massive gauge boson (3 d.o.f.)

Massive Higgs (1 d.o.f.)

Total: 4 d.o.f. ✓

The total number of degrees of freedom is conserved. Nothing was created or destroyed — just rearranged.

The Crucial Point

The gauge boson mass term $\frac{1}{2}g^2v^2B_\mu B^\mu$ did *not* break gauge invariance. It arose from the gauge-invariant interaction $|D_\mu\phi|^2$ when the field ϕ acquired a nonzero vacuum expectation value. The Lagrangian is *still gauge-invariant* — the symmetry is merely “hidden,” not truly broken. This is why the theory remains renormalizable.

8 The Standard Model Higgs: Breaking $SU(2)_L \times U(1)_Y$

The full Standard Model applies the same idea, but now to the electroweak gauge group $SU(2)_L \times U(1)_Y$, which has **four** gauge bosons: three from $SU(2)_L$ ($W_\mu^1, W_\mu^2, W_\mu^3$) and one from $U(1)_Y$ (B_μ).

8.1 The Higgs Doublet

We need to give masses to three gauge bosons ($W^\pm$ and Z) while keeping one massless (the photon γ). This requires *three* Goldstone bosons. Since each complex scalar provides two real degrees of freedom, the minimal choice is a **doublet of complex scalar fields**:

$$\phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} \phi_1 + i\phi_2 \\ \phi_3 + i\phi_4 \end{pmatrix}. \quad (22)$$

This doublet has four real degrees of freedom: three will become the longitudinal polarizations of W^+ , W^- , and Z , and one will become the physical Higgs boson h .

Why a Doublet?

The Higgs field must be able to couple to the $SU(2)_L$ gauge bosons (to give them mass via the covariant derivative). This means it must transform nontrivially under $SU(2)_L$, i.e., it must carry weak isospin. The simplest choice is $I_W = \frac{1}{2}$ (a doublet). The upper component is chosen to be charged (ϕ^+) and the lower neutral (ϕ^0) so that the vacuum can be electrically neutral.

8.2 Symmetry Breaking and the Vacuum

The Higgs potential is:

$$V(\phi) = \mu^2(\phi^\dagger\phi) + \lambda(\phi^\dagger\phi)^2, \quad \mu^2 < 0, \lambda > 0. \quad (23)$$

The vacuum is chosen to be:

$$\langle \phi \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix}, \quad v = \sqrt{\frac{-\mu^2}{\lambda}} \approx 246 \text{ GeV}. \quad (24)$$

Why the Neutral Component Gets the vev

If the charged component ϕ^+ had a nonzero vacuum expectation value, the vacuum would carry electric charge. This would break $U(1)_{\text{EM}}$ and the photon would become massive. Since we observe a massless photon, only the neutral component can develop a vev.

8.3 The Covariant Derivative and Mass Generation

In the unitary gauge, $\phi(x) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v+h(x) \end{pmatrix}$. The covariant derivative is:

$$D_\mu \phi = \left(\partial_\mu + ig_W \mathbf{T} \cdot \mathbf{W}_\mu + i\frac{g'}{2} B_\mu \right) \phi, \quad (25)$$

where $\mathbf{T} = \frac{1}{2}\boldsymbol{\sigma}$ are the $SU(2)$ generators (Pauli matrices) and g' is the $U(1)_Y$ coupling.

The term $(D_\mu\phi)^\dagger(D^\mu\phi)$, when expanded with $\phi = \frac{1}{\sqrt{2}}\begin{pmatrix} 0 \\ v+h \end{pmatrix}$, produces mass terms for the gauge bosons.

The Gauge Boson Masses

After working through the algebra (see Thomson §17.5.4):

$$m_W = \frac{1}{2}g_W v \approx 80 \text{ GeV}, \quad (26)$$

$$m_Z = \frac{1}{2} \frac{g_W}{\cos \theta_W} v = \frac{m_W}{\cos \theta_W} \approx 91 \text{ GeV}, \quad (27)$$

$$m_\gamma = 0. \quad (28)$$

The physical fields are mixtures of the underlying gauge fields:

$$W_\mu^\pm = \frac{1}{\sqrt{2}}(W_\mu^1 \mp iW_\mu^2), \quad (29)$$

$$Z_\mu = \cos \theta_W W_\mu^3 - \sin \theta_W B_\mu, \quad (30)$$

$$A_\mu = \sin \theta_W W_\mu^3 + \cos \theta_W B_\mu. \quad (31)$$

Here $\tan \theta_W = g'/g_W$ is the **weak mixing angle**.

The Prediction $m_W/m_Z = \cos \theta_W$

This relation between the W and Z masses is a *prediction* of the minimal Higgs model (one doublet). Its experimental verification at LEP and the Tevatron was a major triumph and provides compelling evidence for the reality of the Higgs mechanism.

9 Fermion Masses via the Higgs

The mass problem is not restricted to gauge bosons. In the $SU(2)_L$ gauge theory, a fermion mass term $-m\bar{\psi}\psi = -m(\bar{\psi}_R\psi_L + \bar{\psi}_L\psi_R)$ couples left-handed doublets to right-handed singlets. Since these transform differently under $SU(2)_L$, this term also breaks gauge invariance.

Yukawa Couplings

Fermion masses are generated through **Yukawa couplings** — gauge-invariant interactions between the fermion fields and the Higgs doublet:

$$\mathcal{L}_{\text{Yukawa}} = -g_f \bar{L} \phi R + \text{h.c.} \quad (32)$$

When ϕ acquires its vev ($\phi \rightarrow \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v+h \end{pmatrix}$), this becomes:

$$\mathcal{L}_{\text{Yukawa}} \longrightarrow -\frac{g_f v}{\sqrt{2}} \bar{f} f - \frac{g_f}{\sqrt{2}} \bar{f} f h. \quad (33)$$

The first term is a mass: $m_f = g_f v / \sqrt{2}$. The second is a Higgs–fermion–fermion coupling proportional to m_f .

Limitation

The Yukawa coupling constants g_f are free parameters — one for each fermion. The Higgs mechanism tells us *how* fermions get masses but doesn't predict *what* those masses are. This is one of the unsatisfying aspects of the Standard Model.

10 Why ϕ^4 and Renormalizability Matter

This section addresses the last part of the question: *why is it relevant that a scalar field ϕ with ϕ^4 interactions is renormalizable?*

10.1 The Power-Counting Argument

Term Refresher: Renormalizability Criterion

A theory is renormalizable if its coupling constant is **dimensionless** (in natural units $\hbar = c = 1$). The reasoning is:

- (i) The action $S = \int d^4x \mathcal{L}$ must be dimensionless.
- (ii) Therefore $[\mathcal{L}] = [\text{mass}]^4$.
- (iii) For a scalar field, $[\phi] = [\text{mass}]^1$ (from the kinetic term $(\partial_\mu \phi)^2$).

Now examine the ϕ^4 interaction term: $\mathcal{L}_{\text{int}} = \frac{\lambda}{4!} \phi^4$. Since $[\phi^4] = [\text{mass}]^4 = [\mathcal{L}]$, the coupling λ must be **dimensionless**.

Compare with Fermi's four-fermion theory: $\mathcal{L} \sim G_F (\bar{\psi}\psi)^2$. Since $[\psi] = [\text{mass}]^{3/2}$, we get $[(\bar{\psi}\psi)^2] = [\text{mass}]^6$, so $[G_F] = [\text{mass}]^{-2}$. The coupling has *negative* mass dimension $\Rightarrow$ non-renormalizable.

The Key Facts about ϕ^4 Theory

1. ϕ^4 theory is **renormalizable**: only a finite number of types of divergent diagrams appear (those with ≤ 4 external legs), and all infinities can be absorbed into redefinitions of m and λ .
2. ϕ^4 is the **most general renormalizable self-interaction** for a scalar field in 4 spacetime dimensions. A ϕ^5 term would require a coupling with $[\text{mass}]^{-1}$ (non-renormalizable). A ϕ^3 term can be absorbed by a field redefinition.
3. The Higgs potential $V(\phi) = \mu^2 \phi^2 + \lambda \phi^4$ is precisely a ϕ^4 theory.

10.2 Connection to the Higgs Mechanism

Why This Matters for the Higgs

The relevance of ϕ^4 renormalizability to the Higgs mechanism is this:


The Higgs potential must be renormalizable for the entire Standard Model to remain renormalizable.

The whole point of the Higgs mechanism is to give masses to gauge bosons *without* breaking the gauge symmetry that guarantees renormalizability. But if the Higgs potential itself were non-renormalizable (say, ϕ^6), we would re-introduce the very infinities we were trying to avoid.

The fact that ϕ^4 interactions are renormalizable means the Higgs sector does not spoil the beautiful renormalizability of the gauge theory. This is not a coincidence — it's a **consistency requirement**. Griffiths notes (end of Ch. 10) that the potential “has to be quartic in the fields” for renormalizability. Lancaster & Blundell use ϕ^4 theory extensively as their go-to example precisely because it is the simplest interacting scalar field theory that is renormalizable.

The Chain of Logic

Here is the complete logical chain, from symmetry to prediction:



11 Summary: The Higgs Mechanism at a Glance

c	
Ingredient	Role
Higgs doublet $\phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}$	Four real scalar fields, transforming as an $SU(2)_L$ doublet
Higgs potential $V = \mu^2 \phi^\dagger \phi + \lambda (\phi^\dagger \phi)^2$	With $\mu^2 < 0$: creates a nonzero vacuum expectation value $v \approx 246$ GeV
Vacuum expectation value	$\langle \phi \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix}$: breaks $SU(2)_L \times U(1)_Y$ to $U(1)_{EM}$
Three Goldstone bosons	“Eaten” by W^+ , W^- , and Z , providing their longitudinal polarization
Physical Higgs boson h	The remaining degree of freedom. Discovered at the LHC in 2012
Photon	Remains massless: the combination of W^3 and B corresponding to $U(1)_{EM}$
Fermion masses	Generated through Yukawa couplings $g_f \bar{L} \phi R$; where L, R are fermion doublets and singlets

The Bottom Line

The principle of local gauge invariance gives us two enormous gifts: (1) it determines the form of all interactions, and (2) it guarantees renormalizability (predictive power). The Higgs mechanism extends this framework to accommodate massive gauge bosons by introducing a scalar field with a ϕ^4 potential that spontaneously breaks the electroweak symmetry. The use of ϕ^4 (and not higher powers) is essential because only ϕ^4 is renormalizable in four spacetime dimensions. Together, gauge invariance and the Higgs mechanism provide the theoretical foundation for the entire electroweak sector of the Standard Model.

Sources: Griffiths Ch. 10–11; Thomson Ch. 10, 17; Mann Ch. 12, 23; Lancaster & Blundell Ch. 26, 32–34.