

Existence and Uniqueness for ODEs: The Picard-Lindelöf Theorem

Math Foundations — Node 30

1 Setting

Let $D \subset \mathbb{R} \times \mathbb{R}^n$ be an open set, $(t_0, y_0) \in D$, and $f : D \rightarrow \mathbb{R}^n$ a continuous function. We study the initial value problem (IVP)

$$y'(t) = f(t, y(t)), \quad y(t_0) = y_0. \quad (1)$$

A *solution* is a differentiable curve $y : I \rightarrow \mathbb{R}^n$ on an open interval $I \ni t_0$ satisfying (1) pointwise.

2 Lipschitz continuity and Picard iteration

Definition 1 (Lipschitz in y). A function $f : D \rightarrow \mathbb{R}^n$ is *Lipschitz continuous in y* on D if there is $L \geq 0$ such that

$$\|f(t, y_1) - f(t, y_2)\| \leq L \|y_1 - y_2\| \quad \forall (t, y_1), (t, y_2) \in D.$$

The smallest such L is the *Lipschitz constant* of f .

Remark 2. If f is C^1 on D , then $\partial f / \partial y$ being bounded by L on D (in operator norm) implies f is Lipschitz in y with constant L . This is the most common way Lipschitz hypotheses arise in practice.

Definition 3 (Picard iteration). Given the IVP (1), define the sequence of functions $\{y_k\}_{k \geq 0}$ by

$$y_0(t) \equiv y_0, \quad y_{k+1}(t) = y_0 + \int_{t_0}^t f(s, y_k(s)) ds.$$

These are the *Picard iterates* of the IVP.

A continuous function y solves (1) if and only if it is a fixed point of the integral operator $T[y](t) = y_0 + \int_{t_0}^t f(s, y(s)) ds$. So existence and uniqueness become a fixed-point problem for T .

3 The Picard-Lindelöf theorem

Theorem 4 (Picard-Lindelöf). Let $a, b > 0$ and let

$$R = \{(t, y) \in \mathbb{R} \times \mathbb{R}^n : |t - t_0| \leq a, \|y - y_0\| \leq b\}.$$

Suppose f is continuous on R with $\|f(t, y)\| \leq M$ for all $(t, y) \in R$, and Lipschitz in y on R with constant L . Set

$$\alpha = \min\left(a, \frac{b}{M}\right).$$

Then the IVP (1) has a unique solution $y : [t_0 - \alpha, t_0 + \alpha] \rightarrow \mathbb{R}^n$, and the Picard iterates y_k converge to it uniformly on that interval.

Proof sketch. Set $I = [t_0 - \alpha, t_0 + \alpha]$ and let

$$X = \{y \in C(I, \mathbb{R}^n) : \|y(t) - y_0\| \leq b \text{ for all } t \in I\}.$$

Equip $C(I, \mathbb{R}^n)$ with the sup norm $\|y\|_\infty = \sup_{t \in I} \|y(t)\|$. Then X is a closed subset of a Banach space, hence complete. Define $T : X \rightarrow C(I, \mathbb{R}^n)$ by

$$T[y](t) = y_0 + \int_{t_0}^t f(s, y(s)) ds.$$

Step 1 (T maps X into X). For $y \in X$ and $t \in I$,

$$\|T[y](t) - y_0\| \leq \int_{t_0}^t \|f(s, y(s))\| ds \leq M|t - t_0| \leq M\alpha \leq b.$$

Hence $T[y] \in X$.

Step 2 (Contraction in a weighted norm). On $C(I, \mathbb{R}^n)$ define the equivalent norm $\|y\|_w = \sup_{t \in I} e^{-2L|t-t_0|} \|y(t)\|$. Then for $y_1, y_2 \in X$ and $t \geq t_0$ in I ,

$$\begin{aligned} \|T[y_1](t) - T[y_2](t)\| &\leq \int_{t_0}^t \|f(s, y_1(s)) - f(s, y_2(s))\| ds \\ &\leq L \int_{t_0}^t \|y_1(s) - y_2(s)\| ds \\ &\leq L \int_{t_0}^t e^{2L(s-t_0)} ds \cdot \|y_1 - y_2\|_w \\ &= \frac{1}{2} (e^{2L(t-t_0)} - 1) \|y_1 - y_2\|_w. \end{aligned}$$

Multiplying by $e^{-2L(t-t_0)}$ and taking sup over t (the $t < t_0$ case is symmetric) yields

$$\|T[y_1] - T[y_2]\|_w \leq \frac{1}{2} \|y_1 - y_2\|_w.$$

Thus T is a strict contraction on $(X, \|\cdot\|_w)$ with contraction factor $1/2$.

Step 3 (Banach fixed point). By the Banach contraction mapping theorem, T has a unique fixed point $y^* \in X$. By construction y^* is continuous and satisfies the integral equation; since f is continuous, y^* is in fact C^1 and solves (??) pointwise on I . Uniqueness in X follows from the contraction property; any solution must lie in X (by Step 1 applied to it), so uniqueness is global on I .

Finally, the Picard iterates are exactly the sequence $T^k[y_0]$ produced by the contraction mapping theorem, and so converge uniformly on I to y^* . $\square$

4 What goes wrong without Lipschitz

Example 5 (Non-uniqueness). Consider $y' = y^{2/3}$, $y(0) = 0$. Here $f(t, y) = y^{2/3}$ fails to be Lipschitz at $y = 0$ since $\partial_y f = \frac{2}{3}y^{-1/3}$ blows up. The IVP admits the two distinct solutions $y \equiv 0$ and $y(t) = (t/3)^3$, as well as a one-parameter family y_τ that equals 0 for $t \leq \tau$ and $((t - \tau)/3)^3$ for $t > \tau$.

5 Connection to flow-based generative models

Modern generative models (flow matching, ELF, neural ODEs) define

$$\dot{x}(t) = v_\theta(t, x(t))$$

where v_θ is a neural network. On any compact set K , neural networks built from smooth activations are C^1 and hence Lipschitz in x with a finite constant L_K . Theorem ?? therefore guarantees a unique sample trajectory for each starting point x_0 , and hence a well-defined flow ϕ_t on K . Smooth dependence on initial conditions further makes ϕ_t a diffeomorphism — a property used implicitly throughout the literature.