

Gradient and Jacobian

math-foundations / concept 14

1 Setup and motivation

We work in finite-dimensional Euclidean space $\mathbb{R}^n$ with the standard inner product $\langle u, v \rangle = u^\top v$ and induced norm $\|v\| = \sqrt{\langle v, v \rangle}$. For scalar functions, the derivative captures a single rate of change; for vector-valued functions, it captures *all* pairwise rates between input and output coordinates, organized into a matrix.

2 Definitions

Definition 1 (Partial derivative). Let $U \subseteq \mathbb{R}^n$ be open and $f : U \rightarrow \mathbb{R}$. The *partial derivative* of f with respect to x_j at $x \in U$ is

$$\frac{\partial f}{\partial x_j}(x) = \lim_{h \rightarrow 0} \frac{f(x + he_j) - f(x)}{h},$$

where e_j is the j th standard basis vector, when this limit exists.

Definition 2 (Gradient). Let $f : U \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ be differentiable at x . The *gradient* of f at x is the column vector

$$\nabla f(x) = \left(\frac{\partial f}{\partial x_1}(x), \dots, \frac{\partial f}{\partial x_n}(x) \right)^\top \in \mathbb{R}^n.$$

Equivalently, $\nabla f(x)$ is the unique vector such that $D_v f(x) = \langle \nabla f(x), v \rangle$ for every $v \in \mathbb{R}^n$, where $D_v f(x) = \lim_{t \rightarrow 0} (f(x + tv) - f(x))/t$ is the directional derivative.

Definition 3 (Jacobian matrix). Let $F : U \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$ with components $F = (F_1, \dots, F_m)$, differentiable at x . The *Jacobian matrix* of F at x is the $m \times n$ matrix

$$J_F(x) = \begin{pmatrix} \frac{\partial F_1}{\partial x_1}(x) & \cdots & \frac{\partial F_1}{\partial x_n}(x) \\ \vdots & \ddots & \vdots \\ \frac{\partial F_m}{\partial x_1}(x) & \cdots & \frac{\partial F_m}{\partial x_n}(x) \end{pmatrix}, \quad [J_F(x)]_{ij} = \frac{\partial F_i}{\partial x_j}(x).$$

$J_F(x)$ is the matrix of the total derivative $dF_x : \mathbb{R}^n \rightarrow \mathbb{R}^m$ in standard bases.

Definition 4 (Jacobian determinant). If $m = n$, the *Jacobian determinant* of F at x is $\det J_F(x) \in \mathbb{R}$. It is the signed factor by which F locally scales n -dimensional volume at x : a small parallelepiped of volume V at x has image of volume $|\det J_F(x)| V + o(V)$.

3 The gradient and steepest ascent

Lemma 1 (Directional-derivative formula). *If f is differentiable at x , then for every $v \in \mathbb{R}^n$,*

$$D_v f(x) = \langle \nabla f(x), v \rangle.$$

Sketch. Differentiability at x means $f(x+h) = f(x) + \langle \nabla f(x), h \rangle + r(h)$ with $r(h)/\|h\| \rightarrow 0$. Setting $h = tv$ and dividing by t gives the claim in the limit $t \rightarrow 0$. $\square$

Theorem 2 (Direction of maximum directional derivative). *Let $f : U \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ be differentiable at $x \in U$, with $\nabla f(x) \neq 0$. Then among all unit vectors $v \in \mathbb{R}^n$ (i.e. $\|v\| = 1$), the directional derivative $D_v f(x)$ attains its maximum at*

$$v^* = \frac{\nabla f(x)}{\|\nabla f(x)\|},$$

and the maximum value is $\|\nabla f(x)\|$.

Proof. By Lemma 1, $D_v f(x) = \langle \nabla f(x), v \rangle$. The Cauchy–Schwarz inequality states that for any $u, v \in \mathbb{R}^n$,

$$|\langle u, v \rangle| \leq \|u\| \|v\|,$$

with equality iff u and v are linearly dependent. Applying this with $u = \nabla f(x)$ and a unit vector v ,

$$D_v f(x) = \langle \nabla f(x), v \rangle \leq \|\nabla f(x)\| \|v\| = \|\nabla f(x)\|.$$

Equality holds iff v is a positive scalar multiple of $\nabla f(x)$; since $\|v\| = 1$ and $\nabla f(x) \neq 0$, the unique maximizer is $v^* = \nabla f(x)/\|\nabla f(x)\|$. Substituting back gives $D_{v^*} f(x) = \langle \nabla f(x), \nabla f(x)/\|\nabla f(x)\| \rangle = \|\nabla f(x)\|$. $\square$

Example 1. For $f(x, y) = x^2 + 3y^2$, $\nabla f(x, y) = (2x, 6y)$. At $(1, 1)$, the steepest-ascent direction is $(2, 6)/\sqrt{40} = (1, 3)/\sqrt{10}$ with rate $\sqrt{40}$, while $-\nabla f$ points toward the minimum at the origin.

4 The chain rule for vector-valued maps

Theorem 3 (Chain rule). *Let $G : \mathbb{R}^p \rightarrow \mathbb{R}^n$ be differentiable at x , and $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ differentiable at $y = G(x)$. Then $H = F \circ G$ is differentiable at x and*

$$J_H(x) = J_F(G(x)) \cdot J_G(x),$$

an $m \times p$ matrix obtained as the product of an $m \times n$ matrix and an $n \times p$ matrix.

This is the algebraic backbone of reverse-mode automatic differentiation: for a deep composition $F_L \circ F_{L-1} \circ \dots \circ F_1$, the overall Jacobian factors as a product of layer Jacobians, and the loss gradient propagates by successive vector–Jacobian products.

5 Worked example: $F(x, y) = (x^2 - y, e^{xy})$

The partials are $\partial F_1/\partial x = 2x$, $\partial F_1/\partial y = -1$, $\partial F_2/\partial x = ye^{xy}$, $\partial F_2/\partial y = xe^{xy}$. Hence

$$J_F(x, y) = \begin{pmatrix} 2x & -1 \\ ye^{xy} & xe^{xy} \end{pmatrix}, \quad J_F(1, 0) = \begin{pmatrix} 2 & -1 \\ 0 & 1 \end{pmatrix}, \quad \det J_F(1, 0) = 2.$$

By the inverse function theorem, F is a local diffeomorphism near $(1, 0)$, locally doubling area.