

## CLEVELAND METROPARKS Plant Community Assessment Program: Quality Control Form



Project Label:

PCAP

Plot No:

1012

Date Sampled:

07/01/15

Lead:

CKM

Comment required if item answer is NO

Parking/Access outside of Park Boundaries:	Y	<input checked="" type="radio"/> N	If yes, write details in Comments section below
Field journals completed	<input checked="" type="radio"/> Y	N	
Site sketch made on 1:3000 map?	<input checked="" type="radio"/> Y	N	
Check cover page	<input checked="" type="radio"/> Y	N	
X-axis Bearing of plot recorded	<input checked="" type="radio"/> Y	N	
GPS coords. Recorded	<input checked="" type="radio"/> Y	N	
North direction recorded	<input checked="" type="radio"/> Y	N	
Photographs taken?	<input checked="" type="radio"/> Y	N	
Relocated Pins Mapped	<input checked="" type="radio"/> Y	N	
Plot No., Date agreement on all pages?	<input checked="" type="radio"/> Y	N	
Header data completed all pages?	<input checked="" type="radio"/> Y	N	
Cover classes recorded in all intensive modules	<input checked="" type="radio"/> Y	N	
Browse Level By Species	<input checked="" type="radio"/> Y	N	
Woody stem quality control check	<input checked="" type="radio"/> Y	N	Check every line and cross check with the Tree Cover Sheet
Invasive plant quality control check	<input checked="" type="radio"/> Y	N	NA
Ash trees mapped	<input checked="" type="radio"/> Y	N	
Completed Forest Pest/Pathogen Datasheet	<input checked="" type="radio"/> Y	N	
Cover by Strata? (confirm cover type)	<input checked="" type="radio"/> Y	N	
Soil samples collected with matching plot #.	<input checked="" type="radio"/> Y	N	NA
Cross check 2010 information	<input checked="" type="radio"/> Y	N	Highlight any changes from 2010 information
Vouchers labeled on datasheet with initials and number	<input checked="" type="radio"/> Y	N	
Vouchers labeled on collection bag	<input checked="" type="radio"/> Y	N	
Pink flags removed	<input checked="" type="radio"/> Y	N	
Data sheet QA before leaving site?	<input checked="" type="radio"/> Y	N	
Common equipment returned to tub.	<input checked="" type="radio"/> Y	N	
Data sheets scanned?	yes MB		Enter date to left 7/6/15
Final data sheets scanned?			Enter date to left
Buffer Widths measured?	<input checked="" type="radio"/> Y	N	
Web Soil Survey	<input checked="" type="radio"/> Y	N	
Voucher Location	Refrigerator	<input checked="" type="radio"/> Y	N
(# vouchers collected)	Press (#)		Enter number to left
CKM 126-134	Drier	<input checked="" type="radio"/> Y	N
	Identified	<input checked="" type="radio"/> Y	N
	Mounted	<input checked="" type="radio"/> Y	N
	Thrown away	<input checked="" type="radio"/> Y	N

## GRTS point verification: Is plot sampleable?

<input checked="" type="checkbox"/> Yes	Original GRTS point is sampleable
<input type="checkbox"/> No	Original GRTS point lands in a non-sampleable area (fill in category below)
	<input type="checkbox"/> Point falls in a water (i.e. river, lake)
	<input type="checkbox"/> Managed mowed area (i.e. golf course, picnic area, right-of-way)
	<input type="checkbox"/> Paved area (i.e. parkinglot, road)
	<input type="checkbox"/> Unsafe to sample (i.e. steep slope)
	<input type="checkbox"/> Other

## Additional Comments:

--



# CLEVELAND METROPARKS Plant Community Assessment Program - Background Data Sheet


<b>GENERAL INFORMATION</b>	
Project Label: PCAP	
Project Name: 02NC2015	
Plot Name: Interrupted Lurch	
Snake vs. Toad	
Plot No: 1012	
<input type="checkbox"/> Level 4 (no nested corners sampled)	
<input checked="" type="checkbox"/> Level 5 (nested corners sampled)	
Date (mm/dd/yyyy): 07/01/2015	
End date (if > 1 day): / /	
Party	Role**
C. Minney	Plot leader
B. Sweet	Bot. Asst.
M. Busam	Woody Tech
R. Eagle-Malone	Woody Tech
** Roles: Co-leader, Asst., Guide, Owner, Taxonomic, etc.	
PLOT NOT SAMPLED: <input type="checkbox"/> Other	
<input type="checkbox"/> Perm. water <input type="checkbox"/> Paved <input type="checkbox"/> Slope <input type="checkbox"/> Safety	
<b>SAMPLING QUALITY*</b>	
Effort Level:	subjective evaluation of how much effort put into sampling. Hurried plots may still provide good data
<input checked="" type="checkbox"/> Very thorough	
<input type="checkbox"/> Accurate	
<input type="checkbox"/> Hurried	
<b>TAXONOMIC ACCURACY</b>	
<input type="checkbox"/> high	<input type="checkbox"/> low
<input checked="" type="checkbox"/> vascular	<input type="checkbox"/> n/a
<input type="checkbox"/> bryo	<input checked="" type="checkbox"/> lichen
<b>TAXONOMIC STANDARD</b>	
Authority: G&C	Pub Date: 1998

<b>LOCATION</b>	
State: OH	County: Cuyahoga
Quadrangle: Mayfield Heights	
Local Place Name: Ox Lane + Oxbow Lane	
Landowner: CMP	
Data Confidentiality:	
Check one: <input checked="" type="checkbox"/> Public data <input type="checkbox"/> Private Data	
<input type="checkbox"/> Fuzz 100m <input type="checkbox"/> Fuzz 250m <input type="checkbox"/> Fuzz 500m	
Reason:	
If data not public why?	
Source of coordinates: <input type="checkbox"/> MAP <input checked="" type="checkbox"/> GPS	
Coordinate system:	Coord. Units
<input checked="" type="checkbox"/> Lat/Long <input type="checkbox"/> UTM <input type="checkbox"/> StatePlane	<input type="checkbox"/> deg <input type="checkbox"/> deg min
<input type="checkbox"/> Other (specify)	<input type="checkbox"/> m <input type="checkbox"/> ft
Datum: <input checked="" type="checkbox"/> NAD83/WGS84 <input type="checkbox"/> NAD27	
GPS location in plot x=0 to 5, y=-1.0, +1): x = 0 y = 0 (base of plot x=0, y=0)	
Latitude: 41.56544	
Longitude: 81.42216	
Coord. Accuracy: <input checked="" type="checkbox"/> m <input type="checkbox"/> ft	+/- 4
GPS File Name: 1012A	
Plot size for cover data: .1	(hectares)
X-axis Bearing of plot: [256]	°
Depth: (1-5): 4	
Intensive modules: 2, 3, 8, 9	(EDIT IF MODIFIED)
Camera No.: 4	
Photo Nos.: 4444	
Plot placement: <input checked="" type="checkbox"/> GRTS <input type="checkbox"/> Representative	
<input type="checkbox"/> Random <input type="checkbox"/> Stratified Random <input type="checkbox"/> Transect component	
<input type="checkbox"/> Systematic (grid) <input type="checkbox"/> Capture specific feature <input type="checkbox"/> Other	

Minimum required fields in Bold and Underlined

\* Definitions and values in CM PCAP FOM v. 1.0 and CVS Field Guide

OVER



NOTES: Include Layout (any unusual shape details), Location (directions and landscape content), Rationale (why here), and Veg Characterization (description of community, dominants, strata, BROWSE). Additional notes in space on back.

Layout: 2X5

Location: 75 meters southwest of 3-Way Intersection of Ox Lane + Oxbow Lane. Park along road.

Rationale: GRTS

Veg Characterization: The canopy is dominated by sugar maple, ~~large~~ large cordiforms. Red maple with various other individuals intermixed. Shrub layer dominated by Lindera, Sugar Maple and Beech. The herb layer was dominated by ramps and Podophyllum. Lots of Maple seedlings.

Fit= Conf=

Beech-Maple forest

Correct  
SDE 9-14-15

- ☐ Conspicuous inclusions
- ☐ Irregular/pattern mosaic

## HYDROLOGIC REGIME\*

**□ Fresh**

**Additional notes & diagrams:** (Representativeness of plot to the stand, successional status, maturity, etc.)

Additional notes & diagrams: (Representativeness of plot to the stand, successional status, maturity, etc.)

The stand is not quite even-aged but still has some successional species present. ~~Canopy~~ Canopy, shrub and herb layers are thick. The creek to the north is cutting deeply against the bank and is within 2 meters of plot. The western end (uphill) portion of plot borders a depression that is having a ~~seasonal~~ seasonal-submerging effect on a small portion of med 5. From ~~the~~ downed trees either from the eroding bank or whatever the shrub layer is somewhat more thick than surrounding forest.

\*\*L=low, ML=med low, M=med, MH=med high, H=high, VH=very high

**Former Land Use:**

Natural Resources Management FORM NR/2010-01b



Plot no.: 1012

Plot area (ha)



## Cleveland Metroparks


**Br = Browse Level. Use cover classes to describe amount of browse per species over entire plot**

Strata - Cov. entire plot

[illegible]

# EXAMPLES OF PERCENT OF AREA COVERED

The following graphic can be used for various data elements to convey "Amount" or "Quantity". NOTE: Within any given box, each quadrant contains the same total area covered, just different sized objects.



## BROWSE RATING NARRATIVE DESCRIPTION

**LOW OR NONE:** there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.

**MEDIUM LOW** values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

**MEDIUM:** browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.

**MEDIUM HIGH** values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.

**HIGH:** greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.

**VERY HIGH** values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.

**Typical Plot:**

Diagram showing a 50m x 20m plot divided into 10m x 10m modules. The plot is divided into 10 modules (1-10) and 5 corners (1-5). The plot is divided into 10 modules (1-10) and 5 corners (1-5). The plot is divided into 10 modules (1-10) and 5 corners (1-5).

**KEY:**

- Plot Origin
- Post
- Module Number
- Corner Number
- Tape
- Intensive Corner

**Nested Corners**

Diagram showing a 100m x 100m plot divided into 10m x 10m modules. The plot is divided into 10 modules (1-10) and 5 corners (1-5). The plot is divided into 10 modules (1-10) and 5 corners (1-5). The plot is divided into 10 modules (1-10) and 5 corners (1-5).

**Depth 1 = 100m<sup>2</sup>**

**Depth 2 = 10m<sup>2</sup>**

**Depth 3 = 1m<sup>2</sup>**

**Depth 4 = 0.1m<sup>2</sup>**


**Depth 5 = 0.01m<sup>2</sup>**






# EXAMPLES OF PERCENT OF AREA COVERED


This following graphic can be used to record data elements in corner "Amount" or "Quality". NOTE: Within any given box, each quadrant contains the same total area covered, just different sized objects.



cover class	% cover	midpoint
1	solitary or few	0.0001
2	0-1%	0.005
3	1-2%	0.015
4	2-5%	0.035
5	5-10%	0.075
6	10-25%	0.175
7	25-50%	0.375
8	50-75%	0.625
9	75-95%	0.850
10	95-100%	0.975



## Nested Corners



**BROWSE RATING NARRATIVE DESCRIPTION**

**LOW OR NONE:** there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.


**MEDIUM LOW** values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

**MEDIUM:** browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.

**MEDIUM HIGH** values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.

**HIGH:** greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.

**VERY HIGH** values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.










# EXAMPLES OF PERCENT OF AREA COVERED

The following graphic can be used for various data elements to convey "Amount" or "Quantity". NOTE: Within any given but, each quadrant contains the same total area covered, just different sized objects.



cover class	% cover	midpoint
1	solitary or few	0.0001
2	0-1%	0.005
3	1-2%	0.015
4	2-5%	0.035
5	5-10%	0.075
6	10-25%	0.175
7	25-50%	0.375
8	50-75%	0.625
9	75-95%	0.850
10	95-100%	0.975

## Nested Corners



**BROWSE RATING NARRATIVE DESCRIPTION**

**LOW OR NONE:** there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.

**MEDIUM LOW** values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

**MEDIUM:** browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.

**MEDIUM HIGH** values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.

**HIGH:** greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.

**VERY HIGH** values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.

## Page 1 of 1

Project name: OZ NL 2015 Plot no.: 1012

[illegible]



Project Label: PCAP Project name:                      Plot no.:

[illegible]

# CLEVELAND METROPARKS Plant Community Assessment Program Natural Woody Stem Data Sheet

Project Label: PCAP

Project Name: 02NC2015

Plot No.: 1012


Page: 1 of 4



Explain subsample (additional room on back):

mod #	species	c	voucher#	# stems 0-1.4m browed	% sub or super sample	# shrub clumps	size class (cm) woody stems > 1.4m												11 >40 (record each tree)
							1 0-1	2 1-2.5	3 2.5-5	4 5-10	5 10-15	6 15-20	7 20-25	8 25-30	9 30-35	10 35-40			
1	Acer saccharum																		73.0
1	Lindera benzoin																		
1	Fagus grandifolia																		
1	Standing dead																		
1	Acer rubrum																		
1	Ostrya virginiana																		
1	Carya cordiformis																		
1	Fraxinus pennsylvanica																		
1	Prunus serotina																		
1	Acer sp.																		
2	Lindera benzoin																		
2	Fagus grandifolia																		
2	Standing dead																		
2	Prunus serotina																		
2	Acer saccharum																		
2	Magnolia accuminata																		
2	Fraxinus pennsylvanica																		
2	Rubus alleghaniensis																		
3	Fagus grandifolia																		
3	Acer saccharum																		
3	Standing dead																		
3	Lindera benzoin																		
3	Fraxinus sp.																		
3	Carya cordiformis																		

### DBH Measurement Rules



### Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.


Record using the tally system from 1 to 10



1



2



3




4



5

### ASH CANOPY CONDITION

- 1. Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
- 2. Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
- 3. Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
- 4. >50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
- 5. Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.



A

B

C

D

E

### ASH CANOPY BREAKUP CONDITION (for dead trees):

(if an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)

- A:** All main branches contain fine twigs (newly dead).
- B:** Over 50% of main branches have fine twigs.
- C:** Less than 50% of main branches have fine twigs.
- D:** Stem still standing and tertiary main branches present.
- E:** Central stem still standing.



# CLEVELAND METROPARKS Plant Community Assessment Program Natural Woody Stem Data Sheet

Project Label: PCAP

Project Name: 62NC2015

Plot No.: 1012

Page: 2 of 4




Explain subsample (additional room on back):

mod #	species	c	voucher#	# stems 0-1.4m browed	% sub or super sample	# shrub clumps	size class (cm) woody stems >1.4m	1 0-1	2 1-2.5	3 2.5-5	4 5-10	5 10-15	6 15-20	7 20-25	8 25-30	9 30-35	10 35-40	11 >40 (record each tree)
3	Acer rubrum																	45.4
3	Fraxinus sp.																	
3	Liriodendron tulipifera																	
4	Fagus grandifolia																	
4	Lindera benzoin																	
4	Acer saccharum																	
4	Acer rubrum																	
4	Standing dead																	
4	Fraxinus pennsylvanica																	
5	Lindera benzoin																	
5	Standing dead																	
5	Carya bnfifolia																	
5	Fagls grandifolia																	
5	Acer saccharum																	
5	Magnolia acuminata																	
5	Carya sp. cordifolia																	
5	Tilia americana																	
5	Acer saccharum																	
5	Lindera benzoin																	
5	Carya cordifolia																	
5	Standing dead																	
5	Fraxinus pennsylvanica																	
5	Fagus grandifolia																	
5	Crataegus sp.																	

2010 had a lot of this being present in 2015. Most of the trees are large.

### DBH Measurement Rules



### Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.

Record using the tally system from 1 to 10



1



2



3



4



5

### ASH CANOPY CONDITION

- 1. Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
- 2. Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
- 3. Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
- 4. >50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
- 5. Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.



A

B

C

D

E

### ASH CANOPY BREAKUP CONDITION (for dead trees):

(If an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)

- A:** All main branches contain fine twigs (newly dead).
- B:** Over 50% of main branches have fine twigs.
- C:** Less than 50% of main branches have fine twigs.
- D:** Stem still standing and tertiary main branches present.
- E:** Central stem still standing.



# CLEVELAND METROPARKS Plant Community Assessment Program Natural Woody Stem Data Sheet

Project Label: PCAP

Project Name: OZNC2015

Plot No.: 1012

Page: 3

of

4




Explain subsample (additional room on back):

mod #	species	c	voucher#	# stems 0-1.4m browsed	% sub or super sample	# shrub clumps	size class (cm) woody stems > 1.4m	1 0-1	2 1-2.5	3 2.5-5	4 5-10	5 10-15	6 15-20	7 20-25	8 25-30	9 30-35	10 35-40	11 >40 (record each tree)
6	<del>Fraxinus sp.</del>																	
6	<del>Acer sp.</del>																	
7	Acer saccharum																	83.7
7	Fagus grandifolia																	
7	Standing dead																	
7	Lindera benzoin																	
7	Crataegus sp.																	
7	Prunus serotina																	
7	Acer sp.																	
8	Acer saccharum																	73.4
8	Standing dead																	
8	Lindera benzoin																	
8	Fagus grandifolia																	
8	Acer sp.																	
9	Lindera benzoin																	
9	Standing dead																	
9	Acer saccharum																	62.6
9	Sabua cordiformis																	
9	Fagus grandifolia																	
9	Parthenocissus quinquefolia																	
9	Fraxinus sp.																	
10	Lindera benzoin																	
10	Acer saccharum																	
10	Carya cordiformis																	



### DBH Measurement Rules



### Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.

Record using the tally system from 1 to 10



1



2



3



4



5

### ASH CANOPY CONDITION

- 1. Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
- 2. Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
- 3. Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
- 4. >50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
- 5. Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.



A

B

C

D

E

### ASH CANOPY BREAKUP CONDITION (for dead trees):

(If an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)


- A:** All main branches contain fine twigs (newly dead).
- B:** Over 50% of main branches have fine twigs.
- C:** Less than 50% of main branches have fine twigs.
- D:** Stem still standing and tertiary main branches present.
- E:** Central stem still standing.

**Cleveland Metroparks**

**Cleveland Metroparks**

Natural Resources Management FORM NR/2010-03a

### DBH Measurement Rules




### Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.

Record using the tally system from 1 to 10



1



2



3



4



5

### ASH CANOPY CONDITION

1. Healthy, full canopy: A healthy ash canopy is normally thinner than many other trees such as maple.
2. Thinning canopy: There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
3. Dieback: Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
4. >50% Dieback: The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
5. Dead canopy: No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.



A

B

C

D

E

### ASH CANOPY BREAKUP CONDITION (for dead trees):

(If an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)

- A: All main branches contain fine twigs (newly dead).
- B: Over 50% of main branches have fine twigs.
- C: Less than 50% of main branches have fine twigs.
- D: Stem still standing and tertiary main branches present.
- E: Central stem still standing.



Project Label: PCAP

Project Name:

02NC2015

Plot No.:

1012

Date:


14 July 2015

Tree ID	Species	Dead	Voucher #	DBH (cm)	Ht (m)	Ash condition	Dead condition	# Exit holes	Epilimna present	Woodpecker holes
1	No ash present									
2										
3										
4										
5										
6										
7										
8										
9										
10										
11										
12										
13										
14										
15										
16										
17										
18										
19										
20										
21										
22										
23										
24										
25										

\* If Ash Condition scores 5 (dead) provide breakup score (A-E)  
Count EAB exit holes 1.25m x 21.5m  
Woodpecker and epicormic marked present (1) or absent (0)



\*\*\* Change intensive module numbers when necessary



Map all ash trees ≥ 10cm in each module using Tree ID number

CLEVELAND METROPARKS Plant Community Assessment Program: Invasive Species Survey



Tier 1: Early detection/ Rapid response		Presence				GPS	Presence X: yes
		NE	SE	SW	NW		
Microstegium vimineum	Japanese stiltgrass						
Ranunculus ficaria	Lesser Celandine						
Cynanchum louiseae (vine)	Black Swallow-wort						
Butomus umbellatus (wetland)	Flowering Rush						
Heracleum mantegazzianum	Giant Hogweed						
Tier 2: Assess as Needed		# of Plants				comments	# of Plants
		NE	SE	SW	NW		
Acer platanoides	Norway Maple						1: 1-10
Ailanthus altissima	Tree of Heaven						2: 11-50.
Lonicera japonica (vine)	Japanese Honeysuckle						3: 51-100
Lythrum salicaria (wetland)	Purple Loosestrife						4: 101-1,000
Aegopodium podagraria (G-cover)	Bishop's Goutweed						5: >1,000
Celastrus orbiculatus (vine)	Asian Bittersweet						
Torilis sp.	Hedgeparsley						
Conium maculatum	Poison Hemlock						
Rhamnus cathartica	Common Buckthorn (shrub)						
Berberis thunbergii	Japanese Barberry (shrub)						
Alnus glutinosa	European Alder						
Dipsacus laciniatus	Cut-leaf Teasel						
Elaeagnus umbellata	Autumn Olive (shrub)						
Lonicera maackii	Amur Honeysuckle (shrub)						
Euonymus fortunei	Wintercreeper						
Tier 3: Presence is of Interest		# of Plants				comments	# of Plants
		NE	SE	SW	NW		
Convallaria majalis (G-cover)	Lily of the Valley						1: 1-10
Coronilla varia (G-cover)	Crown Vetch						2: 11-50.
Eleutherococcus pentaphyllus	Five-leaf Aralia (shrub)						3: 51-100
Pachysandra terminalis (G-cover)	Japanese Pachysandra						4: 101-1,000
Philadelphus coronarius	Mock Orange (shrub)						5: >1,000
Pulmonaria officinalis (G-cover)	Lungwort						
Rubus phoenicolasius	Wineberry						
Iris pseudacorus (wetland)	Yellow Flag Iris						
Ornithogalum umbellatum	Star of Bethlehem						
Viburnum opulus var. opulus	European Cranberry (shrub)						
Viburnum plicatum	Doublefile Viburnum (shrub)						
Tier 4: Widespread and abundant		Presence				comments	# of Plants
		NE	SE	SW	NW		
Alliaria petiolata	Garlic Mustard						1: 1-10
Ligustrum vulgare	Common Privet (shrub)						2: 11-50.
L. morrowii, L. tatarica	Bush Honeysuckles (shrub)						3: 51-100
Phalaris arundinacea	Reed Canarygrass						4: 101-1,000
Phragmites australis (wetland)	Phragmites						5: >1,000
Polygonum cuspidatum	Japanese Knotweed						
Frangula alnus	Glossy Buckthorn (shrub)						
Rosa multiflora	Multiflora Rose (shrub)						
Typha angustifolia, T. x. glauca	Cattails (wetland)						
Cirsium arvense	Canada thistle						
Dipsacus fullonum	Common Teasel						
Hesperis matronalis	Dame's Rocket						
Vinca minor (G-cover)	Periwinkle						

Note: For Ground-cover plants record "stem #" but in comment field describe # of colonies and patch size (S,M, L)

# CLEVELAND METROPARKS Plant Community Assessment Program Forest Pest and Pathogens Data Sheet

Project Label: \_\_\_\_\_

PCAP \_\_\_\_\_

Project Name: 02NC2015

Plot No.: 1012

Page: 1 of 1



mod #	species	voucher#	# shrub clumps	size class (cm) woody stems > 1m										
				1 0-1	2 1-2.5	3 2.5-5	4 5-10	5 10-15	6 15-20	7 20-25	8 25-30	9 30-35	10 35-40	11 >40 (record each tree)
1	<i>Fagus grandifolia</i>													
2	<i>Fagus grandifolia</i>													
3	<i>Fagus grandifolia</i>													
4														
5	<i>Fagus grandifolia</i>													
6	<i>Fagus grandifolia</i>													
7	<i>Fagus grandifolia</i>													
8	<i>Fagus grandifolia</i>													
9	<i>Fagus grandifolia</i>													
10	<i>Fagus grandifolia</i>													

\* IF EVIDENCE OF PEST OR PATHOGEN RECORD TOTAL SPECIES POPULATION IN THE PLOT EVEN THE NOT INFECTED

Strata	# of stems infected	Severity (H, M, or L)
Tree (size class 3 or above)	100	L
Shrub (size class 2 or below including shrub clumps)	100	M

\* Write None Present if no evidence:

Beech (Fungus)	Asian Longhorned Beetle
Hemlock (HWA)	Other Pest or Pathogen
Walnut (Thousand Canker)	

## Severity

High = more than 50% of leaf/needle cover exhibiting symptoms

Medium = Less than 50% of leaf/needle cover exhibiting symptoms

Low = Only a few leaves or branches are exhibiting symptoms





Original Datasheet

Page: \_\_\_\_\_ of \_\_\_\_\_

**Cleveland Metroparks**  
of

① Herb: 1-1007.  
 Mod = 1007, in fact = mod level  
 ② low security, 1st / mod, 2nd / 1007  
 3rd / 1007

\* Write None Present if no evidence:

Density of Infestation:  
High, med, or low

③ ~~Shrub~~-L, L, M, M, M, L, L  
herb-L  
shrub-M, M, L

④ herb-L  
shrub-M, M, L

⑤ shrub-L  
herb-L, L, L, B

⑥ Shrub-L, H

⑦ Sh-L, M  
herb-M, M

⑧ sh-M, L, M, L, M, L

②  $Sh-M, M, L, L, M, L$

76/sh-med, m, m





STANDING BIOMASS (required for emergent wetland) collected in 0.1m clip plot (33x32 cm) from corner 1 and 3 in each intensive module. Required for VIBI-E score calculation. C7=check when collected

Module #	C7	Corner	Corner

### CLASSIFICATION

G11 - exclusive of Fit and Confidence

Hydroscorable class (WETLANDS ONLY)

<input type="checkbox"/> DEPRESSION	Fit=	Conf=
<input type="checkbox"/> INFUNDICENT <input type="checkbox"/> Beaver <input type="checkbox"/> Human	Fit=	Conf=
<input type="checkbox"/> RIVERINE <input type="checkbox"/> Headwater <input type="checkbox"/> Mainstem <input type="checkbox"/> Channel	Fit=	Conf=
<input type="checkbox"/> SLOPE (ground water hydrology or on a physical slope)	Fit=	Conf=
<input type="checkbox"/> FRINGING <input type="checkbox"/> Reservoir <input type="checkbox"/> Natural Lake	Fit=	Conf=
<input type="checkbox"/> COASTAL (specific subclasses)	Fit=	Conf=
<input type="checkbox"/> BOG (strongly, moderately, weakly ombrotrophic)	Fit=	Conf=

Ohio EPA VIBI Plant Community Class (WETLANDS ONLY)

<input type="checkbox"/> FOREST <input type="checkbox"/> swamp forest <input type="checkbox"/> bog forest <input type="checkbox"/> forest swamp	Fit=	Conf=
<input type="checkbox"/> EMERGENT <input type="checkbox"/> marsh <input type="checkbox"/> wet meadow <input type="checkbox"/> open bog	Fit=	Conf=
<input type="checkbox"/> SHRUB <input type="checkbox"/> shrub swamp <input type="checkbox"/> tall sh. bog <input type="checkbox"/> tall sh. fen	Fit=	Conf=

### MICROTOPOGRAPHIC FEATURE COUNTS - Intensive modules only

Results for microhabitat features. Select one or select two and average the score. NOTE: If mod falls on a slope automatically gets ranked based on steepness (1-3) to begin + any features present  
 Slope 1 = slight elevational grade across module (flat) Slope 2 = both on slope -20° Slope 3 = maximum steepness that can be safely sampled -45°

- 0 feature is absent or functionally absent from the wetland
- 3 feature is present in the wetland in very small amounts or if more common, of low quality
- 7 feature is present in moderate amounts, but not of highest quality, or in small amounts of highest quality
- 10 feature is present in moderate or greater amounts and of highest quality

C.W.D. - count for pieces with minimum 1m length									
mod#	corner	no. of tussocks	no. of hummocks uplands (TTP-Lps)	no. macro. depressions	C.W.D. (2-12 cm)	C.W.D. (13-10cm)	C.W.D. >40 cm	microhab. interspect.	microhab.
		depth 3 1x1m (count)	depth 2 3 (6x3, 16m) (count)	depth 1 10x10m (count)	depth 1 10x10m (count)	depth 1 10x10m (count)	depth 1 10x10m (count)	depth 1 10x10m (rank)	SLOPE 10x10m (rank)
2	4	0	0	1	4 = 14	0 = 0	0 = 0	3	1
3	4	0	0	2	13 = 13	1 = 1	8 = 0	3	1
4	4	0	0	4	10 = 10	1 = 1	1 = 1	4	1
9	4	0	0	2	15 = 16	0 = 0	0 = 0	4	1

NOTE: tussock and hummocks are counted in BOTH nested quadrat corners but counts are aggregated.

### MGBI INDICES (degrees) + for up - for down

FILLED OUT USING OUR PROGRAM - DO NOT FILL OUT IN FIELD

At aspect	N	S	E	W
45 degrees				
90 degrees				
135 degrees				
180 degrees				
225 degrees				
270 degrees				
315 degrees				

\* Landform Index (position within landscape)

\* Terrain Slope Index (slope microtopographic shape)

CROWN COVER (DENSITOMETER): Male 4  
 readings per module facing N, S, E, W. Place dot count in corresponding space. (1 dot per grid square)

Module	N	S	E	W
2	4	6	0	0
3	4	2	1	1
4	4	0	0	0
9	4	3	3	2

# COVER BY STRATA

STRATUM	GENERAL FORM
Tree (generally >5 m)	Tree (overstory), very tall shrubs*, liana, epiphyte)
Shrub (generally 0.5 to 5 m)	Tree (sapling), shrub, liana, epiphyte)
Herb (Field)	Herb, dwarf-shrub**, tree (seedling***)
Floating	Floating
Aquatic (submerged)	Submerged

\*Very tall shrubs are sometimes included in the tree stratum

\*\*Can also include seedlings of shrubs, i.e. all shrubs <0.5m

\*\*\*Tree seedlings are often defined as up to 1.4 m height or as <2.5 cm DBH in which case they would span the herb and shrub layers.





FIGURE 3-20.—Generalized section of Upper Devonian, Mississippian, and Lower Pennsylvanian formations in northeastern Ohio. Asterisks indicate units that are fossiliferous. This composite section represents about 400 meters of rock exposed across the area. The section is not to scale, but the thicknesses indicated are proportional. The term "Waverly" is used in the older literature to refer to Mississippian rocks in Ohio. Some geologists use the European term "Carboniferous," which encompasses the Mississippian and Pennsylvanian Periods of the U.S. Many units have been named within the Cuyahoga Formation, but most units are local and cannot be traced over great distances. The Black Hand Member is a spectacular massive sandstone that is fairly widespread but discontinuous. See Hyde (1953), Hoover (1960), and Collins (1979) for more information on Mississippian rocks in Ohio. See figure 3-18 for explanation of rock types.

**SOIL PIT DESCRIPTION:** Excavate 20 cm plug with shovel. Describe using Munsell chart, visual exam, texture, and odor.

**SOIL SAMPLES** Standard procedure: collect a soil sample of the top 10 cm of soil from center of each intensive module and composite the sample

Soil pit module #      (one per entire plot)

5 cm	matrix color
	mottle color
	%mottle
	oxid roots
	texture*
	redox features**
	hydr. cond.***
20 cm	matrix color
	mottle color
	%mottle
	oxid roots
	texture*
	redox features**
	hydr. cond.***

\* refer to texture classes on reverse side  
 \*\* e.g. hydrogen sulfide odor, glistening, etc.  
 \*\*\* Circle one:

Unindurated S=saturated M=moist D=dry

Notes: include evidence of earthworms (worms, castings, middens)

2 worms present (2) castings, middens  
 3 worms, casting found  
 8 worms and casting present  
 9 castings, no worms noted

Soil Collection Method	Hartman (A, B, C)
2,3,8,9 composited	A
Method Soil Survey Information	
Soil Series/Type	
Soil Series Source	Ohio Soil Survey
Landform type	
Depth to root layer	
Parent Material	
Drainage*	
<input type="checkbox"/> Excessively dr. <input type="checkbox"/> Somewhat excessively <input type="checkbox"/> Well drained <input type="checkbox"/> Moderately well dr. <input type="checkbox"/> Somewhat poorly dr. <input type="checkbox"/> Very poorly dr. <input type="checkbox"/> Impermeable surface	

**SOIL DEPTH MEASUREMENT:** Measure to the nearest 0.1 cm in center of intensive modules. If >30.5 cm, record as >30

	1 liter+ organic depth (cm)	2 liter depth (cm)	water depth (cm)	depth sat soil (cm)
mod#				
2	1.3	1.3	—	—
3	1.1	1.1	—	—
8	0.9	0.9	—	—
9	1.3	1.3	—	—

EARTH SURFACE & GROUND COVER			
Underlying Earth Surface*	Ground Cover		
Grass - 100%	Percent	Height ≤ 100%	Percent
Histocool	0	Coarse Woody Debris**	7
Mineral Soil	100	Fine Woody Debris***	4
Gravel-Cobble*	0	Litter	75
Boulder**	0	Duff (Ferm + Humus)	0
Bedrock	0	Bryophyte-Lichen	1
* Gravel-Cobble = 1/16-10"	Water		0
** Boulder = > 10 in	Bare Soil		1
*** > 5 cm in diameter	Dead Trail		1
**** < 5 cm in diameter	Other		0

TRAIL INFORMATION:	
Record type and cover for each	
Type	%Cover
<input type="checkbox"/> All Purpose	
<input type="checkbox"/> Bridge	
<input type="checkbox"/> Hiking sanctioned	
<input type="checkbox"/> Boulding unsanctioned	
<input type="checkbox"/> Gravel	
<input checked="" type="checkbox"/> Other	1

**COVER BY STRATA**  
 estimate using midpoints of 5, ex: 3, 8, 13 %

Strata	Midnet Range (cm)	Total Cover (%)
Tree	5 - 4	83
Shrub	0.5 - 5	68
Herb	0 - 0.5	68
(Floating)*	—	—
(Aquatic)*	—	—


\* rooted and floating or slightly emerged  
 \*\* submerged most plant mass below surface  
 SEE BACK OF PAGE FOR TYPICAL STRATA DESCRIPTIONS. STRATA CAN VARY BY COVER TYPE.

STAND SIZE	
<input type="checkbox"/> > 600 x plot size	
<input checked="" type="checkbox"/> > 100 x plot size	
<input type="checkbox"/> 10-100 x plot size	
<input type="checkbox"/> 3-10 x plot size	
<input type="checkbox"/> 1-3 x plot size	
<input type="checkbox"/> < plot size	




# PERCENT MOTTLES (USE CLASS CODES):

Class	Code	Criteria: % of Surface Area Covered
Conv.	NASIS	
Few	f	< 2
Common	c	2 to < 20
Many	m	≥ 20



2%





20%

**SOIL TEXTURE:** Record the code for the soil texture of the 5 cm and 20 cm layers. To estimate texture, collect a soil sample from the appropriate layer and moisten it with water to the consistency of modeling clay/wet newspaper; the sample should be wet enough that all of the particles are saturated but excess water does not freely flow from the sample when squeezed. Attempt to roll the sample into a ball. If the soil will not stay in a ball and has a grainy texture, the texture is either sandy or coarse sandy. If the soil does form a ball, squeeze the sample between your fingers and attempt to form a self-supporting ribbon. Samples which form both a ball and a ribbon should be coded as clayey; samples which form a ball but not a ribbon should be coded as loamy.

- 0= Organic
- 1= Loamy
- 2= Clayey
- 3= Sandy
- 4= Coarse Sand
- 9= Not measured - make plot note


**Geomorphic Component - Three-dimensional descriptors of parts of landforms or microfeatures that are best applied to areas. Unique descriptors are available for Hills, Terraces, Mountains, and Flat Plains; e.g., (for Hills) nose slope or NS.**

Hills	Code	NASIS
PDP		
interfluvial	IF	IF
head slope	HS	HS
nose slope	NS	NS
side slope	SS	SS
base slope	—	BS



**Hillslope - Profile Position (Hillslope Position in PDP) - Two-dimensional descriptors of parts of line segments (i.e., slope position) along a transect that runs up and down the slope; e.g., backslope or BS. This is best applied to transects or points, not areas.**

Position	Code
summit	SU
shoulder	SH
backslope	BS
footslope	FS
toeslope	TS



**HYDROLOGIC REGIME** Modified from Grossman et al 1998. (Frequency and duration of flooding.)

**UPLAND:** Not a wetland. Very rarely flooded.

**INTERMITTENTLY/SEASONALLY SATURATED:** Dry at least once per year. Surface water is seldom present, but substrate is saturated to surface for extended periods during the growing season.

**PERMANENTLY/SEMPERMANENTLY SATURATED:** Dry less than once per year. Surface water is seldom present, but substrate is saturated to surface for extended periods during the growing season. Equivalent to Cowardin's Saturated modifier.

**OCCASIONALLY FLOODED:** Surface water can be present for brief periods during growing season, but not in most years. Often characterizes flood-plain upper terraces.

**TEMPORARILY FLOODED:** Surface water present for brief periods during growing season, but water table usually lies well below soil surface. Often characterizes flood-plain levees and lower terraces. Equivalent to Cowardin's Temporary modifier.

**INTERMITTENTLY FLOODED:** Substrate is usually exposed, but surface water can be present for variable periods without detectable seasonal periodicity. Inundation is not predictable to a given season and is dependent upon highly localized rain storms. This modifier was developed for use in the arid West for water regimes of Playa lakes, intermittent streams, and dry washes but can be used in other parts of the U.S. where appropriate. This modifier can be applied to both wetland and non-wetland situations. Equivalent to Cowardin's Intermittently Flooded modifier.

**SEMPERMANENTLY FLOODED (exposed <1/year):** Surface water persists throughout the growing season in most years. Land surface is normally saturated when water level drops below soil surface. Includes Cowardin's Intermittently Exposed and Semipermanently Flooded modifiers.

**PERMANENTLY FLOODED:** Water covers the land surface at all times of the year in all years. Equivalent to Cowardin's "permanently flooded".

**UNKNOWN:** The hydrologic regime cannot be determined from the available information.