

CLEVELAND METROPARKS Plant Community Assessment Program: Quality Control Form



Project Label: _____

PCAP

Plot No: 1054Date Sampled: 07/20/15Lead: LANCE

Comment required if item answer is NO

Parking/Access outside of Park Boundaries:		<input checked="" type="radio"/> Y	<input checked="" type="radio"/> N	If yes, write details in Comments section below
Field journals completed		<input checked="" type="radio"/> Y	<input type="radio"/> N	
Site sketch made on 1:3000 map?		<input checked="" type="radio"/> Y	<input type="radio"/> N	
Check cover page	X-axis Bearing of plot recorded	<input checked="" type="radio"/> Y	<input type="radio"/> N	
	GPS coords. Recorded	<input checked="" type="radio"/> Y	<input type="radio"/> N	
	North direction recorded	<input checked="" type="radio"/> Y	<input type="radio"/> N	
	Photographs taken?	<input checked="" type="radio"/> Y	<input type="radio"/> N	
	Relocated Pins Mapped	<input checked="" type="radio"/> Y	<input type="radio"/> N	
Plot No., Date agreement on all pages?		<input checked="" type="radio"/> Y	<input type="radio"/> N	
Header data completed all pages?		<input checked="" type="radio"/> Y	<input type="radio"/> N	
Cover classes recorded in all Intensive modules		<input checked="" type="radio"/> Y	<input type="radio"/> N	
Browse Level By Species		<input checked="" type="radio"/> Y	<input type="radio"/> N	
Woody stem quality control check		<input checked="" type="radio"/> Y	<input type="radio"/> N	Check every line and cross check with the Tree Cover Sheet
Invasive plant quality control check		<input checked="" type="radio"/> Y	<input type="radio"/> N	N/A
Ash trees mapped		<input checked="" type="radio"/> X	<input type="radio"/> N	N/A
Completed Forest Pest/Pathogen Datasheet		<input checked="" type="radio"/> Y	<input type="radio"/> N	
Cover by Strata? (confirm cover type)		<input checked="" type="radio"/> Y	<input type="radio"/> N	
Soil samples collected with matching plot #.		<input checked="" type="radio"/> Y	<input type="radio"/> N	
Cross check 2010 information		<input checked="" type="radio"/> Y	<input type="radio"/> N	Highlight any changes from 2010 information
Vouchers labeled on datasheet with initials and number		<input checked="" type="radio"/> Y	<input type="radio"/> N	
Vouchers labeled on collection bag		<input checked="" type="radio"/> Y	<input type="radio"/> N	
Pink flags removed		<input checked="" type="radio"/> Y	<input type="radio"/> N	
Data sheet QA before leaving site?		<input checked="" type="radio"/> Y	<input type="radio"/> N	
Common equipment returned to tub.		<input checked="" type="radio"/> Y	<input type="radio"/> N	
Data sheets scanned?				Enter date to left
Final data sheets scanned?				Enter date to left
Buffer Widths measured?		<input checked="" type="radio"/> Y	<input type="radio"/> N	
Web Soil Survey		<input checked="" type="radio"/> Y	<input type="radio"/> N	
Voucher Location	Refrigerator	<input checked="" type="radio"/> Y	<input type="radio"/> N	
(# vouchers collected)	Press (#)			Enter number to left
	Drier	<input checked="" type="radio"/> Y	<input type="radio"/> N	
	Identified	<input checked="" type="radio"/> Y	<input type="radio"/> N	
	Mounted	<input checked="" type="radio"/> Y	<input type="radio"/> N	
	Thrown away	<input checked="" type="radio"/> Y	<input type="radio"/> N	

GRTS point verification: Is plot sampleable?

<input checked="" type="checkbox"/> Yes	Original GRTS point is sampleable
<input type="checkbox"/> No	Original GRTS point lands in a non-sampleable area (fill in category below)
	<input type="checkbox"/> Point falls in a water (i.e. river, lake)
	<input type="checkbox"/> Managed mowed area (i.e. golf course, picnic area, right-of-way)
	<input type="checkbox"/> Paved area (i.e. parkinglot, road)
	<input type="checkbox"/> Unsafe to sample (i.e. steep slope)
	<input type="checkbox"/> Other

Additional Comments:

--

b

1

GENERAL INFORMATION		LOCATION																													
Project Label: PCAP	State: OH County: Cuyahoga	Local Place Names:																													
Project Name: 02RR2015	Quadrangle: Lakewood	Landowner: CMP																													
Plot Name: Mosquito Heaven																															
Plot No.: 1054																															
<input type="checkbox"/> Level 4 (no nested corners sampled)																															
<input checked="" type="checkbox"/> Level 5 (nested corners sampled)																															
Date (mm/dd/yyyy): 07/22/2015																															
End date (if > 1 day): 1																															
Party: H. Lance	Role: Plot leader																														
M. Getgey	Bot. Asst.																														
E. Knauts	Crew																														
M. Busam	Crew																														
Plot NOT SAMPLED: <input type="checkbox"/> Other <input type="checkbox"/> Perm. water <input type="checkbox"/> Paved <input type="checkbox"/> Slope <input type="checkbox"/> Safety																															
SAMPLING QUALITY* Effort Level: <input checked="" type="checkbox"/> Every thorough <input type="checkbox"/> Accurate <input type="checkbox"/> Hurried subjective evaluation of how much effort put into sampling. Hurried plots may still provide good data																															
TAXONOMIC ACCURACY <table border="1"> <tr> <td>high</td> <td>modera.</td> <td>low</td> <td>not simpl</td> </tr> <tr> <td><input checked="" type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> </tr> <tr> <td>vascul.</td> <td></td> <td></td> <td>n/a</td> </tr> <tr> <td><input checked="" type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> </tr> <tr> <td>dryo</td> <td></td> <td></td> <td></td> </tr> <tr> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> </tr> <tr> <td>lichen</td> <td></td> <td></td> <td><input checked="" type="checkbox"/></td> </tr> </table>		high	modera.	low	not simpl	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	vascul.			n/a	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	dryo				<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	lichen			<input checked="" type="checkbox"/>		
high	modera.	low	not simpl																												
<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>																												
vascul.			n/a																												
<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>																												
dryo																															
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>																												
lichen			<input checked="" type="checkbox"/>																												
TAXONOMIC STANDARD Authority: G&C Pub Date: 1998																															

Reason: If data not public why? Source of coordinates: <input type="checkbox"/> MAP <input checked="" type="checkbox"/> GPS Coordinate system: <input checked="" type="checkbox"/> Lat/Long <input type="checkbox"/> UTM <input type="checkbox"/> StatePlane <input type="checkbox"/> Other (specify) Datum: <input checked="" type="checkbox"/> NAD83/WGS84 <input type="checkbox"/> NAD27	GPS location in plot x=0 to 5, y=-1.0, +1): x = 0 y = 0 (base of plot x=0, y=0) Latitude: 41.41938 Longitude: 81.85876 Coord. Accuracy: <input checked="" type="checkbox"/> m <input type="checkbox"/> ft GPS File Name: 1054A Plot size for cover data: 0.1 (hectares) X-axis Bearing of plot: 262 ° Depth: (1-5): 4 Intensive modules: 2, 3, 8, 9 (EDIT IF MODIFIED) Camera No.: 3 Photo Nos.: 0141 Plot placement: <input checked="" type="checkbox"/> GRTS <input type="checkbox"/> Representative <input type="checkbox"/> Random <input type="checkbox"/> Stratified Random <input type="checkbox"/> Transect component <input type="checkbox"/> Systematic (grid) <input type="checkbox"/> Capture specific feature <input type="checkbox"/> Other
---	---

Layout → 2x5
 Location → Park just north of the Brookpark Rd. overpass. Walk under the overpass for approx. 100 m. Plot is ~40 m from overpass.
 Rationale → GRTS; PCAP re-sample
 Veg. Characteristics → ^{Early} High quality floodplain community. Black snakeroot and wingstem completely dominate the herbaceous layer.
 Canopy is primarily black walnut, with →

Diagram Key: Plot origin (0,0) point, GPS location point, photo taken, with direction, location of permanent posts

NOTES: Include Layout (any unusual shape details), Location (directions and landscape content), Rationale (why here), and Veg Characterization (description of community, dominants, strata, BROWSE). Additional notes in space on back.

OVER

*Definitions and values in CM PCAP FOM v. 1.0 and CVS Field Guide

CLEVELAND METROPARKS Plant Community Assessment Program - Background Data Sheet

Project Label: PCAP

Project Name: 02882015

Plot No.: 1054

Cleveland Metroparks

Page 2 of 2

MODIFIED NATURE RESERVE CLASS*

CODE (on separate form): L-01

Fit= Conf=

COMMUNITY NAME:

Mesic Floodplain Forest

HOMOGENEITY

☒ Homogeneous
 ☐ Compositional trend across the plot

☐ Conspicuous inclusions
 ☐ Irregular/pattern mosaic

DISTURBANCES

type*	severity**	yrs ago	% of plot	description
Human	L	0	100%	trash
Natural				
Fire				
Cut				
Animal	M	0	100%	browse
Other	L	0	100%	overpass influence (noise)

**L=low, M=med low, M=med, MH=med high, H=high, VH=very high

Current Land Use: PARK

Former Land Use: UNKNOWN

HYDROLOGIC REGIME*

☐ Upland (seldom flooded)
 ☒ Intermittently/seasonally saturated (seldom flooded)
 ☐ Permanently/Semipermanent saturated (dry < 1/yr, seldom flooded)
 ☐ Occasionally flooded (< 1/yr)
 ☐ Temporarity flooded

☐ Intermittently flooded
 ☐ Semipermanently flooded
 ☐ Permanently flooded
 ☐ Tidal/Seiche flooded daily
 ☐ Tidal/Seiche flooded monthly
 ☐ Tidal/Seiche flooded irregular (e.g. wind, storms)
 ☐ Unknown

SALINITY*

☐ Saltwater
 ☐ Brackish
 ☒ Fresh
 ☐ Upland (n/a)

(by default unless plot is a wetland)

Additional notes & diagrams: (Representativeness of plot to the stand, successional status, maturity, etc.)

box elder and a few buckeyes as well.
Shrub layer dominated by box elder.

Page 1 of 2

Plot area (ha)




Cleveland Metroparks

Br = Browse Level. Use cover classes to describe amount of browse per species over entire plot

[illegible]




EXAMPLES OF PERCENT OF AREA COVERED

The following graphic can be used to convert data elements to canopy "Amount" or "Quantity". NOTE: Within any given box, each quadrant contains the same total area covered, just different sized objects.



cover class	% cover	midpoint
1	solitary or few	0.0001
2	0-1%	0.005
3	1-2%	0.015
4	2-5%	0.035
5	5-10%	0.075
6	10-25%	0.175
7	25-50%	0.375
8	50-75%	0.625
9	75-95%	0.850
10	95-100%	0.975

Nested Corners



BROWSE RATING NARRATIVE DESCRIPTION

LOW OR NONE: there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.

MEDIUM LOW values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

MEDIUM: browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.

MEDIUM HIGH values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.

HIGH: greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.

VERY HIGH values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.

Project Label: PCAP

Project name: 028205

Plot no.: 1054

Total modules: 10

Intensive modules: 4

Plot configuration: 2x5

Plot area (ha): .1




Cleveland Metroparks

Br = Browse Level. Use cover classes to describe amount of browse per species over entire plot

S	H	(F)	(A)	Br	Species	C	Voucher #	Estimate for each intensive module:				Estimate for each intensive module:				Estimate for each intensive module:				Estimate for each intensive module:				R	mod
								%open water	depth	cover	mod	%unvegetated open water	depth	cover	mod	%unveg. ground (bare soil)	depth	cover	mod	%unveg. filter (bare filter)	depth	cover	mod		
4	2				Pinus serotina				1	4	2		1	4	2		1	4	2		1	4	2		
5	2				Vitis riparia				1	4	2		1	4	2		1	4	2		1	4	2		
7	2				Ranunculus sp.				1	4	2		1	4	2		1	4	2		1	4	2		
7	2				Poa sp. ^{Elymus} maritima ^{maritima}				1	4	2		1	4	2		1	4	2		1	4	2		
4	2				Lindera benzoin				1	4	2		1	4	2		1	4	2		1	4	2		
1					Trillium sp.				1	4	2		1	4	2		1	4	2		1	4	2		
1					Carya sp.				1	4	2		1	4	2		1	4	2		1	4	2		
2					Allium tricoccum				1	4	2		1	4	2		1	4	2		1	4	2		
2					Fraxinus pennsylvanica				1	4	2		1	4	2		1	4	2		1	4	2		
1					Sedum ternatum				1	4	2		1	4	2		1	4	2		1	4	2		
1					Geum canadense				1	4	2		1	4	2		1	4	2		1	4	2		
2					Rhus virginiana				1	4	2		1	4	2		1	4	2		1	4	2		
3					Fraxinus americana				1	4	2		1	4	2		1	4	2		1	4	2		
2					Geranium maculatum				1	4	2		1	4	2		1	4	2		1	4	2		
1					Thalictrum pubescens				1	4	2		1	4	2		1	4	2		1	4	2		
4	1				Crataegus sp.				1	4	2		1	4	2		1	4	2		1	4	2		
1					Aster sp.				1	4	2		1	4	2		1	4	2		1	4	2		
1					Canna cordiformis				1	4	2		1	4	2		1	4	2		1	4	2		
1					Osmorhiza longistylis				1	4	2		1	4	2		1	4	2		1	4	2		

EXAMPLES OF PERCENT OF AREA COVERED

The following graphic can be used for various data elements to convey "Amount" or "Quantity". NOTE: Within any given box, each quadrant contains the same field area covered, just different sized objects.



BROWSE RATING NARRATIVE DESCRIPTION

LOW OR NONE: there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.



MEDIUM LOW values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

MEDIUM: browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.

MEDIUM HIGH values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.

HIGH: greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.

VERY HIGH values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.



Page 1 of 1

SQL 9-27-15

% COVER												Presence of tree species (X)		mod		mod		mod		mod		R	
Strata - Cov. entire plot												c		Voucher #								R	
T		Br		Species																			

CLEVELAND METROPARKS Plant Community Assessment Program Natural Woody Stem Data Sheet

Project Label: PCAP

Project Name: 02R2015

Plot No.: 1054

Page: 1

of

3




Explain subsample (additional room on back):

mod #	species	c	voucher#	# stems 0-1.4m browed	% sub or super sample	# shrub clumps	size class (cm) woody stems > 1.4m													
							1 0-1	2 1-2.5	3 2.5-5	4 5-10	5 10-15	6 15-20	7 20-25	8 25-30	9 30-35	10 35-40	11 >40 (record each tree)			
1	Asclepias glabra			.																
1	standing dead																			
1	Vitis sp. riparia																			
1	Fraxinus americana																			
1	Rhus serotina																			
1	Acer nigrum																			
1	Fraxinus sp. seedling			.																
2	Asclepias glabra																			
2	standing dead																			
2	Acer negundo			.																
2	Juglans nigra																			
2	Vitis ligustrice																			
2	Rubus odoratus			.																
2	Ligustrum vulgare																			
3	Acer negundo			.																
3	standing dead																			
3	Vitis ligustrice			11																
3	Juglans nigra			.																
3	Asclepias glabra			.																
4	Acer negundo			.																
4	standing dead																			
4	Rhus serotina																			
4	Vitis ligustrice																			
4	Ligustrum vulgare																			

75.3

46.6

DBH Measurement Rules



Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.


Record using the tally system from 1 to 10




1



2



3




4



5

ASH CANOPY CONDITION

- 1. Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
- 2. Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
- 3. Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
- 4. >50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
- 5. Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.



A

B

C

D

E

ASH CANOPY BREAKUP CONDITION (for dead trees):

(If an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)


- All main branches contain fine twigs (newly dead).
- Over 50% of main branches have fine twigs.
- Less than 50% of main branches have fine twigs.
- Stem still standing and tertiary main branches present.
- Central stem still standing.

Cleveland Microprints

[illegible][illegible]

70.1, 41.8, 69.7

DBH Measurement Rules



Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.


Record using the tally system from 1 to 10



1



2



3




4



5

ASH CANOPY CONDITION

1. **Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
2. **Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
3. **Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
4. **>50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
5. **Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.



A

B

C

D

E

ASH CANOPY BREAKUP CONDITION (for dead trees):

(if an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)


- A: All main branches contain fine twigs (newly dead).
- B: Over 50% of main branches have fine twigs.
- C: Less than 50% of main branches have fine twigs.
- D: Stem still standing and tertiary main branches present.
- E: Central stem still standing.

10

۴۰

63.1

DBH Measurement Rules



Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.


Record using the tally system from 1 to 10



1



2



3



4



5

ASH CANOPY CONDITION

- 1. Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
- 2. Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
- 3. Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
- 4. >50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
- 5. Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.



A

B

C

D

E


ASH CANOPY BREAKUP CONDITION (for dead trees):

(If an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)


- All main branches contain fine twigs (newly dead).
- Over 50% of main branches have fine twigs.
- Less than 50% of main branches have fine twigs.
- Stem still standing and tertiary main branches present.
- Central stem still standing.

Module ID	Tree ID	Species	DBH (cm)	Ht @ DBH	Ash condition	Dead condition	ASH ONLY		
							# Ext holes	Epicormic present	Woodpecker holes
	1	<u>None present</u>							
	2								
	3								
	4								
	5								
	6								
	7								
	8								
	9								
	10								
	11								
	12								
	13								
	14								
	15								
	16								
	17								
	18								
	19								
	20								
	21								
	22								
	23								
	24								
	25								

* If Ash Condition scores 5 (dead) provide breakup score (A-E)
 Count EAB exit holes 1.25m² x 21.5m
 Woodpecker and epicormic marked present (1) or absent (0)



*** Change intensive module numbers when necessary



Map all ash trees ≥ 10cm in each module using Tree ID number

CLEVELAND METROPARKS Plant Community Assessment Program: Invasive Species Survey



Tier 1: Early detection/ Rapid response		Presence				GPS
		NE	SE	SW	NW	
Microstegium vimineum	Japanese stiltgrass					
Ranunculus ficaria	Lesser Celandine					
Cynanchum louiseae (vine)	Black Swallow-wort					
Butomus umbellatus (wetland)	Flowering Rush					
Heracleum mantegazzianum	Giant Hogweed					
Tier 2: Assess as Needed		# of Plants				comments
		NE	SE	SW	NW	
Acer platanoides	Norway Maple					
Ailanthus altissima	Tree of Heaven					
Lonicera japonica (vine)	Japanese Honeysuckle					
Lythrum salicaria (wetland)	Purple Loosestrife					
Aegopodium podagraria (G-cover)	Bishop's Goutweed					
Celastrus orbiculatus (vine)	Asian Bittersweet					
Torilis sp.	Hedgeparsley					
Conium maculatum	Poison Hemlock					
Rhamnus cathartica	Common Buckthorn (shrub)					
Berberis thunbergii	Japanese Barberry (shrub)					
Alnus glutinosa	European Alder					
Dipsacus laciniatus	Cut-leaf Teasel					
Elaeagnus umbellata	Autumn Olive (shrub)					
Lonicera maackii	Amur Honeysuckle (shrub)					
Euonymus fortunei	Wintercreeper					
Tier 3: Presence is of Interest		# of Plants				comments
		NE	SE	SW	NW	
Convallaria majalis (G-cover)	Lily of the Valley					
Coronilla varia (G-cover)	Crown Vetch					
Eleutherococcus pentaphyllus	Five-leaf Aralia (shrub)					
Pachysandra terminalis (G-cover)	Japanese Pachysandra					
Philadelphus coronarius	Mock Orange (shrub)					
Pulmonaria officinalis (G-cover)	Lungwort					
Rubus phoenicolasius	Wineberry					
Iris pseudacorus (wetland)	Yellow Flag Iris					
Ornithogalum umbellatum	Star of Bethlehem					
Viburnum opulus var. opulus	European Cranberry (shrub)					
Viburnum plicatum	Doublefile Viburnum (shrub)					
Tier 4: Widespread and abundant		Presence				comments
		NE	SE	SW	NW	
Alliaria petiolata	Garlic Mustard					
Ligustrum vulgare	Common Privet (shrub)					
L. morrowii, L. tatarica	Bush Honeysuckles (shrub)					
Phalaris arundinacea	Reed Canarygrass					
Phragmites australis (wetland)	Phragmites					
Polygonum cuspidatum	Japanese Knotweed					
Frangula alnus	Glossy Buckthorn (shrub)					
Rosa multiflora	Multiflora Rose (shrub)					
Typha angustifolia, T. x. glauca	Cattails (wetland)					
Cirsium arvense	Canada thistle					
Dipsacus fullonum	Common Teasel					
Hesperis matronalis	Dame's Rocket					
Vinca minor (G-cover)	Periwinkle					

Presence
X: yes

of Plants
1: 1-10
2: 11-50.
3: 51-100
4: 101-1,000
5: >1,000

of Plants
1: 1-10
2: 11-50.
3: 51-100
4: 101-1,000
5: >1,000

of Plants
1: 1-10
2: 11-50.
3: 51-100
4: 101-1,000
5: >1,000

Note: For Ground-cover plants record "stem #" but in comment field describe # of colonies and patch size (S,M, L)

Project Label: PCAPProject Name: 622P2015Plot No.: 1054Page: 1 of 1

mod #	species	voucher#	# shrub clumps	size class (cm) woody stems > 1m										
				1 0-<1	2 1-<2.5	3 2.5-<5	4 5-<10	5 10-<15	6 15-<20	7 20-<25	8 25-<30	9 30-<35	10 35-<40	11 >40 (record each tree)
1	<i>None present</i>													
2														
3														
4														
5														
6														
7														
8														
9														
10														

* IF EVIDENCE OF PEST OR PATHOGEN RECORD TOTAL SPECIES POPULATION IN THE PLOT EVEN THE NOT INFECTED

Strata	# of stem infected	Severity (H, M, or L)
Tree (size class 3 or above)		
Shrub (size class 2 or below including shrub clumps)		

* Write None Present if no evidence:

<i>None</i>	Beech (Fungus)	<i>None</i>	Asian Longhorned Beetle
<i>None</i>	Hemlock (HWA)	<i>None</i>	Other Pest or Pathogen
<i>None</i>	Walnut (Thousand Canker)		

Severity

High = more than 50% of leaf/needle cover exhibiting symptoms

Medium = Less than 50% of leaf/needle cover exhibiting symptoms

Low = Only a few leaves or branches are exhibiting symptoms

COVER BY STRATA

STRATUM	GENERAL FORM
Tree (generally >5 m)	Tree (overstory), very tall shrubs*, liana, epiphyte)
Shrub (generally 0.5 to 5 m)	Tree (sapling), shrub, liana, epiphyte)
Herb (Field)	Herb, dwarf-shrub**, tree (seedling***)
Floating	Floating
Aquatic (submerged)	Submerged

*Very tall shrubs are sometimes included in the tree stratum
 **Can also include seedlings of shrubs, i.e. all shrubs <0.5m
 ***Tree seedlings are often defined as up to 1.4 m height or as <2.5 cm DBH in which case they would span the herb and shrub layers.





FIGURE 3-20.—Generalized section of Upper Devonian, Mississippian, and Lower Pennsylvanian formations in northeastern Ohio. Asterisks indicate units that are fossiliferous. This composite section represents about 400 meters of rock exposed across the area. The section is not to scale, but the thicknesses indicated are proportional. The term "Waverly" is used in the older literature to refer to Mississippian rocks in Ohio. Some geologists use the European term "Carboniferous," which encompasses the Mississippian and Pennsylvanian Periods of the U.S. Many units have been named within the Cuyahoga Formation, but most units are local and cannot be traced over great distances. The Black Hand Member is a spectacular massive sandstone that is fairly widespread but discontinuous. See Hyle (1953), Hoover (1960), and Collias (1979) for more information on Mississippian rocks in Ohio. See figure 3-18 for explanation of rock types.

SOIL PIT DESCRIPTION: Excavate 20 cm plug with shovel. Describe using Munsell chart, visual exam, texture, and odor

SOIL SAMPLES Standard procedure: collect a soil sample of the top 10 cm of soil from center of each intensive module and composite the sample

Soil pit module # _____ (one per entire plot)

5 cm	matrix color	
	mottle color	
	%mottle	
	oxid roots	Y N
	texture*	
	redox features**	Y N
	hydr. cond.***	I S M D
20 cm	matrix color	
	mottle color	
	%mottle	
	oxid roots	Y N
	texture*	
	redox features**	Y N
	hydr. cond.***	I S M D

* refer to texture classes on reverse side
 ** e.g. hydrogen sulfide odor, gleying, etc.
 *** Circle one:
 I-indurated S-saturated M-moist D-dry

Notes: include evidence of earthworms (worms, castings, middens)

2-castings present
 3-castings & worms present
 8-worms & castings present
 9-castings present

Soil Collection Method	Horizon (A, B, C)
2,3,4,9 completed	A

Web Soil Survey Information

Soil Series/Type:

Soil Series Source: Ohio Soil Survey

Landform type:

Depth to root layer:

Parent Material

DETAILED*

- ☐ Excessively dr ☐ Somewhat excessively
☐ Well drained ☐ Moderately well dr
☐ Somewhat poorly dr ☐ Very poorly dr
☐ Impermeable surface

SOIL DEPTH MEASUREMENT: Measure to the nearest 0.1 cm in center of intensive modules. If >30.5 cm, record as >30

module	1 liter+ organic depth (cm)	2 liter depth (cm)	water depth (cm)	depth sat soil (cm)
2	0	0	0	0
3	0	0	0	0
8	0	0	0	0
9	0.2	0.2	0	0

EARTH SURFACE & GROUND COVER

Underlying Earth Surface*	Ground Cover	percent
Gravel - 100%	percent (Each ≤ 100%)	percent
limestone	Coarse Woody Debris***	6%
Mineral Soil	Fine Woody Debris****	3%
Gravel-Cobble*	Litter	3%
Boulder**	Duff (Ferm - Humus)	3%
Bedrock	Bryophyte Lichen	3%
* Gravel-Cobble = 1/16-10"	Water	3%
** Boulder = > 10 in	Bare Soil	3%
*** > 5 cm in diameter	Round Trail	3%
**** < 5 cm in diameter	Other	3%

COVER BY STRATA

estimate using midpoints of 5, 9, 3, 8, 13

Strata	Height Range (m)	Total Cover (%)
T tree	5 -	88%
Shrub	2 - 5	68%
Herb	0 - 2	98%
(Floating)*	-	-
(Aquatic)*	-	-

* rooted and floating or slightly emersed
 ** submerged, most plant mass below surface
 SEE BACK OF PAGE FOR TYPICAL STRATA DESCRIPTIONS. STRATA CAN VARY BY COVER TYPE.

STAND SIZE


- ☐ >600 x plot size
☒ > 100 x plot size
☐ 10-100 x plot size
☐ 3-10 x plot size
☐ 1-3 x plot size
☐ < plot size

TRAIL INFORMATION:

Type	%Cover
All Purpose	
Bridle	
Hiking sanctioned	
Boatleg unsanctioned	
Gravel	
Deer	

PERCENT MOTTLES (USE CLASS CODES):

Class	Conv.	Code NASIS	Criteria: % of Surface Area Covered
Few	f	#	< 2
Common	c	#	2 to < 20
Many	m	#	≥ 20





2%



20%


Geomorphic Component - Three-dimensional descriptors of parts of landforms or microfeatures that are best applied to areas. Unique descriptors are available for Hills, Terraces, Mountains, and Flat Plains; e.g., (for Hills) nose slope or NS.

Hills	Code POP	Code NASIS
interfluvial	IF	IF
head slope	HS	HS
nose slope	NS	NS
side slope	SS	SS
base slope	—	BS



Hillslope - Profile Position (Hillslope Position in PDP) - Two-dimensional descriptors of parts of line segments (i.e., slope position) along a transect that runs up and down the slope; e.g., backslope or BS. This is best applied to transects or points, not areas.

Position	Code
summit	SU
shoulder	SH
backslope	BS
footslope	FS
toeslope	TS



HYDROLOGIC REGIME Modified from Grossman et al 1998. (Frequency and duration of flooding.)

UPLAND: Not a wetland. Very rarely flooded.

INTERMITTENTLY/SEASONALLY SATURATED: Dry at least once per year. Surface water is seldom present, but substrate is saturated to surface for extended periods during the growing season.

PERMANENTLY/SEMIPERMANENTLY SATURATED: Dry less than once per year. Surface water is seldom present, but substrate is saturated to surface for extended periods during the growing season. Equivalent to Cowardin's Saturated modifier.

OCCASIONALLY FLOODED: Surface water can be present for brief periods during growing season, but not in most years. Often characterizes flood-plain upper terraces.

TEMPORARILY FLOODED: Surface water present for brief periods during growing season, but water table usually lies well below soil surface. Often characterizes flood-plain levees and lower terraces. Equivalent to Cowardin's Temporary modifier.

INTERMITTENTLY FLOODED: Substrate is usually exposed, but surface water can be present for variable periods without detectable seasonal periodicity. Inundation is not predictable to a given season and is dependent upon highly localized rain storms. This modifier was developed for use in the arid West for water regimes of Playa lakes, intermittent streams, and dry washes but can be used in other parts of the U.S. where appropriate. This modifier can be applied to both wetland and non-wetland situations. Equivalent to Cowardin's Intermittently Flooded modifier.

SEMIPERMANENTLY FLOODED (exposed <1/year): Surface water persists throughout the growing season in most years. Land surface is normally saturated when water level drops below soil surface. Includes Cowardin's Intermittently Exposed and Semipermanently Flooded modifiers.

PERMANENTLY FLOODED: Water covers the land surface at all times of the year in all years. Equivalent to Cowardin's "permanently flooded".

UNKNOWN: The hydrologic regime cannot be determined from the available information.