

CLEVELAND METROPARKS Plant Community Assessment Program: Quality Control Form



Project Label: _____

PCAP _____

Plot No: 1067Date Sampled: 07/14/15Lead: LANCE

Comment required if item answer is NO

Parking/Access outside of Park Boundaries:	Y <input checked="" type="radio"/> N <input type="radio"/>	If yes, write details in Comments section below
Field journals completed	Y <input checked="" type="radio"/> N <input type="radio"/>	
Site sketch made on 1:3000 map?	Y <input checked="" type="radio"/> N <input type="radio"/>	
Check cover page	Y <input checked="" type="radio"/> N <input type="radio"/>	
X-axis Bearing of plot recorded	Y <input checked="" type="radio"/> N <input type="radio"/>	
GPS coords. Recorded	Y <input checked="" type="radio"/> N <input type="radio"/>	
North direction recorded	Y <input checked="" type="radio"/> N <input type="radio"/>	
Photographs taken?	Y <input checked="" type="radio"/> N <input type="radio"/>	
Relocated Pins Mapped	Y <input checked="" type="radio"/> N <input type="radio"/>	
Plot No., Date agreement on all pages?	Y <input checked="" type="radio"/> N <input type="radio"/>	
Header data completed all pages?	Y <input checked="" type="radio"/> N <input type="radio"/>	
Cover classes recorded in all Intensive modules	Y <input checked="" type="radio"/> N <input type="radio"/>	
Browse Level By Species	Y <input checked="" type="radio"/> N <input type="radio"/>	
Woody stem quality control check	Y <input checked="" type="radio"/> N <input type="radio"/>	Check every line and cross check with the Tree Cover Sheet
Invasive plant quality control check	Y <input checked="" type="radio"/> N <input type="radio"/>	N/A
Ash trees mapped	Y <input checked="" type="radio"/> N <input type="radio"/>	N/A
Completed Forest Pest/Pathogen Datasheet	Y <input checked="" type="radio"/> N <input type="radio"/>	
Cover by Strata? (confirm cover type)	Y <input checked="" type="radio"/> N <input type="radio"/>	
Soil samples collected with matching plot #.	Y <input checked="" type="radio"/> N <input type="radio"/>	N/A
Cross check 2010 information	Y <input checked="" type="radio"/> N <input type="radio"/>	Highlight any changes from 2010 information
Vouchers labeled on datasheet with initials and number	Y <input checked="" type="radio"/> N <input type="radio"/>	
Vouchers labeled on collection bag	Y <input checked="" type="radio"/> N <input type="radio"/>	
Pink flags removed	Y <input checked="" type="radio"/> N <input type="radio"/>	
Data sheet QA before leaving site?	Y <input checked="" type="radio"/> N <input type="radio"/>	
Common equipment returned to tub.	Y <input checked="" type="radio"/> N <input type="radio"/>	
Data sheets scanned?		Enter date to left
Final data sheets scanned?		Enter date to left
Buffer Widths measured?	Y <input checked="" type="radio"/> N <input type="radio"/>	
Web Soil Survey	Y <input checked="" type="radio"/> N <input type="radio"/>	
Voucher Location	Refrigerator <input checked="" type="radio"/> N <input type="radio"/>	
(# vouchers collected)	Press (#)	Enter number to left
	Drier	Y <input checked="" type="radio"/> N <input type="radio"/>
	Identified	Y <input checked="" type="radio"/> N <input type="radio"/>
	Mounted	Y <input checked="" type="radio"/> N <input type="radio"/>
	Thrown away	Y <input checked="" type="radio"/> N <input type="radio"/>

GRTS point verification: Is plot sampleable?

<input checked="" type="checkbox"/> Yes	Original GRTS point is sampleable
<input type="checkbox"/> No	Original GRTS point lands in a non-sampleable area (fill in category below)
	<input type="checkbox"/> Point falls in a water (i.e. river, lake)
	<input type="checkbox"/> Managed mowed area (i.e. golf course, picnic area, right-of-way)
	<input type="checkbox"/> Paved area (i.e. parking lot, road)
	<input type="checkbox"/> Unsafe to sample (i.e. steep slope)
	<input type="checkbox"/> Other

Additional Comments:

--

Nested Corners

MOD 2 (2.4) 4.2

MOD 3 (2.4) 4.2

MOD 8 (2.4) 4.2

MOD 9 (4.2)

highlight

CLEVELAND METROPARKS Plant Community Assessment Program - Background Data Sheet

GENERAL INFORMATION	
Project Label:	PCAP
Project Name:	D2MS2015
Plot Name:	Wend Place to
Find Plot - Plot	
Plot No.:	1067
<input type="checkbox"/> Level 4 (no nested corners sampled) <input checked="" type="checkbox"/> Level 5 (nested corners sampled)	
Date (mm/dd/yyyy):	07/14/2015
End date (if > 1 day):	/ /
Party:	Role**
A. Lance	Plot leader
R. Eagle	Crew
D. Sweet	Bot. Asst.
E. Krauss	Crew
** Roles: Co-leader, Asst. Guide, Owner, Taxonomist, etc.	
PLOT NOT SAMPLED:	<input type="checkbox"/> Other
<input type="checkbox"/> Perm. water <input type="checkbox"/> Paved <input type="checkbox"/> Slope <input type="checkbox"/> Safety	
SAMPLING QUALITY*	
Effort Level:	subjective evaluation of how much effort put into sampling. Hurried plots may still provide good data
<input checked="" type="checkbox"/> Very thorough	
<input type="checkbox"/> Accurate	
<input type="checkbox"/> Hurried	
TAXONOMIC ACCURACY	
	high modera. low not simpl
vascular:	✓
bryo:	✓
lichen:	✓
TAXONOMIC STANDARD	
Authority:	G&C Pub Date: 1998

Minimum required fields in Bold and Underlined

LOCATION	
State:	OH County: Cuyahoga
Quadrangle:	Berea
Local Place Names:	Ohio Turnpike
Landowner:	CMP
Data Confidentiality:	<input checked="" type="checkbox"/> Public data <input type="checkbox"/> Private Data
Check one:	<input checked="" type="checkbox"/> Fuzz 100m <input type="checkbox"/> Fuzz 250m <input type="checkbox"/> Fuzz 500m
Reason:	
If data not public why?	
Source of coordinates:	<input type="checkbox"/> MAP <input checked="" type="checkbox"/> GPS
Coordinate system:	Coord. Units
<input checked="" type="checkbox"/> Lat/Long <input type="checkbox"/> UTM <input type="checkbox"/> StatePlane	<input type="checkbox"/> deg <input type="checkbox"/> deg min <input type="checkbox"/> Other (specify)
Datum:	<input checked="" type="checkbox"/> NAD83/WGS84 <input type="checkbox"/> NAD27
GPS location in plot (x=0 to 5, y=-1, 0, +1):	
x = 0 y = 0 (base of plot x=0, y=0)	
Latitude: 41.34475 - 41.34457	
Longitude: 81.24745 - 81.24691	
Coord. Accuracy:	m ft +- 2
GPS File Name:	1067AA
Plot size for cover data:	(hectares)
X-axis Bearing of plot:	114°
Depth: (1-5):	4
Intensive modules:	2, 3, 8, 9 (EDIT IF MODIFIED)
Camera No.:	3
Photo Nos.:	133
Plot placement:	<input checked="" type="checkbox"/> GRTS <input type="checkbox"/> Representative
<input type="checkbox"/> Random <input type="checkbox"/> Stratified Random <input type="checkbox"/> Transect component	
<input type="checkbox"/> Systematic (grid) <input type="checkbox"/> Capture specific feature <input type="checkbox"/> Other	


*Definitions and values in CM PCAP FOM v. 1.0 and CVS Field Guide

<p>Layout → 2x5</p> <p>Location → Approximately 425 m west of Eastland Rd. / I-80 overpass.</p> <p>Rationale → GRTS; PCAP re-sample Veg. Characteristics → Successional, disturbed plot located just west of the Ohio Turnpike. Red maple dominates the canopy; with tulip and red oak also present.</p> <p>→</p>	<p>Diagram Key: Plot origin (0,0) point, GPS location point, photo taken, with direction, location of permanent posts</p> <p>NOTES: Include Layout (any unusual shape details), Location (directions and landscape content), Rationale (why here), and Veg Characterization (description of community, dominants, strata, BROWSE). Additional notes in space on back.</p>
---	---

OVER


EXAMPLES OF PERCENT OF AREA COVERED

The following graphic can be used for various data elements to convey "Amount" or "Quality". NOTE: Within any given box, each quadrant contains the same total area covered, just different sized objects.



cover class	% cover	midpoint
1	solitary or few	0.0001
2	0-1%	0.005
3	1-2%	0.015
4	2-5%	0.035
5	5-10%	0.075
6	10-25%	0.175
7	25-50%	0.375
8	50-75%	0.625
9	75-95%	0.850
10	95-100%	0.975

Nested Corners



BROWSE RATING NARRATIVE DESCRIPTION

LOW OR NONE: there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.



MEDIUM LOW values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

MEDIUM: browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.

MEDIUM HIGH values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.

HIGH: greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.

VERY HIGH values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.



Page 2 of 2

1

1.



**Cleveland
Metroparks**


Br = Browse Level. Use cover classes to describe amount of browse per species over entire plot

Strata - Cov. entire plot



[illegible]

EXAMPLES OF PERCENT OF AREA COVERED

The following graphic can be used for various data elements to convey "Amount" or "Quality". NOTE: Within any given box, each quadrat contains the same total area covered, just different sized objects.



cover class	% cover	midpoint
1	solitary or few	0.0001
2	0-1%	0.005
3	1-2%	0.015
4	2-5%	0.035
5	5-10%	0.075
6	10-25%	0.175
7	25-50%	0.375
8	50-75%	0.625
9	75-95%	0.850
10	95-100%	0.975



BROWSE RATING NARRATIVE DESCRIPTION

LOW OR NONE: there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.

MEDIUM LOW values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

MEDIUM: browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.

MEDIUM HIGH values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.

HIGH: greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.

VERY HIGH values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.

Page of

Plot no.: _____

[illegible]

CLEVELAND METROPARKS Plant Community Assessment Program Natural Woody Stem Data Sheet

Project Label: PCAP

Project Name: PCMS2015

Plot No.: 1067

Page: 1 of 3




Explain subsample (additional room on back):

mod #	species	c	voucher#	# stems 0-1.4m browsed	% sub or super sample	# shrub clumps	size class (cm) woody stems > 1.4m	1 0-1	2 1-2.5	3 2.5-5	4 5-10	5 10-15	6 15-20	7 20-25	8 25-30	9 30-35	10 35-40	11 >40 (record each tree)
1	standing dead																	
1	Lindera benzoin			12														90.0
1	Acer rubrum																	60.2
1	Liriodendron tulipifera																	
1	standing dead																	
1	Acer rubrum																	45.6
1	Liriodendron tulipifera																	
1	Lindera benzoin			13														
1	Aster multiflorus			12														
1	standing dead																	
1	Lindera benzoin			13														53.7, 82.4
1	Acer rubrum																	
1	Corylus sp.			1														
1	Alnus incana			1														
1	standing dead																	
1	Lindera benzoin			13														41.5
1	Acer rubrum																	
1	Parthenocissus quinquefolia			3														
1	standing dead																	
1	Nyssa sylvatica																	
1	Lindera benzoin			9														
1	Acer rubrum																	
1	Acer rubrum			2														
1	Acer rubrum			3														

- Runus taken into the plot but still alive

DBH Measurement Rules



Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.

Record using the tally system from 1 to 10



1



2



3



4



5

ASH CANOPY CONDITION

1. **Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
2. **Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
3. **Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
4. **>50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
5. **Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.



A

B

C

D

E

ASH CANOPY BREAKUP CONDITION (for dead trees):

(If an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)

- A: All main branches contain fine twigs (newly dead).
- B: Over 50% of main branches have fine twigs.
- C: Less than 50% of main branches have fine twigs.
- D: Stem still standing and tertiary main branches present.
- E: Central stem still standing.

CLEVELAND METROPARKS Plant Community Assessment Program Natural Woody Stem Data Sheet

Project Label: PCAP

Project Name: 02M52015

Plot No.: 1067


Page: 2 of 3



Explain subsample (additional room on back):

mod #	species	c	voucher#	# stems 0-1.4m browed	% sub or super sample	# shrub clumps	size class (cm) woody stems >1.4m	1 0-1	2 1-2.5	3 2.5-5	4 5-10	5 10-15	6 15-20	7 20-25	8 25-30	9 30-35	10 35-40	11 >40 (record each tree)
✓6	Carpinus caroliniana			2														
✓6	standing dead																	
✓6	Smilax latifolia			3														
✓6	Acer rubrum																	
✓6	Acerus rubra																	97.6
✓6	Lindera benzoin			5														
✓6	Palmettoisus quinquefolia			6														
✓6	Rhus serotina			1														
✓7	standing dead																	
✓7	Nyssa sylvatica																	
✓7	Acerus rubra																	96.9
✓7	Acer rubrum																	105.9
✓7	Lindera benzoin			7														
✓7	Palmettoisus quinquefolia			1														
✓8	standing dead																	
✓8	Lindera benzoin			4														
✓8	Acerus rubra																	
✓8	Rubus strigosus			3														
✓8	Liriodendron			2														
✓8	Rosa multiflora			4														
✓8	Smilax latifolia			1														
✓8	Palmettoisus quinquefolia			2														
✓9	standing dead																	
✓9	Rhus serotina																	

DBH Measurement Rules



Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.

Record using the tally system from 1 to 10



1



2



3




4



5

ASH CANOPY CONDITION

- 1. Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
- 2. Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
- 3. Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
- 4. >50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
- 5. Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.



A

B

C

D

E

ASH CANOPY BREAKUP CONDITION (for dead trees):

(if an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)

- A:** All main branches contain fine twigs (newly dead).
- B:** Over 50% of main branches have fine twigs.
- C:** Less than 50% of main branches have fine twigs.
- D:** Stem still standing and tertiary main branches present.
- E:** Central stem still standing.

CLEVELAND METROPARKS Plant Community Assessment Program Natural Woody Stem Data Sheet

Project Label: PCAP

Project Name: 02M52015

Plot No.: 1067


Page: 3 of 3



Explain subsample (additional room on back):

plot #	species	c	voucher#	# stems 0-1.4m browed	% sub or super sample	# shrub clumps	size class (cm) woody stems > 1.4m										
							1 0-1	2 1-2.5	3 2.5-5	4 5-10	5 10-15	6 15-20	7 20-25	8 25-30	9 30-35	10 35-40	11 >40 (record each tree)
9	Acer rubrum			115													49.8
1	Lindera benzoin																
4	Fraxinus pennsylvanica			56													
9	Salix fluviatilis			114													
9	Smilax latifolia			1													
4	Fraxinus sp. seedling			1													
9	Asimina triloba			1													
10	Toxicodendron radicans																
10	Acer rubrum																81.2
10	Rhus typhina																
10	Crataegus sp.			1													
10	Lindera benzoin			3													
10	Hammamelis virginiana			X													
10	Staphylea trifolia			1													
10	Rosa multiflora			1													
10	Fraxinus sp.			1													

DBH Measurement Rules



Woody Stem Deer Browse


Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.

Record using the tally system from 1 to 10



ASH CANOPY CONDITION

- 1. Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
- 2. Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
- 3. Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
- 4. >50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
- 5. Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.



A

B

C

D

E

ASH CANOPY BREAKUP CONDITION (for dead trees):


(If an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)

- A:** All main branches contain fine twigs (newly dead).
- B:** Over 50% of main branches have fine twigs.
- C:** Less than 50% of main branches have fine twigs.
- D:** Stem still standing and tertiary main branches present.
- E:** Central stem still standing.


Tree ID	Species	Per c	Voucher #	DBH (cm)	Ht @ DBH	Ash condition	Dead condition	# Exit holes	Epicormic present	Woodpecker holes
1	None present									
2										
3										
4										
5										
6										
7										
8										
9										
10										
11										
12										
13										
14										
15										
16										
17										
18										
19										
20										
21										
22										
23										
24										
25										

ASH ONLY

* If Ash Condition scores 5 (dead) provide breakup score (A-E)
Count EAB exit holes 1.25m² x 21.5m
Woodpecker and epicormic marked present (1) or absent (0)



*** Change intensive module numbers when necessary



Map all ash trees ≥ 10cm in each module using Tree ID number

CLEVELAND METROPARKS Plant Community Assessment Program: Invasive Species Survey



Tier 1: Early detection/ Rapid response		Presence				GPS	Presence X: yes
		NE	SE	SW	NW		
Microstegium vimineum	Japanese stiltgrass						
Ranunculus ficaria	Lesser Celandine						
Cynanchum louiseae (vine)	Black Swallow-wort						
Butomus umbellatus (wetland)	Flowering Rush						
Heracleum mantegazzianum	Giant Hogweed						
Tier 2: Assess as Needed		# of Plants				comments	# of Plants
		NE	SE	SW	NW		
Acer platanoides	Norway Maple						1: 1-10
Ailanthus altissima	Tree of Heaven						2: 11-50.
Lonicera japonica (vine)	Japanese Honeysuckle						3: 51-100
Lythrum salicaria (wetland)	Purple Loosestrife						4: 101-1,000
Aegopodium podagraria (G-cover)	Bishop's Goutweed						5: >1,000
Celastrus orbiculatus (vine)	Asian Bittersweet						
Torilis sp.	Hedgeparsley						
Conium maculatum	Poison Hemlock						
Rhamnus cathartica	Common Buckthorn (shrub)						
Berberis thunbergii	Japanese Barberry (shrub)						
Alnus glutinosa	European Alder						
Dipsacus laciniatus	Cut-leaf Teasel						
Elaeagnus umbellata	Autumn Olive (shrub)						
Lonicera maackii	Amur Honeysuckle (shrub)						
Euonymus fortunei	Wintercreeper						
Tier 3: Presence is of Interest		# of Plants				comments	# of Plants
		NE	SE	SW	NW		
Convallaria majalis (G-cover)	Lily of the Valley						1: 1-10
Coronilla varia (G-cover)	Crown Vetch						2: 11-50.
Eleutherococcus pentaphyllus	Five-leaf Aralia (shrub)						3: 51-100
Pachysandra terminalis (G-cover)	Japanese Pachysandra						4: 101-1,000
Philadelphus coronarius	Mock Orange (shrub)						5: >1,000
Pulmonaria officinalis (G-cover)	Lungwort						
Rubus phoenicolasius	Wineberry						
Iris pseudacorus (wetland)	Yellow Flag Iris						
Ornithogalum umbellatum	Star of Bethlehem						
Viburnum opulus var. opulus	European Cranberry (shrub)						
Viburnum plicatum	Doublefile Viburnum (shrub)						
Tier 4: Widespread and abundant		Presence				comments	# of Plants
		NE	SE	SW	NW		
Alliaria petiolata	Garlic Mustard						1: 1-10
Ligustrum vulgare	Common Privet (shrub)						2: 11-50.
L. morrowii, L. tatarica	Bush Honeysuckles (shrub)						3: 51-100
Phalaris arundinacea	Reed Canarygrass						4: 101-1,000
Phragmites australis (wetland)	Phragmites						5: >1,000
Polygonum cuspidatum	Japanese Knotweed						
Frangula alnus	Glossy Buckthorn (shrub)						
Rosa multiflora	Multiflora Rose (shrub)						
Typha angustifolia, T. x.glauca	Cattails (wetland)						
Cirsium arvense	Canada thistle						
Dipsacus fullonum	Common Teasel						
Hesperis matronalis	Dame's Rocket						
Vinca minor (G-cover)	Periwinkle						

Note: For Ground-cover plants record "stem #" but in comment field describe # of colonies and patch size (S,M, L)

Project Label: PCAPProject Name: 02MS2015Plot No.: 1067Page: 1 of 1

mod #	species	voucher#	# shrub clumps	size class (cm) woody stems > 1m										
				1 0<1	2 1-<2.5	3 2.5-<5	4 5-<10	5 10 - <15	6 15 - <20	7 20 - <25	8 25 - <30	9 30 - <35	10 35 - <40	11 >40 (record each tree)
1	<u>None Present</u>													
2														
3														
4														
5														
6														
7														
8														
9														
10														

* IF EVIDENCE OF PEST OR PATHOGEN RECORD TOTAL SPECIES POPULATION IN THE PLOT EVEN THE NOT INFECTED

Strata	# of stem infected	Severity (H,M, or L)
Tree (size class 3 or above)		
Shrub (size class 2 or below including shrub clumps)		

* Write None Present if no evidence:

_____ Beech (Fungus) _____ Asian Longhorned Beetle

_____ Hemlock (HWA) _____ Other Pest or Pathogen

_____ Walnut (Thousand Canker)

Severity

High = more than 50% of leaf/needle cover exhibiting symptoms



Medium = Less than 50% of leaf/needle cover exhibiting symptoms

Low = Only a few leaves or branches are exhibiting symptoms

COVER BY STRATA

STRATUM	GENERAL FORM
Tree (generally >5 m)	Tree (overstory), very tall shrubs*, liana, epiphyte)
Shrub (generally 0.5 to 5 m)	Tree (sapling), shrub, liana, epiphyte)
Herb (Field)	Herb, dwarf-shrub**, tree (seedling***)
Floating	Floating
Aquatic (submerged)	Submerged

*Very tall shrubs are sometimes included in the tree stratum
 **Can also include seedlings of shrubs, i.e. all shrubs <0.5m
 ***Tree seedlings are often defined as up to 1.4 m height or as <2.5 cm DBH in which case they would span the herb and shrub layers.



SOIL PIT DESCRIPTION: Excavate 20 cm plug with shovel. Describe using Munsell chart, visual exam, texture, and odor.

SOIL SAMPLES Standard procedure: collect a soil sample of the top 10 cm of soil from center of each intensive module and composite the sample

Soil pit module # (one per entire plot)

5 cm	matrix color	
	mottle color	
	%mottle	
	oxid roots	Y N
	texture*	
	redox features**	Y N
	hyd. cond.***	I S M D
20 cm	matrix color	
	gristle color	
	%mottle	
	oxid roots	Y N
	texture*	
	redox features**	Y N
	hyd. cond.***	I S M D

Soil Collection Module/Section (A, B, C)	A
2.3.4.9 composite	
Hyd. Soil Survey Information	
Soil Series/Type:	
Soil Series Source: Ohio Soil Survey	
Landform type:	
Depth to root layer:	
Parent Material:	
DRAINAGE*	
<input type="checkbox"/> Excessively dr. <input type="checkbox"/> Somewhat excessively <input type="checkbox"/> Well drained <input type="checkbox"/> Moderately well dr. <input type="checkbox"/> Somewhat poorly dr. <input type="checkbox"/> Very poorly dr. <input type="checkbox"/> Impermeable surface	

SOIL DEPTH MEASUREMENT: Measure to the nearest 0.1 cm in center of intensive modules. If >30.5 cm, record as >30

	1 liter+ organic depth (cm)	2 liter depth (cm)	water depth (cm)	depth soil (cm)
mod				
2	1.5	1.5	0	0
3	1.9	1.9	8	0
8	1.2	1.2	6	0
9	1.4	6.4	0	0

2 - castings present
3 - No. worms/castings
8 - worms present
9 - castings + midens present

8 - worms present

EARTH SURFACE & GROUND COVER

Underlying Earth Surface*	Ground Cover	percent
Sum = 100%	percent (Each ≤ 100%)	percent
Hierosol	Coarse Woody Debris***	10%
Mineral Soil	Fine Woody Debris****	8%
Gravel-Cobble*	Litter	90%
Boulder**	Duff (Ferm. + Humus)	—
Bedrock	Bryophyte-Lichen	1%
* Gravel-Cobble = 1/16-10"	Water	0%
** Boulder = > 10 in	Bare Soil	1%
*** > 5 cm in diameter	Road/Trail	0%
**** < 5 cm in diameter	Other	0%

TRAIL INFORMATION:

record type and cover for each

Type	% Cover
<input type="checkbox"/> All Purpose	
<input type="checkbox"/> Bridle	
<input type="checkbox"/> Hiking sanctioned	
<input type="checkbox"/> Boodleg unsanctioned	
<input type="checkbox"/> Gravel	
<input type="checkbox"/> Deer	

COVER BY STRATA

%

estimate using midpoints of 5, ex: 3, 8, 13

Strata	Height Range (m)	Total Cover (%)
Tree	5 -	88%
Shrub	5 - 5	18%
Herb	0 - 1.5	18%
(Floating)*	-	
(Aquatic)*	-	

* rooted and floating or slightly emerged
** submersed, most plant mass below surface


SEE BACK OF PAGE FOR TYPICAL STRATA DESCRIPTIONS. STRATA CAN VARY BY COVER TYPE

STAND SIZE


- ☐ > 600 x plot size
- ☐ > 100 x plot size
- ☐ 10-100 x plot size
- ☒ 5-10 x plot size
- ☐ 1-3 x plot size
- ☐ < plot size

PERCENT MOTTLES (USE CLASS CODES):

Class	Code	Criteria: % of Surface Area Covered
Conv.	NASIS	
Few	f	< 2
Common	c	2 to < 20
Many	m	≥ 20



2%



20%


SOIL TEXTURE: Record the code for the soil texture of the 5 cm and 20 cm layers. To estimate texture, collect a soil sample from the appropriate layer and moisten it with water to the consistency of modeling clay/wet newspaper; the sample should be wet enough that all of the particles are saturated but excess water does not freely flow from the sample when squeezed. Attempt to roll the sample into a ball. If the soil will not stay in a ball and has a grainy texture, the texture is either sandy or coarse sandy. If the soil does form a ball, squeeze the sample between your fingers and attempt to form a self-supporting ribbon. Samples which form both a ball and a ribbon should be coded as clayey; samples which form a ball but not a ribbon should be coded as loamy.

- 0= Organic
- 1= Loamy
- 2= Clayey
- 3= Sandy
- 4= Coarse Sand
- 9= Not measured - make plot note

Geomorphic Component - Three-dimensional descriptors of parts of landforms or microfeatures that are best applied to areas. Unique descriptors are available for Hills, Terraces, Mountains, and Flat Plains; e.g., (for Hills) nose slope or NS.



Hills	Code	NASIS
PDP		
interfluvio	IF	IF
head slope	HS	HS
nose slope	NS	NS
side slope	SS	SS
base slope	—	BS

Terraces	Code
riser	RI
tread	TR



Hillslope - Profile Position (Hillslope Position in PDP) - Two-dimensional descriptors of parts of line segments (i.e., slope position) along a transect that runs up and down the slope; e.g., backslope or BS. This is best applied to transects or points, not areas.

Position	Code
summit	SU
shoulder	SH
backslope	BS
footslope	FS
toeslope	TS



HYDROLOGIC REGIME Modified from Grossman et al 1998. (Frequency and duration of flooding.)

UPLAND: Not a wetland. Very rarely flooded.

INTERMITTENTLY/SEASONALLY SATURATED: Dry at least once per year. Surface water is seldom present, but substrate is saturated to surface for extended periods during the growing season.

PERMANENTLY/SEMI-PERMANENTLY SATURATED: Dry less than once per year. Surface water is seldom present, but substrate is saturated to surface for extended periods during the growing season. Equivalent to Cowardin's Saturated modifier.

OCCASIONALLY FLOODED: Surface water can be present for brief periods during growing season, but not in most years. Often characterizes flood-plain upper terraces.

TEMPORARILY FLOODED: Surface water present for brief periods during growing season, but water table usually lies well below soil surface. Often characterizes flood-plain levees and lower terraces. Equivalent to Cowardin's Temporary modifier.

INTERMITTENTLY FLOODED: Substrate is usually exposed, but surface water can be present for variable periods without detectable seasonal periodicity. Inundation is not predictable to a given season and is dependent upon highly localized rain storms. This modifier was developed for use in the arid West for water regimes of Playa lakes, intermittent streams, and dry washes but can be used in other parts of the U.S. where appropriate. This modifier can be applied to both wetland and non-wetland situations. Equivalent to Cowardin's Intermittently Flooded modifier.

SEMI-PERMANENTLY FLOODED (exposed <1/year): Surface water persists throughout the growing season in most years. Land surface is normally saturated when water level drops below soil surface. Includes Cowardin's Intermittently Exposed and Semipermanently Flooded modifiers.

PERMANENTLY FLOODED: Water covers the land surface at all times of the year in all years. Equivalent to Cowardin's "permanently flooded".

UNKNOWN: The hydrologic regime cannot be determined from the available information.