

CLEVELAND METROPARKS Plant Community Assessment Program: Quality Control Form



Project Label:

PCAP

Plot No:

1064

Date Sampled:

7/27/15

Lead:

CKM

Comment required if item answer is NO

Parking/Access outside of Park Boundaries:	Y	<input checked="" type="radio"/> N	If yes, write details in Comments section below
Field journals completed	<input checked="" type="radio"/> Y	N	
Site sketch made on 1:3000 map?	<input checked="" type="radio"/> Y	N	
Check cover page	<input checked="" type="radio"/> Y	N	
X-axis Bearing of plot recorded	<input checked="" type="radio"/> Y	N	
GPS coords. Recorded	<input checked="" type="radio"/> Y	N	
North direction recorded	<input checked="" type="radio"/> Y	N	
Photographs taken?	<input checked="" type="radio"/> Y	N	
Relocated Pins Mapped	<input checked="" type="radio"/> Y	N	
Plot No., Date agreement on all pages?	<input checked="" type="radio"/> Y	N	
Header data completed all pages?	<input checked="" type="radio"/> Y	N	
Cover classes recorded in all Intensive modules	<input checked="" type="radio"/> Y	N	
Browse Level By Species	<input checked="" type="radio"/> Y	N	
Woody stem quality control check	<input checked="" type="radio"/> Y	N	Check every line and cross check with the Tree Cover Sheet
Invasive plant quality control check	Y	N	NA
Ash trees mapped	<input checked="" type="radio"/> Y	N	
Completed Forest Pest/Pathogen Datasheet	<input checked="" type="radio"/> Y	N	
Cover by Strata? (confirm cover type)	<input checked="" type="radio"/> Y	N	
Soil samples collected with matching plot #.	Y	N	NA
Cross check 2010 information	<input checked="" type="radio"/> Y	N	Highlight any changes from 2010 information
Vouchers labeled on datasheet with initials and number	<input checked="" type="radio"/> Y	N	
Vouchers labeled on collection bag	<input checked="" type="radio"/> Y	N	
Pink flags removed	<input checked="" type="radio"/> Y	N	
Data sheet QA before leaving site?	<input checked="" type="radio"/> Y	N	
Common equipment returned to tub.	Y	N	
Data sheets scanned?			Enter date to left
Final data sheets scanned?			Enter date to left
Buffer Widths measured?	Y	N	
Web Soil Survey	Y	N	
Voucher Location	Refrigerator	Y	N
(# vouchers collected)	Press (#)		Enter number to left
CKM 239-240	Drier	Y	N
	Identified	Y	N
	Mounted	Y	N
	Thrown away	Y	N

GRTS point verification: Is plot sampleable?

<input type="checkbox"/> Yes	Original GRTS point is sampleable
<input type="checkbox"/> No	Original GRTS point lands in a non-sampleable area (fill in category below)
	<input type="checkbox"/> Point falls in a water (i.e. river, lake)
	<input type="checkbox"/> Managed mowed area (i.e. golf course, picnic area, right-of-way)
	<input type="checkbox"/> Paved area (i.e. parkinglot, road)
	<input type="checkbox"/> Unsafe to sample (i.e. steep slope)
	<input type="checkbox"/> Other

Additional Comments:

--


GENERAL INFORMATION			
Project Label: PCAP			
Project Name: 025C2015			
Plot Name: Fallen Spicebush			
Plot No.: 1064			
<input type="checkbox"/> Level 4 (no nested corners sampled) <input checked="" type="checkbox"/> Level 5 (nested corners sampled)			
Date (mm/dd/yyyy): 7/27/2015			
End date (if > 1 day): / /			
Party	Role**		
C. Minney	Plot leader		
M. Geitgey	Bot. Asst.		
E. Knauss	Woody Tech		
T. Cochran	Woody Tech		
** Roles: Co-leader, Asst. Graphic, Owner, Taxonomist, etc.			
PLOT NOT SAMPLED: <input type="checkbox"/> Other			
<input type="checkbox"/> Perm water <input type="checkbox"/> Paved <input type="checkbox"/> Slope <input type="checkbox"/> Safety SAMPLING QUALITY*			
Effort Level:	subjective evaluation of how much effort put into sampling. Hurried plots may still provide good data		
<input checked="" type="checkbox"/> Very thorough			
<input type="checkbox"/> Accurate			
<input type="checkbox"/> Hurried			
TAXONOMIC ACCURACY			
	high	modera.	low
vascul.	<input checked="" type="checkbox"/>		not sampl
bryo		<input checked="" type="checkbox"/>	n/a
lichen			<input checked="" type="checkbox"/>
TAXONOMIC STANDARD			
Authority:	G&C	Pub Date:	1998

LOCATION	
State: OH	County: Cuyahoga
Quadrangle: Chagrin Falls	
Local Place Names: Horse Trail and Arborlane	
Landowner: CMP	
Data Confidentiality:	
<input checked="" type="checkbox"/> Public data <input type="checkbox"/> Private Data <input type="checkbox"/> Fuzz 100m <input type="checkbox"/> Fuzz 250m <input type="checkbox"/> Fuzz 500m	
Reasons:	
If data not public why?	
Source of coordinates: <input type="checkbox"/> MAP <input checked="" type="checkbox"/> GPS	Coord. Units
Coordinate system:	<input checked="" type="checkbox"/> Lat/Long <input type="checkbox"/> UTM <input type="checkbox"/> StaticPlane <input type="checkbox"/> deg <input type="checkbox"/> deg min <input type="checkbox"/> Other (specify) <input type="checkbox"/> m <input type="checkbox"/> ft <input type="checkbox"/>
Datum: <input checked="" type="checkbox"/> NAD83/MGSR4 <input type="checkbox"/> NAD27	
GPS location in plot x=0 to 5, y=-1.0,+1):	
x = 0 y = 0 (base of plot x=0, y=0)	
Latitude: 41.41441	
Longitude: 81.42733	
Coord. Accuracy: <input checked="" type="checkbox"/> m <input type="checkbox"/> ft +- 3	
GPS File Name: 1064A	
Plot size for cover data: -1 (hectares)	
X-axis Bearing of plot: [03]°	
Depth: (1-5): 4	
Intensive modules: 2, 3, 8, 9 (EDIT IF MODIFIED)	
Camera No.: 4	
Photo Nos.: CH670	
Plot placement: <input checked="" type="checkbox"/> GRTS <input type="checkbox"/> Representative	
<input type="checkbox"/> Random <input type="checkbox"/> Stratified Random <input type="checkbox"/> Transect component <input type="checkbox"/> Systematic (grid) <input type="checkbox"/> Capture specific feature <input type="checkbox"/> Other	

Minimum required fields in Bold and Underlined

*Definitions and values in CM PCAP FOM v. 1.0 and CVS Field Guide

OVER



NOTES: Include Layout (any unusual shape details), Location (directions and landscape content), Rationale (why here), and Veg Characterization (description of community, dominants, strata, BROWSE). Additional notes in space on back.

Layout: 2x5

Location: From Arbor lane turn right on Bedford Chagrin Pkwy driving ~1000ft to grassy area just down slope of bridal trail crossing. Park here if ground is dry. Plot 80m off road

Rationale: GRTS

Veg Characterization: The canopy is dominated by Sugar maple with several other trees spread evenly throughout. The shrub layer is dominated by sugar maple and beech. The herb layer is very sparse dominated by May-apple and tree seedlings.

CLEVELAND METROPARKS Plant Community Assessment Program - Background Data Sheet



Page 2 of 2

Plot No.: 1064

Project Name: 02 SC 2015

Project Label: PCAP

MODIFIED NATURE RESERVE CLASS*

CODE (on separate form):

Fit= Conf=

CO2

COMMUNITY NAME:

Beech-Maple Forest

HOMOGENEITY

☒ Homogeneous ☐ Compositional trend across the plot

☐ Conspicuous inclusions ☐ Irregular/pattern mosaic

DISTURBANCES

type*	severity**	yrs ago	% of plot	description
Human				
Natural				
Fire				
Cut				
Animal	VH	0	100	Deer Browse
Other				

**L=low, ML=med low, M=med, MH=med high, H=high, VH=very high

Current Land Use: CMP

Former Land Use:

HYDROLOGIC REGIME*

☒ Upland (seldom flooded)

☐ Intermittently/seasonally saturated

(seldom flooded)

☐ Permanently/Semipermanent, saturated

(dry <1/yr, seldom flooded)

☐ Occasionally flooded (<1/yr)

☐ Temporarily flooded

☐ Intermittently flooded

☐ Semipermanently flooded

☐ Permanently flooded

☐ Tidal/Seiche flooded daily

☐ Tidal/Seiche flooded monthly

☐ Tidal/Seiche flooded irregular

(e.g. wind, storms)

☐ Unknown


(by default unless plot is a wetland)

Additional notes & diagrams: (Representativeness of plot to the stand, successional status, maturity, etc.)

The age is un-even aged. There are some really nice mature trees in the plot including three nice Cherrys. The canopy has many small sugar maples and a few beech coming in as the next generation. Plot hit hard by worms and deer browse, the herb layer sparse and not diverse. Shrub layer thick but some trees dying out.



EXAMPLES OF PERCENT OF AREA COVERED

The following graphic can be used to convert data elements to canopy "Amount" or "Quantity". NOTE: Within any given bar, each quadrant contains the same but area covered, just different sized objects.



cover class	% cover	midpoint
1	solitary or few	0.0001
2	0-1%	0.005
3	1-2%	0.015
4	2-5%	0.035
5	5-10%	0.075
6	10-25%	0.175
7	25-50%	0.375
8	50-75%	0.625
9	75-95%	0.850
10	95-100%	0.975

Nested Corners



BROWSE RATING NARRATIVE DESCRIPTION

LOW OR NONE: there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.

MEDIUM LOW values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

MEDIUM: browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.


MEDIUM HIGH values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.

HIGH: greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.

VERY HIGH values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.



EXAMPLES OF PERCENT OF AREA COVERED

The following graphic can be used for various data elements to convey "Amount" or "Quantity". NOTE: Within any given box, each quadrant contains the same total area covered, just different sized objects.



cover class	% cover	midpoint
1	solitary or few	0.0001
2	0-1%	0.005
3	1-2%	0.015
4	2-5%	0.035
5	5-10%	0.075
6	10-25%	0.175
7	25-50%	0.375
8	50-75%	0.625
9	75-95%	0.850
10	95-100%	0.975

Nested Corners



BROWSE RATING NARRATIVE DESCRIPTION

LOW OR NONE: there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.

MEDIUM LOW values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

MEDIUM: browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.

MEDIUM HIGH values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.

HIGH: greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.

VERY HIGH values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.

Page 7 of 7

Plot no.: 1064

25
6
109
5

Plot no.: _____

[illegible]

CLEVELAND METROPARKS Plant Community Assessment Program Natural Woody Stem Data Sheet

Project Label: PCAP

Project Name: OASC2015

Plot No.: 1064

Page: 1 of 3


Cleveland Metroparks

Explain subsample (additional room on back):

mod #	species	c	voucher#	# stems 0-1.4m browased	% sub or super sample	# shrub clumps	size class (cm) woody stems > 1.4m	1 0-1	2 1-2.5	3 2.5-5	4 5-10	5 10-15	6 15-20	7 20-25	8 25-30	9 30-35	10 35-40	11 >40 (record each tree)
1	STANDING DEAD																	
1	Fagus grandifolia																	
1	Acer saccharum																	
1	Quercus rubra																	
1	Acer rubrum																	84.8
1	NO BROWSE																	
2	Acer saccharum																	
2	STANDING DEAD																	
2	Fagus NO BROWSE																	
3	Fagus grandifolia																	
3	STANDING DEAD																	
3	Acer saccharum																	47.4
3	NO BROWSE																	
4	Fagus grandifolia																	
4	Acer saccharum																	
4	STANDING DEAD																	
4	Carya cordiformis																	14.8
4	Liriodendron																	
5	Acer rubrum																	
5	Acer saccharum																	
5	STANDING DEAD																	
5	Fagus grandifolia																	
5	Quercus rubra																	79.6
5	Fraxinus sp.																	

*2010 MISID'd as Carya glabra

DBH Measurement Rules



Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.


Record using the tally system from 1 to 10



1



2



3




4



5

ASH CANOPY CONDITION

- 1. Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
- 2. Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
- 3. Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
- 4. >50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
- 5. Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.



A

B

C

D

E

ASH CANOPY BREAKUP CONDITION (for dead trees):

(If an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)

- A:** All main branches contain fine twigs (newly dead).
- B:** Over 50% of main branches have fine twigs.
- C:** Less than 50% of main branches have fine twigs.
- D:** Stem still standing and tertiary main branches present.
- E:** Central stem still standing.

CLEVELAND METROPARKS Plant Community Assessment Program Natural Woody Stem Data Sheet

Project Label: PCAP

Project Name: 02.5C2015

Plot No.: 1064


Page: 2 of 3



Explain subsample (additional room on back):

mod #	species	c	voucher#	# stems 0-1.4m browsed	% sub or super sample	# shrub clumps	size class (cm) woody stems >1.4m	1 0-1	2 1-2.5	3 2.5-5	4 5-10	5 10-15	6 15-20	7 20-25	8 25-30	9 30-35	10 35-40	11 >40 (record each tree)
✓ 10	Prunus serotina																	103.4
✓ 10	STANDING DEAD																	
✓ 10	Fagus grandifolia																	
✓ 10	Acer saccharum																	74.0
✓ 10	Ostrya virginiana																	
✓ 10	Vitis aestivalis																	
✓ 10	Quercus rubra																	
✓ 10	Acer rubrum																	
✓ 10	NO BEAUSE																	
✓ 10	Fagus grandifolia																	
✓ 10	Acer saccharum																	40.6, 42.5
✓ 10	Vitis aestivalis																	
✓ 10	Lindera benzoin																	
✓ 10	Eucalyptus alatus																	
✓ 10	Pyrus sp.																	
✓ 10	Carpinus caroliniana																	
✓ 10	Fagus grandifolia																	
✓ 10	Acer saccharum																	55.9, 46.0
✓ 10	STANDING DEAD																	
✓ 10	Carpinus caroliniana																	
✓ 10	Ostrya virginiana																	
✓ 10	Pinus strobus																	63.1
✓ 10	Lindera benzoin																	
✓ 10	Carya ovata																	44.3

DBH Measurement Rules



Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.


Record using the tally system from 1 to 10



1



2



3




4



5

ASH CANOPY CONDITION

1. **Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
2. **Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
3. **Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
4. **>50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
5. **Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.



A

B

C

D

E

ASH CANOPY BREAKUP CONDITION (for dead trees):

(if an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)

- A: All main branches contain fine twigs (newly dead).
- B: Over 50% of main branches have fine twigs.
- C: Less than 50% of main branches have fine twigs.
- D: Stem still standing and tertiary main branches present.
- E: Central stem still standing.


1) $\alpha T / \alpha U$

Clareland Markets

Clareland Markets

mod #	species	browsed	sample	chumps
c	voucher#			
		0-1	1-2.5	2.5-5
		5-10	10-15	15-20
		20-25	25-30	30-35
		35-40		
		>40 (record each time)		

DBH Measurement Rules



Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.


Record using the tally system from 1 to 10



1



2



3



4



5

ASH CANOPY CONDITION

- 1. Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
- 2. Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
- 3. Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
- 4. >50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
- 5. Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.



A

B

C

D

E


ASH CANOPY BREAKUP CONDITION (for dead trees):

(If an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)


- All main branches contain fine twigs (newly dead).
- Over 50% of main branches have fine twigs.
- Less than 50% of main branches have fine twigs.
- Stem still standing and tertiary main branches present.
- Central stem still standing.

Tree ID	Species	Dead	c	Voucher #	DBH (cm)	Ht @ DBH	ASH ONLY			
							Ash condition	Dead condition	# Exit holes	Epicormic present
1	<u>None present</u>									
2										
3										
4										
5										
6										
7										
8										
9										
10										
11										
12										
13										
14										
15										
16										
17										
18										
19										
20										
21										
22										
23										
24										
25										

• If Ash Condition scores 5 (dead) provide breakup score (A-E)
 Count EAB exit holes 1.25m² x 21.5m
 Woodpecker and epicormic marked present (1) or absent (0)



*** Change intensive module numbers when necessary



Map all ash trees ≥ 10cm in each module using Tree ID number

CLEVELAND METROPARKS Plant Community Assessment Program: Invasive Species Survey



Tier 1: Early detection/ Rapid response		Presence				GPS	Presence X: yes
		NE	SE	SW	NW		
Microstegium vimineum	Japanese stiltgrass						
Ranunculus ficaria	Lesser Celandine						
Cynanchum louiseae (vine)	Black Swallow-wort						
Butomus umbellatus (wetland)	Flowering Rush						
Heracleum mantegazzianum	Giant Hogweed						
Tier 2: Assess as Needed		# of Plants				comments	# of Plants
		NE	SE	SW	NW		
Acer platanoides	Norway Maple						1: 1-10
Ailanthus altissima	Tree of Heaven						2: 11-50.
Lonicera japonica (vine)	Japanese Honeysuckle						3: 51-100
Lythrum salicaria (wetland)	Purple Loosestrife						4: 101-1,000
Aegopodium podagraria (G-cover)	Bishop's Goutweed						5: >1,000
Celastrus orbiculatus (vine)	Asian Bittersweet						
Torilis sp.	Hedgeparsley						
Conium maculatum	Poison Hemlock						
Rhamnus cathartica	Common Buckthorn (shrub)						
Berberis thunbergii	Japanese Barberry (shrub)						
Alnus glutinosa	European Alder						
Dipsacus laciniatus	Cut-leaf Teasel						
Elaeagnus umbellata	Autumn Olive (shrub)						
Lonicera maackii	Amur Honeysuckle (shrub)						
Euonymus fortunei	Wintercreeper						
Tier 3: Presence is of Interest		# of Plants				comments	# of Plants
		NE	SE	SW	NW		
Convallaria majalis (G-cover)	Lily of the Valley						1: 1-10
Coronilla varia (G-cover)	Crown Vetch						2: 11-50.
Eleutherococcus pentaphyllus	Five-leaf Aralia (shrub)						3: 51-100
Pachysandra terminalis (G-cover)	Japanese Pachysandra						4: 101-1,000
Philadelphus coronarius	Mock Orange (shrub)						5: >1,000
Pulmonaria officinalis (G-cover)	Lungwort						
Rubus phoenicolasius	Wineberry						
Iris pseudacorus (wetland)	Yellow Flag Iris						
Ornithogalum umbellatum	Star of Bethlehem						
Viburnum opulus var. opulus	European Cranberry (shrub)						
Viburnum plicatum	Doublefile Viburnum (shrub)						
Tier 4: Widespread and abundant		Presence				comments	# of Plants
		NE	SE	SW	NW		
Alliaria petiolata	Garlic Mustard						1: 1-10
Ligustrum vulgare	Common Privet (shrub)						2: 11-50.
L. morrowii, L. tatarica	Bush Honeysuckles (shrub)						3: 51-100
Phalaris arundinacea	Reed Canarygrass						4: 101-1,000
Phragmites australis (wetland)	Phragmites						5: >1,000
Polygonum cuspidatum	Japanese Knotweed						
Frangula alnus	Glossy Buckthorn (shrub)						
Rosa multiflora	Multiflora Rose (shrub)						
Typha angustifolia, T. x.glauca	Cattails (wetland)						
Cirsium arvense	Canada thistle						
Dipsacus fullonum	Common Teasel						
Hesperis matronalis	Dame's Rocket						
Vinca minor (G-cover)	Periwinkle						

Note: For Ground-cover plants record "stem #" but in comment field describe # of colonies and patch size (S,M, L)

CLEVELAND METROPARKS Plant Community Assessment Program Forest Pest and Pathogens Data Sheet

Project Label:

PCAP

Project Name:

02SC 2015

Plot No.:

1064

Page:

1 of 1



mod #	species	voucher#	# shrub clumps	size class (cm) woody stems > 1m										
				1 0-<1	2 1-<2.5	3 2.5-<5	4 5-<10	5 10 - <15	6 15 - <20	7 20 - <25	8 25 - <30	9 30 - <35	10 35 - <40	11 >40 (record each tree)
1														
2														
3														
4														
5														
6														
7														
8														
9														
10														

* IF EVIDENCE OF PEST OR PATHOGEN RECORD TOTAL SPECIES POPULATION IN THE PLOT EVEN THE NOT INFECTED

Strata	# of stem infected	Severity (H, M, or L)
Tree (size class 3 or above)		
Shrub (size class 2 or below including shrub clumps)		

* Write None Present if no evidence:

~~NONE PRESENT~~ Beech (Fungus)

~~NONE PRESENT~~ Asian Longhorned Beetle

Hemlock (HWA)

Other Pest or Pathogen

Walnut (Thousand Canker)

Severity

High = more than 50% of leaf/needle cover exhibiting symptoms

Medium = Less than 50% of leaf/needle cover exhibiting symptoms

Low = Only a few leaves or branches are exhibiting symptoms

STANDING BIOMASS (required for canopy wetland) collected in 0 in clip plot (32x32 cm) from corners 1 and 3 in each intensive module. Required for VIBR-E score calculation. C7=check when collected

Module #	C7	Corner	Corner

CLASSIFICATION

FTT = excellent, F and Confidence

Hydrogeomorphic class (WETLANDS ONLY)

<input type="checkbox"/> DEPRESSION	Fit =	Conf =
<input type="checkbox"/> ANPOUNDMENT <input type="checkbox"/> Beaver <input type="checkbox"/> Human	Fit =	Conf =
<input type="checkbox"/> RIVERINE <input type="checkbox"/> Headwater <input type="checkbox"/> Mainstem <input type="checkbox"/> Channel	Fit =	Conf =
<input type="checkbox"/> SLOPE (ground water in ditches or on a physical slope)	Fit =	Conf =
<input type="checkbox"/> FRINGING <input type="checkbox"/> Reservoir <input type="checkbox"/> Natural Lake	Fit =	Conf =
<input type="checkbox"/> COASTAL (specific subclases)	Fit =	Conf =
<input type="checkbox"/> BOG (strongly, moderately, weakly ombrotrophic)	Fit =	Conf =

Ohio EPA VIBR Plant Community Class (WETLANDS ONLY)

<input type="checkbox"/> FOREST <input type="checkbox"/> swamp forest <input type="checkbox"/> bog forest <input type="checkbox"/> forest seep	Fit =	Conf =
<input type="checkbox"/> EMERGENT <input type="checkbox"/> marsh <input type="checkbox"/> wet meadow <input type="checkbox"/> open bog	Fit =	Conf =
<input type="checkbox"/> SHRUB <input type="checkbox"/> shrub swamp <input type="checkbox"/> tall sh. bog <input type="checkbox"/> tall sh. fen	Fit =	Conf =

MICROTOPOGRAPHIC FEATURE COUNTS - Intensive modules only

Plants for microhabitat features. Select one or select two and average the scores. NOTE: If mod tabs on a slope automatically gate ranked based on steepness (1-3) to begin any features present
Slope 1 = slight elevational grade across module (1M) Slope 2 = tabs on slope -20° Slope 3 = maximum steepness that can be safely sampled -45°

- feature is absent or functionally absent from the wetland
- feature is present in the wetland in very small amounts or if more common, of low quality
- feature is present in moderate amounts, but not of highest quality, or in small amounts of highest quality
- feature is present in moderate or greater amounts and of highest quality

		no. of		no. of		no. macro.		C.W.D.		C.W.D.		C.W.D.		microhab.		microhab.	
		tussocks		hummocks		depressions		(2-12 cm)		(12-18cm)		>40 cm		interspers.		SLOPE	
		depth 3		depth 2		depth 1		depth 1		depth 1		depth 1		depth 1		1hr:10m	
		1x1m		3 lckx1.6cm		10x10m		10x10m		10x10m		10x10m		10x10m		(rank)	
module	corner	(count)	(count)	(count)	(count)	(count)	(count)	(count)	(count)	(count)	(count)	(count)	(count)	(rank)	(rank)	(rank)	(rank)
2		0	0	0	0	2	2	15	0	0	0	0	0	2	2	1	1
3		0	0	0	0	2	2	10	0	0	0	0	0	2	2	1	1
4		0	0	0	0	2	2	7	0	0	0	0	0	2	2	1	1

NOTE: tussock and hummocks are counted in BDTT nested quadrat corners but counts are aggregated.

MONAB INDICES (degrees) + for up - for down

FILLED OUT USING GIS PROGRAM - DO NOT FILL OUT IN FIELD

AI aspect	N	SE	E	SE	S	SW	W	NW
+45 degrees								
+90 degrees								
+135 degrees								
+180 degrees								
+225 degrees								
+270 degrees								
+315 degrees								

Landform Index (position within landscape)
Tern Shape Index (aka microtopographic shape)

CROWN COVER (DENSIMETER) NALE-4
readings per module facing N, S, E, W. Place dot count in
corresponding space. (4 dots per grid square)

Module	N	S	E	W
2	0	0	0	0
3	0	0	0	0
4	0	0	0	0
9	1	1	2	1

COVER BY STRATA

STRATUM	GENERAL FORM
Tree (generally >5 m)	Tree (overstory), very tall shrubs*, liana, epiphyte)
Shrub (generally 0.5 to 5 m)	Tree (sapling), shrub, liana, epiphyte)
Herb (Field)	Herb, dwarf-shrub**, tree (seedling***)
Floating	Floating
Aquatic (submerged)	Submerged

*Very tall shrubs are sometimes included in the tree stratum
 **Can also include seedlings of shrubs, i.e. all shrubs <0.5m
 ***Tree seedlings are often defined as up to 1.4 m height or as <2.5 cm DBH in which case they would span the herb and shrub layers.







FIGURE 3-20.—Generalized section of Upper Devonian, Mississippian, and Lower Pennsylvanian formations in northeastern Ohio. Asterisks indicate units that are fossiliferous. This composite section represents about 400 meters of rock exposed across the area. The section is not to scale, but the thicknesses indicated are proportional. The term "Waverly" is used in the older literature to refer to Mississippian rocks in Ohio. Some geologists use the European term "Carboniferous," which encompasses the Mississippian and Pennsylvanian Periods of the U.S. Many units have been named within the Cuyahoga Formation, but most units are local and cannot be traced over great distances. The Black Hand Member is a spectacular massive sandstone that is fairly widespread but discontinuous. See Hyde (1953), Hoover (1960), and Collins (1979) for more information on Mississippian rocks in Ohio. See figure 3-18 for explanation of rock types.

PERCENT MOTTLES (USE CLASS CODES):

Class	Code	Criteria: % of Surface Area Covered
Few	f	< 2
Common	c	2 to < 20
Many	m	≥ 20



2%




20%

SOIL TEXTURE: Record the code for the soil texture of the 5 cm and 20 cm layers. To estimate texture, collect a soil sample from the appropriate layer and moisten it with water to the consistency of modeling clay/wet newspaper; the sample should be wet enough that all of the particles are saturated but excess water does not freely flow from the sample when squeezed. Attempt to roll the sample into a ball. If the soil will not stay in a ball and has a grainy texture, the texture is either sandy or coarse sandy. If the soil does form a ball, squeeze the sample between your fingers and attempt to form a self-supporting ribbon. Samples which form both a ball and a ribbon should be coded as clayey; samples which form a ball but not a ribbon should be coded as loamy.

- 0= Organic
- 1= Loamy
- 2= Clayey
- 3= Sandy
- 4= Coarse Sand
- 9= Not measured - make plot note


Geomorphic Component - Three-dimensional descriptors of parts of landforms or microfeatures that are best applied to areas. Unique descriptors are available for Hills, Terraces, Mountains, and Flat Plains: e.g., (for Hills) nose slope or NS.

Hills	Code	NASIS
interfluvial	IF	IF
head slope	HS	HS
nose slope	NS	NS
side slope	SS	SS
base slope	BS	BS



Hillslope - Profile Position (Hillslope Position in PDP) - Two-dimensional descriptors of parts of line segments (i.e., slope position) along a transect that runs up and down the slope: e.g., *backslope* or *BS*. This is best applied to transects or points, not areas.

Position	Code
summit	SU
shoulder	SH
backslope	BS
footslope	FS
toeslope	TS



HYDROLOGIC REGIME Modified from Grossman et al 1998. (Frequency and duration of flooding.)

UPLAND: Not a wetland. Very rarely flooded.

INTERMITTENTLY/SEASONALLY SATURATED: Dry at least once per year. Surface water is seldom present, but substrate is saturated to surface for extended periods during the growing season.

PERMANENTLY/SEMIPERMANENTLY SATURATED: Dry less than once per year. Surface water is seldom present, but substrate is saturated to surface for extended periods during the growing season. Equivalent to Cowardin's Saturated modifier.

OCCASIONALLY FLOODED: Surface water can be present for brief periods during growing season, but not in most years. Often characterizes flood-plain upper terraces.

TEMPORARILY FLOODED: Surface water present for brief periods during growing season, but water table usually lies well below soil surface. Often characterizes flood-plain levees and lower terraces. Equivalent to Cowardin's Temporary modifier.

INTERMITTENTLY FLOODED: Substrate is usually exposed, but surface water can be present for variable periods without detectable seasonal periodicity. Inundation is not predictable to a given season and is dependent upon highly localized rain storms. This modifier was developed for use in the arid West for water regimes of Playa lakes, intermittent streams, and dry washes but can be used in other parts of the U.S. where appropriate. This modifier can be applied to both wetland and non-wetland situations. Equivalent to Cowardin's Intermittently Flooded modifier.

SEMIPERMANENTLY FLOODED (exposed <1/year): Surface water persists throughout the growing season in most years. Land surface is normally saturated when water level drops below soil surface. Includes Cowardin's Intermittently Exposed and Semipermanently Flooded modifiers.

PERMANENTLY FLOODED: Water covers the land surface at all times of the year in all years. Equivalent to Cowardin's "permanently flooded".

UNKNOWN: The hydrologic regime cannot be determined from the available information.