

CLEVELAND METROPARKS Plant Community Assessment Program: Quality Control Form

Cleveland Metroparks

Project Label:

PCAP

Plot No:

1074

Date Sampled:

8/3/15

Lead:

CKM

Comment required if item answer is NO

Parking/Access outside of Park Boundaries:	<input checked="" type="radio"/> Y	N	If yes, write details in Comments section below
Field journals completed	<input checked="" type="radio"/> Y	N	
Site sketch made on 1:3000 map?	<input checked="" type="radio"/> Y	N	
Check cover page	<input checked="" type="radio"/> Y	N	
X-axis Bearing of plot recorded	<input checked="" type="radio"/> Y	N	
GPS coords. Recorded	<input checked="" type="radio"/> Y	N	
North direction recorded	<input checked="" type="radio"/> Y	N	
Photographs taken?	<input checked="" type="radio"/> Y	N	
Relocated Pins Mapped	<input checked="" type="radio"/> Y	N	
Plot No., Date agreement on all pages?	<input checked="" type="radio"/> Y	N	
Header data completed all pages?	<input checked="" type="radio"/> Y	N	
Cover classes recorded in all Intensive modules	<input checked="" type="radio"/> Y	N	
Browse Level By Species	<input checked="" type="radio"/> Y	N	
Woody stem quality control check	<input checked="" type="radio"/> Y	N	Check every line and cross check with the Tree Cover Sheet
Invasive plant quality control check	<input checked="" type="radio"/> Y	N	NA
Ash trees mapped	<input checked="" type="radio"/> Y	N	
Completed Forest Pest/Pathogen Datasheet	<input checked="" type="radio"/> Y	N	
Cover by Strata? (confirm cover type)	<input checked="" type="radio"/> Y	N	
Soil samples collected with matching plot #.	<input checked="" type="radio"/> Y	N	NA
Cross check 2010 information	<input checked="" type="radio"/> Y	N	Highlight any changes from 2010 information
Vouchers labeled on datasheet with initials and number	<input checked="" type="radio"/> Y	N	
Vouchers labeled on collection bag	<input checked="" type="radio"/> Y	N	
Pink flags removed	<input checked="" type="radio"/> Y	N	
Data sheet QA before leaving site?	<input checked="" type="radio"/> Y	N	
Common equipment returned to tub.	<input checked="" type="radio"/> Y	N	
Data sheets scanned?			Enter date to left
Final data sheets scanned?			Enter date to left
Buffer Widths measured?	<input checked="" type="radio"/> Y	N	
Web Soil Survey	<input checked="" type="radio"/> Y	N	
Voucher Location	Refrigerator	<input checked="" type="radio"/> Y	N
(# vouchers collected)	Press (#)		Enter number to left
CKM 307-313	Drier	<input checked="" type="radio"/> Y	N
	Identified	<input checked="" type="radio"/> Y	N
	Mounted	<input checked="" type="radio"/> Y	N
	Thrown away	<input checked="" type="radio"/> Y	N

GRTS point verification: Is plot sampleable?

<input checked="" type="checkbox"/> Yes	Original GRTS point is sampleable
<input type="checkbox"/> No	Original GRTS point lands in a non-sampleable area (fill in category below)
	<input type="checkbox"/> Point falls in a water (i.e. river, lake)
	<input type="checkbox"/> Managed mowed area (i.e. golf course, picnic area, right-of-way)
	<input type="checkbox"/> Paved area (i.e. parking lot, road)
	<input type="checkbox"/> Unsafe to sample (i.e. steep slope)
	<input type="checkbox"/> Other

Additional Comments:

Found all pins, might have parked on Airport property? Near intersection of Rude and Spafford. Paved road beside fenced runway light.

Last data sheet said recycling area which is closer but small and a truck was pulled in as we left.

D

P

1. The first part of the paper is devoted to a discussion of the

main results of the paper. The second part is devoted to a

discussion of the main results of the paper. The third part is

devoted to a discussion of the main results of the paper. The

fourth part is devoted to a discussion of the main results of the

paper. The fifth part is devoted to a discussion of the main results

of the paper. The sixth part is devoted to a discussion of the main

results of the paper. The seventh part is devoted to a discussion of

the main results of the paper. The eighth part is devoted to a

discussion of the main results of the paper. The ninth part is

CLEVELAND METROPARKS Plant Community Assessment Program - Background Data Sheet

GENERAL INFORMATION	
Project Label:	PCAP
Project Name:	02KR2015
Plot Name:	Plane Spotting
Plot No.:	1074
<input type="checkbox"/> Level 4 (no nested corners sampled) <input checked="" type="checkbox"/> Level 5 (nested corners sampled)	
Date (mm/dd/yyyy):	8/3/2015
End date (if > 1 day):	/ /
Party:	Role**
C. Minney	Plot leader
D. Sweet	Bot. Asst.
R. Eagle-Malone	Woody Tech
E. Knauss	Woody Tech
** Roles: Co-leader, Asst. Guide, Observer, Taxonomist, etc.	
PLOT NOT SAMPLED: <input type="checkbox"/> Other	
<input type="checkbox"/> Fern, water <input type="checkbox"/> Paved <input type="checkbox"/> Slope <input type="checkbox"/> Safety	
SAMPLING QUALITY*	
Effort Level:	subjective evaluation of how much effort put into sampling. Hurried plots may still provide good data
<input checked="" type="checkbox"/> Very thorough	
<input type="checkbox"/> Accurate	
<input type="checkbox"/> Hurried	
TAXONOMIC ACCURACY	
	high modera. low not simpl
vascular:	<input checked="" type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> n/a
bryo:	<input type="checkbox"/> <input checked="" type="checkbox"/> <input type="checkbox"/>
lichen:	<input type="checkbox"/> <input type="checkbox"/> <input checked="" type="checkbox"/>
TAXONOMIC STANDARD	
Authority:	G&C Pub Date: 1998

Minimum required fields in Bold and Underlined

LOCATION	
State:	OH County: Cuyahoga
Quadrangle:	North Olmsted
Local Place Names:	End of Cleveland, Airport Runway
Landowner:	CMP
Data Confidentiality:	
Check one:	<input checked="" type="checkbox"/> Public data <input type="checkbox"/> Private Data
<input type="checkbox"/> Fuzz 100m	<input type="checkbox"/> Fuzz 250m <input type="checkbox"/> Fuzz 500m
Reason:	
If data not public why?	
Source of coordinates:	<input type="checkbox"/> MAP <input checked="" type="checkbox"/> GPS
Coordinate system:	Coord. Units
<input checked="" type="checkbox"/> Lat/Long <input type="checkbox"/> UTM <input type="checkbox"/> StatePlane	<input type="checkbox"/> deg <input type="checkbox"/> deg min <input type="checkbox"/> m <input type="checkbox"/> ft <input type="checkbox"/>
<input type="checkbox"/> Other (specify):	
Datum:	<input checked="" type="checkbox"/> NAD83/WGS84 <input type="checkbox"/> NAD27
GPS location in plot x=0 to 5, y=-1.0 to +1.0:	
x = 0 y = 0 (base of plot x=0, y=0)	
Latitude:	41.39828
Longitude:	81.87908
Coord. Accuracy:	X m <input type="checkbox"/> ft + - 3
GPS File Name:	1074A
Plot size for cover data:	.01 (hectares)
X-axis Bearing of plot:	17
Depth: (1-5):	4
Intensive modules:	2, 3, 8, 9 1 (EDIT IF MODIFIED)
Camera No.:	4
Photo No.:	4705
Plot placement:	<input checked="" type="checkbox"/> GRTS <input type="checkbox"/> Representative
<input type="checkbox"/> Random <input type="checkbox"/> Stratified Random <input type="checkbox"/> Transect component	
<input type="checkbox"/> Systematic (grid) <input type="checkbox"/> Capture specific feature <input type="checkbox"/> Other	

*Definitions and values in CM PCAP FOM v. 1.0 and CVS Field Guide

Layout: 1x1 (5m x 20m)

Location: ~500m NE of intersection of Ruple and Spafford. Park at pull off near intersection or Recycling Area where Ruple turns NW. We parked at the former because of limited space at the Recycling Center. Plot starts at end of guard rail, Plot is ~40m off Ruple. as you're traveling north along Ruple.

Rationale: GRTS

Veg Characterization: The canopy was dominated by Sugar Maple with an Elm and Fraxinus. Shrub layer had Rose, Sugar Maple, Cherry and Vitis. The herb layer was thick with graminoids near tree line and sparse in the understory.

Diagram key:

- Plot origin (0,0) point
- GPS location point
- photo taken, with direction
- location of permanent posts

NOTES: Include Layout (any unusual shape details), Location (directions and landscape content), Rationale (why here), and Veg Characterization (description of community, dominants, strata, BROWSE). Additional notes in space on back.

OVER

CLEVELAND METROPARKS Plant Community Assessment Program - Background Data Sheet

Project Name: **02R2015**

Project Label: _____ PCAP _____

Plot No.: **1074**

Page 2 of 2

MODIFIED NATURE RESERVE CLASS*

CODE (on separate form): **C03**

COMMUNITY NAME: **Sugar Maple Forest**

HOMOGENEITY

☒ Homogeneous ☐ Compositional trend across the plot

☒ Conspicuous inclusions ☐ Irregular/pattern mosaic

PCAP **Fit=** **Conf=**

Disturbances

type*	severity**	yrs ago	% of plot	description
Human	M	0	20	Bottles and trash in plot, most old
Natural				
Fire				
Cut	L	10+	1	Cut stump probably older than 5 years
Animal	L	0	100	Deer Browse
Other	M	?	20	Road slag dumped and some cut logs rolled through plot

**L=low, M=med low, M=med, MH=med high, H=high, VH=very high

Current Land Use: **CMP**

Former Land Use: _____

HYDROLOGIC REGIME*

☒ Upland (seldom flooded)

☐ Intermittently/seasonally saturated (seldom flooded)

☐ Permanently/Semipermanent, saturated (dry <1/yr, seldom flooded)

☐ Occasionally flooded (<1/yr)

☐ Temporarily flooded

(by default unless plot is a wetland)

SALINITY*

☐ Saltwater

☐ Brackish

☐ Fresh

☒ Upland (n/a)

DISTURBANCES

☐ Intermittently flooded

☐ Semipermanently flooded

☐ Permanently flooded

☐ Tidal/Seiche flooded daily

☐ Tidal/Seiche flooded monthly

☐ Tidal/Seiche flooded irregular (e.g. wind, storms)


☐ Unknown

Additional notes & diagrams: (Representativeness of plot to the stand, successional status, maturity, etc.)

The plot is small, the stand is young and mostly even-aged, 10-20% of the plot is not fully covered by trees due to the edge which undulates along the right side margin. This side has some weedy invaders, mainly rose, Festuca and Lolium? Moving from the tree line to the interior the slope descends gradually then dramatically (near vertical). Assume that is why browse is low. The wooded interior portion is sparse with areas of gradual, slight erosion and gravel/road slag/asphalt consisting mostly of weedy invaders and a lone population of Bloodroot. Down slope there is a population of Phragmites. Garlic Mustard establishing in plot.


EXAMPLES OF PERCENT OF AREA COVERED

The following graphic can be used for various data elements to convey "Amount" or "Quantity". NOTE: Within any given box, each quadrant contains the same total area covered, just different sized objects.



cover class	% cover	mid point
1	solitary or few	0.0001
2	0-1%	0.005
3	1-2%	0.015
4	2-5%	0.035
5	5-10%	0.075
6	10-25%	0.175
7	25-50%	0.375
8	50-75%	0.625
9	75-85%	0.850
10	85-100%	0.975

Nested Corners



BROWSE RATING NARRATIVE DESCRIPTION

LOW OR NONE: there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.



MEDIUM LOW values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

MEDIUM: browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.

MEDIUM HIGH values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.


HIGH: greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.

VERY HIGH values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.




EXAMPLES OF PERCENT OF AREA COVERED

The following graphic can be used for various data elements to carry "Amount" or "Quantity". NOTE: Within any given box, each quadrant contains the same bold area covered, just different sized objects.



cover class	% cover	midpoint
1	solitary or few	0.0001
2	0-1%	0.005
3	1-2%	0.015
4	2-5%	0.035
5	5-10%	0.075
6	10-25%	0.175
7	25-50%	0.375
8	50-75%	0.625
9	75-85%	0.850
10	85-100%	0.975

Nested Corners



BROWSE RATING NARRATIVE DESCRIPTION

LOW OR NONE: there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.


MEDIUM LOW values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

MEDIUM: browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.

MEDIUM HIGH values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.

HIGH: greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.

VERY HIGH values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.



Page 1 of 1

Page 1 of 1

Natural Resource Management FORM NR/2010-02a

Page 1 of 1

Project name: 0282015

Plot no.: 1074

1

9 3 2 2 2

CLEVELAND METROPARKS Plant Community Assessment Program Natural Woody Stem Data Sheet

Project Label: PCAP

Project Name: 02PR2015

Plot No.: 1079


Page: 1 of 1



Explain subsample (additional room on back):

mod #	species	c	voucher#	# stems 0-1.4m browed	% sub or super sample	# shrub clumps	size class (cm) woody stems > 1.4m	1 0-1	2 1-2.5	3 2.5-5	4 5-10	5 10-15	6 15-20	7 20-25	8 25-30	9 30-35	10 35-40	11 >40 (record each tree)
1	<i>Robinia pseudacacia</i>																	
1	<i>Vitis</i>		CKM307															
1	<i>Rosa pratincola</i>																	
1	<i>Quercus macrocarpa</i> sp.																	
1	<i>Acer spicatum</i>																	
1	<i>Rosa mollis</i>																	
1	<i>Staphylea trifolia</i>																	
1	<i>Prunus serotina</i>																	
1	<i>Ulmus rubra</i>																	
1	<i>Fraxinus americana</i>																	
1	<i>Acer negundo</i>																	

DBH Measurement Rules



Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.


Record using the tally system from 1 to 10




1



2



3



4



5

ASH CANOPY CONDITION

- 1. Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
- 2. Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
- 3. Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
- 4. >50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
- 5. Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.



A

B

C

D

E

ASH CANOPY BREAKUP CONDITION (for dead trees):

(If an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)

- A:** All main branches contain fine twigs (newly dead).
- B:** Over 50% of main branches have fine twigs.
- C:** Less than 50% of main branches have fine twigs.
- D:** Stem still standing and tertiary main branches present.
- E:** Central stem still standing.

Project Label: PCAP


Project Name: 06R2015

PLOT No.: 1079


Date: 8/3/15

Tree Module ID	Species	Dead	c	Voucher #	DBH (cm)	HT @ DBH	Ash condition	*Dead condition	# EAB exit holes	Epicormic present	Woodpecker holes
1	Fraxinus americana				17.9		2		0	0	1
2											
3											
4											
5											
6											
7											
8											
9											
10											
11											
12											
13											
14											
15											
16											
17											
18											
19											
20											
21											
22											
23											
24											
25											

ASH ONLY



*** Change intensive module numbers when necessary



Map all ash trees ≥ 10cm in each module using Tree ID number

- If Ash Condition scores 5 (dead) provide breakup score (A-E)
- Count EAB exit holes 1.25m² x 21.5m
- Woodpecker and epicormic marked present (1) or absent (0)



CLEVELAND METROPARKS Plant Community Assessment Program: Invasive Species Survey



Tier 1: Early detection/ Rapid response		Presence				GPS	Presence X: yes
		NE	SE	SW	NW		
Microstegium vimineum	Japanese stiltgrass						
Ranunculus ficaria	Lesser Celandine						
Cynanchum louiseae (vine)	Black Swallow-wort						
Butomus umbellatus (wetland)	Flowering Rush						
Heracleum mantegazzianum	Giant Hogweed						
Tier 2: Assess as Needed		# of Plants				comments	# of Plants 1: 1-10 2: 11-50. 3: 51-100 4: 101-1,000 5: >1,000
		NE	SE	SW	NW		
Acer platanoides	Norway Maple						
Ailanthus altissima	Tree of Heaven						
Lonicera japonica (vine)	Japanese Honeysuckle						
Lythrum salicaria (wetland)	Purple Loosestrife						
Aegopodium podagraria (G-cover)	Bishop's Goutweed						
Celastrus orbiculatus (vine)	Asian Bittersweet						
Torilis sp.	Hedgeparsley						
Conium maculatum	Poison Hemlock						
Rhamnus cathartica	Common Buckthorn (shrub)						
Berberis thunbergii	Japanese Barberry (shrub)						
Alnus glutinosa	European Alder						
Dipsacus laciniatus	Cut-leaf Teasel						
Elaeagnus umbellata	Autumn Olive (shrub)						
Lonicera maackii	Amur Honeysuckle (shrub)						
Euonymus fortunei	Wintercreeper						
Tier 3: Presence is of Interest		# of Plants				comments	# of Plants 1: 1-10 2: 11-50. 3: 51-100 4: 101-1,000 5: >1,000
		NE	SE	SW	NW		
Convallaria majalis (G-cover)	Lily of the Valley						
Coronilla varia (G-cover)	Crown Vetch						
Eleutherococcus pentaphyllus	Five-leaf Aralia (shrub)						
Pachysandra terminalis (G-cover)	Japanese Pachysandra						
Philadelphus coronarius	Mock Orange (shrub)						
Pulmonaria officinalis (G-cover)	Lungwort						
Rubus phoenicolasius	Wineberry						
Iris pseudacorus (wetland)	Yellow Flag Iris						
Ornithogalum umbellatum	Star of Bethlehem						
Viburnum opulus var. opulus	European Cranberry (shrub)						
Viburnum plicatum	Doublefile Viburnum (shrub)						
Tier 4: Widespread and abundant		Presence				comments	# of Plants 1: 1-10 2: 11-50. 3: 51-100 4: 101-1,000 5: >1,000
		NE	SE	SW	NW		
Alliaria petiolata	Garlic Mustard						
Ligustrum vulgare	Common Privet (shrub)						
L. morrowii, L. tatarica	Bush Honeysuckles (shrub)						
Phalaris arundinacea	Reed Canarygrass						
Phragmites australis (wetland)	Phragmites						
Polygonum cuspidatum	Japanese Knotweed						
Frangula alnus	Glossy Buckthorn (shrub)						
Rosa multiflora	Multiflora Rose (shrub)						
Typha angustifolia, T. x.glauca	Cattails (wetland)						
Cirsium arvense	Canada thistle						
Dipsacus fullonum	Common Teasel						
Hesperis matronalis	Dame's Rocket						
Vinca minor (G-cover)	Periwinkle						

Note: For Ground-cover plants record "stem #" but in comment field describe # of colonies and patch size (S,M, L)

CLEVELAND METROPARKS Plant Community Assessment Program Forest Pest and Pathogens Data Sheet



Project Label

PCAP

Project Name

028205

Plot No.

1034

Page

1

of

1

mod #	species	voucher#	# shrub clumps	size class (cm) woody stems >1m										
				1 0<1	2 1-2.5	3 2.5-5	4 5-10	5 10-15	6 15-20	7 20-25	8 25-30	9 30-35	10 35-40	11 >40 (record each tree)
1	None present													
2														
3														
4														
5														
6														
7														
8														
9														
10														

* IF EVIDENCE OF PEST OR PATHOGEN RECORD TOTAL SPECIES POPULATION IN THE PLOT EVEN THE NOT INFECTED

Strata	# of stem infected	Severity (H,M, or L)
Tree (size class 3 or above)		
Shrub (size class 2 or below including shrub clumps)		

* Write None Present if no evidence:

None

Beech (Fungus)

SRE 10-21-15

None

Asian Longhorned Beetle

Hemlock (HWA)

Other Pest or Pathogen

Walnut (Thousand Canker)

Severity

High = more than 50% of leaf/needle cover exhibiting symptoms

Medium = Less than 50% of leaf/needle cover exhibiting symptoms

Low = Only a few leaves or branches are exhibiting symptoms

COVER BY STRATA

STRATUM	GENERAL FORM
Tree (generally >5 m)	Tree (overstory), very tall shrubs*, liana, epiphyte)
Shrub (generally 0.5 to 5 m)	Tree (sapling), shrub, liana, epiphyte)
Herb (Field)	Herb, dwarf-shrub**, tree (seedling***)
Floating	Floating
Aquatic (submerged)	Submerged

*Very tall shrubs are sometimes included in the tree stratum
 **Can also include seedlings of shrubs, i.e. all shrubs <0.5m
 ***Tree seedlings are often defined as up to 1.4 m height or as <2.5 cm DBH in which case they would span the herb and shrub layers.





FIGURE 3-20.—Generalized section of Upper Devonian, Mississippian, and Lower Pennsylvanian formations in northeastern Ohio. Asterisks indicate units that are fossiliferous. This composite section represents about 400 meters of rock exposed across the area. The section is not to scale, but the thicknesses indicated are proportional. The term "Waverly" is used in the older literature to refer to Mississippian rocks in Ohio. Some geologists use the European term "Carboniferous," which encompasses the Mississippian and Pennsylvanian Periods of the U.S. Many units have been named within the Cuyahoga Formation, but most units are local and cannot be traced over great distances. The Black Hand Member is a spectacular massive sandstone that is fairly widespread but discontinuous. See Hyde (1953), Hoover (1960), and Collins (1979) for more information on Mississippian rocks in Ohio. See figure 3-18 for explanation of rock types.

SOIL PIT DESCRIPTION: Excavate 20 cm plug with shovel. Describe using Munsell chart, visual exam, texture, and odor

SOIL SAMPLES Standard procedure: collect a soil sample of the top 10 cm of soil from center of each intensive module and composite the sample

Soil pit module # (one per entire plot)

5 cm	matrix color
	mottle color
	%mottle
	oxid roots
	texture*
	redox features**
	hydr. cond.***
20 cm	matrix color
	mottle color
	%mottle
	oxid roots
	texture*
	redox features**
	hydr. cond.***

* refer to texture classes on reverse side
 ** e.g. hydrogen sulfide odor, gleying, etc.
 *** Circle one:
 I=indurated S=saturated M=moist D=dry
 Notes: include evidence of earthworms (worms, castings, middens)

(castings present)

Soil Collection Method	Horizon (A, B, C)
2,3,8,9 composited	A
Moist Soil Survey Laboratory	
Soil Series/Type	
Soil Series Source: Ohio Soil Survey	
Landform Type	
Depth to root layer	
Parent Material	
Drainage*	
<input type="checkbox"/> Excessively dr. <input type="checkbox"/> Somewhat excessively <input type="checkbox"/> Well drained <input type="checkbox"/> Moderately well dr. <input type="checkbox"/> Somewhat poorly dr. <input type="checkbox"/> Very poorly dr. <input type="checkbox"/> Impermeable surface	

SOIL DEPTH MEASUREMENT: Measure to the nearest 0.1 cm in center of intensive modules. If >30.5 cm, record as >30

	1 liter+ organic depth (cm)	2 liter depth (cm)	water depth (cm)	depth set soil (cm)
1	44.5	80.5	0	0

EARTH SURFACE & GROUND COVER			
Underlying Earth Surface*	Ground Cover		Percent
Gravel - 100%	Percent	(Each ≤ 100%)	
Historic		Coarse Woody Debris***	0
Mineral Soil		Fine Woody Debris****	14
Gravel-Cobble*	18	Litter	38
Boulder**	0	Duff (Ferm + Humus)	0
Bedrock	0	Bryophyte-Lichen	1
* Gravel-Cobble - 1/16-10"		Water	0
** Boulder = > 10 in		Bare Soil	17
*** > 5 cm in diameter		Road/Trail	0
**** < 5 cm in diameter		Other	0

TRAIL INFORMATION:	
record type and cover for each	
Type	%Cover
<input type="checkbox"/> All Purpose	
<input type="checkbox"/> Bridle	
<input type="checkbox"/> Hiking sanctioned	
<input type="checkbox"/> Boodleg unsanctioned	
<input type="checkbox"/> Gravel	
<input type="checkbox"/> Deer	

COVER BY STRATA
 estimate using midpoints of 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 49, 53, 57, 61, 65, 69, 73, 77, 81, 85, 89, 93, 97, 100

Strata	Height Range (m)	Total Cover (%)
Tree	5 - 9	88
Shrub	0.5 - 5	43
Herb	0 - 0.5	48
(Floating)*		
(Aquatic)*		


* rooted and floating or slightly emergent
 ** submerged, most plant mass below surface

SEE BACK OF PAGE FOR TYPICAL STRATA DESCRIPTIONS. STRATA CAN VARY BY COVER TYPE.

STAND SIZE	
<input type="checkbox"/> >600 x plot size	
<input type="checkbox"/> > 100 x plot size	
<input checked="" type="checkbox"/> 10-100 x plot size	
<input type="checkbox"/> 3-10 x plot size	
<input type="checkbox"/> 1-3 x plot size	
<input type="checkbox"/> < plot size	

PERCENT MOTTLES (USE CLASS CODES):

Class	Code	Criteria: % of Surface Area Covered
Few	f	< 2
Common	c	2 to < 20
Many	m	≥ 20



2%





20%

SOIL TEXTURE: Record the code for the soil texture of the 5 cm and 20 cm layers. To estimate texture, collect a soil sample from the appropriate layer and moisten it with water to the consistency of modeling clay/wet newspaper; the sample should be wet enough that all of the particles are saturated but excess water does not freely flow from the sample when squeezed. Attempt to roll the sample into a ball. If the soil will not stay in a ball and has a grainy texture, the texture is either sandy or coarse sandy. If the soil does form a ball, squeeze the sample between your fingers and attempt to form a self-supporting ribbon. Samples which form both a ball and a ribbon should be coded as clayey; samples which form a ball but not a ribbon should be coded as loamy.

- 0= Organic
- 1= Loamy
- 2= Clayey
- 3= Sandy
- 4= Coarse Sand
- 9= Not measured - make plot note


Geomorphic Component - Three-dimensional descriptors of parts of landforms or microfeatures that are best applied to areas. Unique descriptors are available for Hills, Terraces, Mountains, and Flat Plains; e.g., (for Hills) nose slope or NS.

Hills	Code	NASIS
Interfluvial	IF	IF
head slope	HS	HS
nose slope	NS	NS
side slope	SS	SS
base slope	BS	BS



Hillslope - Profile Position (Hillslope Position in PDP) - Two-dimensional descriptors of parts of line segments (i.e., slope position) along a transect that runs up and down the slope; e.g., backslope or BS. This is best applied to transects or points, not areas.

Position	Code
summit	SU
shoulder	SH
backslope	BS
footslope	FS
toeslope	TS



HYDROLOGIC REGIME Modified from Grossman et al 1998. (Frequency and duration of flooding.)

UPLAND: Not a wetland. Very rarely flooded.

INTERMITTENTLY/SEASONALLY SATURATED: Dry at least once per year. Surface water is seldom present, but substrate is saturated to surface for extended periods during the growing season.

PERMANENTLY/SEMI-PERMANENTLY SATURATED: Dry less than once per year. Surface water is seldom present, but substrate is saturated to surface for extended periods during the growing season. Equivalent to Cowardin's Saturated modifier.

OCCASIONALLY FLOODED: Surface water can be present for brief periods during growing season, but not in most years. Often characterizes flood-plain upper terraces.

TEMPORARILY FLOODED: Surface water present for brief periods during growing season, but water table usually lies well below soil surface. Often characterizes flood-plain levees and lower terraces. Equivalent to Cowardin's Temporary modifier.

INTERMITTENTLY FLOODED: Substrate is usually exposed, but surface water can be present for variable periods without detectable seasonal periodicity. Inundation is not predictable to a given season and is dependent upon highly localized rain storms. This modifier was developed for use in the arid West for water regimes of Playa lakes, intermittent streams, and dry washes but can be used in other parts of the U.S. where appropriate. This modifier can be applied to both wetland and non-wetland situations. Equivalent to Cowardin's Intermittently Flooded modifier.

SEMI-PERMANENTLY FLOODED (exposed <1/year): Surface water persists throughout the growing season in most years. Land surface is normally saturated when water level drops below soil surface. Includes Cowardin's Intermittently Exposed and Semipermanently Flooded modifiers.

PERMANENTLY FLOODED: Water covers the land surface at all times of the year in all years. Equivalent to Cowardin's "permanently flooded".

UNKNOWN: The hydrologic regime cannot be determined from the available information.