

CLEVELAND METROPARKS Plant Community Assessment Program: Quality Control Form



Project Label: PCAP

Plot No: 1013

Date Sampled: 07/01

Lead: LANCE

Comment required if item answer is NO

| | | | |
|--|--------------------------------------|---|--|
| Parking/Access outside of Park Boundaries: | <input checked="" type="radio"/> Y | <input type="radio"/> N | If yes, write details in Comments section below |
| Field journals completed | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| Site sketch made on 1:3000 map? | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| Check cover page | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| X-axis Bearing of plot recorded | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| GPS coords. Recorded | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| North direction recorded | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| Photographs taken? | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| Relocated Pins Mapped | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| Plot No., Date agreement on all pages? | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| Header data completed all pages? | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| Cover classes recorded in all Intensive modules | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| Browse Level By Species | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| Woody stem quality control check | <input checked="" type="radio"/> Y | <input type="radio"/> N | Check every line and cross check with the Tree Cover Sheet |
| Invasive plant quality control check | <input type="radio"/> Y | <input type="radio"/> N | N/A |
| Ash trees mapped | <input type="radio"/> Y | <input type="radio"/> N | N/A |
| Completed Forest Pest/Pathogen Datasheet | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| Cover by Strata? (confirm cover type) | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| Soil samples collected with matching plot #. | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| Cross check 2010 information | <input checked="" type="radio"/> Y | <input type="radio"/> N | Highlight any changes from 2010 information |
| Vouchers labeled on datasheet with initials and number | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| Vouchers labeled on collection bag | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| Pink flags removed | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| Data sheet QA before leaving site? | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| Common equipment returned to tub. | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| Data sheets scanned? | <input checked="" type="radio"/> Yes | <input type="radio"/> No | Enter date to left 7/6/15 |
| Final data sheets scanned? | <input type="radio"/> Yes | <input type="radio"/> No | Enter date to left |
| Buffer Widths measured? | <input type="radio"/> Y | <input type="radio"/> N | |
| Web Soil Survey | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| Voucher Location | <input checked="" type="radio"/> Y | <input type="radio"/> N | |
| (# vouchers collected) | Press (#) | | Enter number to left |
| | Drier | <input type="radio"/> Y <input type="radio"/> N | |
| | Identified | <input type="radio"/> Y <input type="radio"/> N | |
| | Mounted | <input type="radio"/> Y <input type="radio"/> N | |
| | Thrown away | <input type="radio"/> Y <input type="radio"/> N | |

GRTS point verification: Is plot sampleable?

| | |
|---|---|
| <input checked="" type="checkbox"/> Yes | Original GRTS point is sampleable |
| <input type="checkbox"/> No | Original GRTS point lands in a non-sampleable area (fill in category below) |
| | <input type="checkbox"/> Point falls in a water (i.e. river, lake) |
| | <input type="checkbox"/> Managed mowed area (i.e. golf course, picnic area, right-of-way) |
| | <input type="checkbox"/> Paved area (i.e. parkinglot, road) |
| | <input type="checkbox"/> Unsafe to sample (i.e. steep slope) |
| | <input type="checkbox"/> Other |

Additional Comments:

2015 Collect Sam's soils

PARK ON SIDE OF VALLEY PKWY.

Q

Q

8/1/14

1

1

1

1

1

1

1

1

1

1

1

| GENERAL INFORMATION | | LOCATION | |
|--|---|-------------------------|--|
| Project Label: PCAP | State: OH | County: Cuyahoga | |
| Project Name: QAB/2015 | Quadrangle: | | |
| Plot Name: Bay Band Plot | Local Place Names: Home at Valley Pkwy. 1521 Valley Pkwy. | | |
| Plot No.: 1013 | Landowner: CMR | | |
| <input type="checkbox"/> Level 4 (no nested corners sampled) | Data Confidentiality: | | |
| <input checked="" type="checkbox"/> Level 5 (nested corners sampled) | Check one: <input checked="" type="checkbox"/> Public data <input type="checkbox"/> Private Data | | |
| Date (mm/dd/yyyy): 07/01/2015 | <input type="checkbox"/> Fuzz 100m <input type="checkbox"/> Fuzz 250m <input type="checkbox"/> Fuzz 500m | | |
| End date (if > 1 day): 1/1 | Reasons: | | |
| Party: A. Lance | If data not public why? | | |
| Role: Plot leader | Source of coordinates: <input type="checkbox"/> MAP <input checked="" type="checkbox"/> GPS | | |
| M. Goffey | Coordinate system: Coord. Units | | |
| T. Cochran | <input checked="" type="checkbox"/> Lat/Long <input type="checkbox"/> UTM <input type="checkbox"/> StatePlane <input type="checkbox"/> deg <input type="checkbox"/> deg min | | |
| E. Krauss | <input type="checkbox"/> Other (specify): m <input type="checkbox"/> ft | | |
| | Datum: <input checked="" type="checkbox"/> NAD83/WGS84 <input type="checkbox"/> NAD27 | | |
| Plot NOT SAMPLED: <input type="checkbox"/> Other | GPS location in plot x=0 to 5, y=1.0, +1): | | |
| <input type="checkbox"/> Perm. water <input type="checkbox"/> Paved <input type="checkbox"/> Slope <input type="checkbox"/> Safety | x = 0 y = 0 (base of plot x=0, y=0) | | |
| SAMPLING QUALITY* | Latitude: 41.30081 | | |
| Effort Level: <input checked="" type="checkbox"/> Very thorough | Longitude: 81.69030 | | |
| <input type="checkbox"/> Accurate | Coord. Accuracy: m <input type="checkbox"/> ft | | |
| <input type="checkbox"/> Hurried | GPS File Name: 1013A | | |
| subjective evaluation of how much effort put into sampling. Hurried plots may still provide good data. | Plot size for cover data: .05 (hectares) | | |
| | X-axis Bearing of plot: [98] ° | | |
| TAXONOMIC ACCURACY | Depth: (1-5): 4 | | |
| <input type="checkbox"/> high <input type="checkbox"/> modera. <input type="checkbox"/> low <input type="checkbox"/> not simpl | Intensive modules: 2-3-8-9 1-2-3-4 (DIT IF MODIFIED) | | |
| vascul. <input checked="" type="checkbox"/> | Camera No.: 3 | | |
| bryo <input type="checkbox"/> | Photo Nos.: 0112 | | |
| lichen <input type="checkbox"/> | Plot placement: <input checked="" type="checkbox"/> GRTS <input type="checkbox"/> Representative | | |
| TAXONOMIC STANDARD | <input type="checkbox"/> Random <input type="checkbox"/> Stratified Random <input type="checkbox"/> Transect component | | |
| Authority: G&C Pub Date: 1998 | <input type="checkbox"/> Systematic (grid) <input type="checkbox"/> Capture specific feature <input type="checkbox"/> Other | | |

* Definitions and values in CM PCAP FOM v. 1.0 and CVS Field Guide


Layout → 1x5

Location → Plot is located next to Valley Parkway in the front yard of a private home.

Rationale → GRTS; PCAP re-sample

Veg. Characteristics → Diversity of trees in the canopy include beech, sugar maple, bitternut hickory, shagbark hickory, American elm. Sparse shrub layer.

Herb layer is also sparse with Prenanthes, Jack-in-the-pulpit, and several Sedges dominating.




NOTES: Include Layout (any unusual shape details), Location (directions and landscape content), Rationale (why here), and Veg Characterization (description of community, dominants, strata, BROWSE). Additional notes in space on back.

OVER

| CLEVELAND METROPARKS Plant Community Assessment Program - Background Data Sheet | | | |
|--|---------------|-------------------------------|----------------------------|
| Project Label: <u>PCAP</u> | | Project Name: <u>02Br2015</u> | |
| Plot No.: <u>1013</u> | | Page 2 of 2 | |
| MODIFIED NATURESERVE CLASS* | | | |
| CODE (on separate form): <u>C-02</u> Fit= Conf= | | | |
| COMMUNITY NAME: <u>Beech-Maple Forest</u> | | | |
| HOMOGENEITY | | | |
| <input checked="" type="checkbox"/> Homogeneous <input type="checkbox"/> Compositional trend across the plot | | | |
| <input checked="" type="checkbox"/> Conspicuous inclusions <input type="checkbox"/> Irregular/pattern mosaic | | | |
| DISTURBANCES | | | |
| type* | severity** | yrs ago | % of plot description |
| Human | M | 0 | 100% trash |
| Natural | | | |
| Fire | | | |
| Get | MH | 0 | 100% browse |
| Animal | MH | 0 | 100% browse |
| Other | | | |
| **L=low, ML=med low, M=med, MH=med high, H=high, VH=very high | | | |
| Current Land Use: <u>PARK (CONNECTS Br with MS)</u> | | | |
| Former Land Use: <u>UNKNOWN</u> | | | |
| HYDROLOGIC REGIME* | | | |
| <input checked="" type="checkbox"/> Upland (seldom flooded) <input type="checkbox"/> Intermittently flooded | | | |
| <input type="checkbox"/> Intermittently/seasonally saturated (seldom flooded) <input type="checkbox"/> Semipermanently flooded | | | |
| <input type="checkbox"/> Permanently/Semipermanent, saturated (dry <1/yr, seldom flooded) <input type="checkbox"/> Permanently flooded | | | |
| <input type="checkbox"/> Occasionally flooded (<1/yr) <input type="checkbox"/> Tidal/Seiche flooded daily | | | |
| <input type="checkbox"/> Temporarity flooded (e.g. wind, storms) <input type="checkbox"/> Tidal/Seiche flooded monthly | | | |
| <input type="checkbox"/> Unknown <input type="checkbox"/> Tidal/Seiche flooded irregular | | | |
| (by default unless plot is a wetland) | | | |
| SALINITY* | | | |
| <input type="checkbox"/> Saltwater | | | |
| <input type="checkbox"/> Brackish | | | |
| <input type="checkbox"/> Fresh | | | |
| <input checked="" type="checkbox"/> Upland (n/a) | | | |
| Additional notes & diagrams: (Representativeness of plot to the stand, successional status, maturity, etc.) | | | |
| <div style="font-size: 2em; color: blue; transform: rotate(-90deg); position: absolute; left: 50%; top: 50%;"> Small creek flows through the back of mod 2. </div> | | | |




EXAMPLES OF PERCENT OF AREA COVERED

The tabular graphic can be used in various data elements to convey "Amount" or "Quality". NOTE: Within any given box, each quadrant contains the same total area covered, just different sized objects.



| cover class | % cover | midpoint |
|-------------|-----------------|----------|
| 1 | solitary or few | 0.0001 |
| 2 | 0-1% | 0.005 |
| 3 | 1-2% | 0.015 |
| 4 | 2-5% | 0.035 |
| 5 | 5-10% | 0.075 |
| 6 | 10-25% | 0.175 |
| 7 | 25-50% | 0.375 |
| 8 | 50-75% | 0.625 |
| 9 | 75-95% | 0.850 |
| 10 | 95-100% | 0.975 |

Nested Corners



BROWSE RATING NARRATIVE DESCRIPTION

LOW OR NONE: there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.

MEDIUM LOW values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

MEDIUM: browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.


MEDIUM HIGH values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.

HIGH: greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.


VERY HIGH values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.

EXAMPLES OF PERCENT OF AREA COVERED

The following graphic can be used for various data elements to convey "Amount" or "Quantity". NOTE: Within any given box, each quadrant contains the same total area covered, just different sized objects.



| cover class | % cover | mid point |
|-------------|-----------------|-----------|
| 1 | solitary or few | 0.0001 |
| 2 | 0-1% | 0.005 |
| 3 | 1-2% | 0.015 |
| 4 | 2-5% | 0.035 |
| 5 | 5-10% | 0.075 |
| 6 | 10-25% | 0.175 |
| 7 | 25-50% | 0.375 |
| 8 | 50-75% | 0.625 |
| 9 | 75-85% | 0.850 |
| 10 | 95-100% | 0.975 |



BROWSE RATING NARRATIVE DESCRIPTION

LOW OR NONE: there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.


MEDIUM LOW values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

MEDIUM: browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.

MEDIUM HIGH values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.

HIGH: greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.

VERY HIGH values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.



Page 1 of 1Plot no.: 1013[illegible]

CLEVELAND METROPARKS Plant Community Assessment Program Natural Woody Stem Data Sheet

Project Label: PCAP

Project Name: 02B12015

Plot No.: 1013

Page: 1 of 2



Explain subsample (additional room on back):


| mod # | species | c | voucher# | # stems 0-1.4m browed | % sub or super sample | # shrub clumps | size class (cm) woody stems > 1.4m | | | | | | | | | | |
|-------|--|---|----------|-----------------------------|-----------------------------|----------------------|------------------------------------|------------|------------|-----------|---------------|---------------|---------------|---------------|---------------|----------------|------------------------------|
| | | | | | | | 1 0-1 | 2 1-2.5 | 3 2.5-5 | 4 5-10 | 5 10 - <15 | 6 15 - <20 | 7 20 - <25 | 8 25 - <30 | 9 30 - <35 | 10 35 - <40 | 11 >40 (record each tree) |
| 1 | Fagus grandifolia | | | 1 | | | | | | | | | | | | | |
| 1 | Ulmus americana | | | | | | | | | | | | | | | | |
| 1 | Corya cordiformis | | | | | | | | | | | | | | | | |
| 1 | Acer saccharum | | | | | | | | | | | | | | | | |
| 1 | Corya glabra | | | | | | | | | | | | | | | | |
| 1 | Parthenocissus quinquefolia | | | 2 | | | | | | | | | | | | | |
| 2 | Corya ovata | | | | | | | | | | | | | | | | |
| 2 | Acer saccharum | | | | | | | | | | | | | | | | |
| 2 | standing dead | | | | | | | | | | | | | | | | |
| 2 | Corya cordiformis | | | | | | | | | | | | | | | | |
| 2 | Acer saccharum | | | 1 | | | | | | | | | | | | | |
| 2 | Fagus grandifolia | | | | | | | | | | | | | | | | |
| 2 | Ulmus americana | | | | | | | | | | | | | | | | |
| 2 | Rosa multiflora | | | 2 | | | | | | | | | | | | | |
| 2 | Fraxinus sp. seedling | | | 1 | | | | | | | | | | | | | |
| 3 | Ulmus americana | | | | | | | | | | | | | | | | |
| 3 | Fagus grandifolia | | | 1 | | | | | | | | | | | | | |
| 3 | Corya cordiformis | | | | | | | | | | | | | | | | |
| 3 | Acer saccharum | | | | | | | | | | | | | | | | |
| 3 | Corya ovata | | | | | | | | | | | | | | | | |
| 3 | standing dead | | | | | | | | | | | | | | | | |
| 3 | Fraxinus sp. seedling | | | 1 | | | | | | | | | | | | | |
| 3 | Rosa multiflora | | | 2 | | | | | | | | | | | | | |
| 4 | standing dead | | | | | | | | | | | | | | | | |

Is this an *Carya* (2015)
or an *Ash*?
(2010)

Is this an *Corya* (2015)
or an *Ash*?
(2010)

Previously taken a 10-15
a meter

DBH Measurement Rules



Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.


Record using the tally system from 1 to 10



1




2



3



4



5

ASH CANOPY CONDITION

- 1. Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
- 2. Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
- 3. Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
- 4. >50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
- 5. Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.



A

B

C

D

E

ASH CANOPY BREAKUP CONDITION (for dead trees):

(If an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)


- A:** All main branches contain fine twigs (newly dead).
- B:** Over 50% of main branches have fine twigs.
- C:** Less than 50% of main branches have fine twigs.
- D:** Stem still standing and tertiary main branches present.
- E:** Central stem still standing.

✓

Specialized Marketing

3aCM PCAP Natural Woody Stem Data Sheet ver 2.0.xls last revised 5/29/2012 jim

DBH Measurement Rules



Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.

Record using the tally system from 1 to 10



1



2



3



4



5

ASH CANOPY CONDITION

1. **Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
2. **Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
3. **Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
4. **>50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
5. **Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.



A

B

C

D

E

ASH CANOPY BREAKUP CONDITION (for dead trees):

(If an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)


- A: All main branches contain fine twigs (newly dead).
- B: Over 50% of main branches have fine twigs.
- C: Less than 50% of main branches have fine twigs.
- D: Stem still standing and tertiary main branches present.
- E: Central stem still standing.

| Module ID | Tree ID | Species | Dead | Voucher # | DBH (cm) | HT @ DBH | Ash condition | Dead condition | # Exit holes | Epiloric present | Woodpecker holes |
|-----------|---------|---------|------|-----------|----------|----------|---------------|----------------|--------------|------------------|------------------|
| 1 | | DONE | | | | | | | | | |
| 2 | | | | | | | | | | | |
| 3 | | | | | | | | | | | |
| 4 | | | | | | | | | | | |
| 5 | | | | | | | | | | | |
| 6 | | | | | | | | | | | |
| 7 | | | | | | | | | | | |
| 8 | | | | | | | | | | | |
| 9 | | | | | | | | | | | |
| 10 | | | | | | | | | | | |
| 11 | | | | | | | | | | | |
| 12 | | | | | | | | | | | |
| 13 | | | | | | | | | | | |
| 14 | | | | | | | | | | | |
| 15 | | | | | | | | | | | |
| 16 | | | | | | | | | | | |
| 17 | | | | | | | | | | | |
| 18 | | | | | | | | | | | |
| 19 | | | | | | | | | | | |
| 20 | | | | | | | | | | | |
| 21 | | | | | | | | | | | |
| 22 | | | | | | | | | | | |
| 23 | | | | | | | | | | | |
| 24 | | | | | | | | | | | |
| 25 | | | | | | | | | | | |

* If Ash Condition scores 5 (dead) provide breakup score (A-E)
Count EAB exit holes 1.25m² x 21.5m
Woodpecker and epicormic marked present (1) or absent (0)

ASH ONLY


Baseline



*** Change intensive module numbers when necessary



Map all ash trees ≥ 10cm in each module using Tree ID number



CLEVELAND METROPARKS Plant Community Assessment Program: Invasive Species Survey



| Tier 1: Early detection/ Rapid response | | Presence | | | | GPS |
|---|-----------------------------|-------------|----|----|----|----------|
| | | NE | SE | SW | NW | |
| Microstegium vimineum | Japanese stiltgrass | | | | | |
| Ranunculus ficaria | Lesser Celandine | | | | | |
| Cynanchum louiseae (vine) | Black Swallow-wort | | | | | |
| Butomus umbellatus (wetland) | Flowering Rush | | | | | |
| Heracleum mantegazzianum | Giant Hogweed | | | | | |
| Tier 2: Assess as Needed | | # of Plants | | | | comments |
| | | NE | SE | SW | NW | |
| Acer platanoides | Norway Maple | | | | | |
| Ailanthus altissima | Tree of Heaven | | | | | |
| Lonicera japonica (vine) | Japanese Honeysuckle | | | | | |
| Lythrum salicaria (wetland) | Purple Loosestrife | | 4 | | | |
| Aegopodium podagraria (G-cover) | Bishop's Goutweed | | | | | |
| Celastrus orbiculatus (vine) | Asian Bittersweet | | | | | |
| Torilis sp. | Hedgeparsley | | | | | |
| Conium maculatum | Poison Hemlock | | | | | |
| Rhamnus cathartica | Common Buckthorn (shrub) | | | | | |
| Berberis thunbergii | Japanese Barberry (shrub) | | | | | |
| Alnus glutinosa | European Alder | | | | | |
| Dipsacus laciniatus | Cut-leaf Teasel | | | | | |
| Elaeagnus umbellata | Autumn Olive (shrub) | | | | | |
| Lonicera maackii | Amur Honeysuckle (shrub) | | | | | |
| Euonymus fortunei | Wintercreeper | | | | | |
| Tier 3: Presence is of Interest | | # of Plants | | | | comments |
| | | NE | SE | SW | NW | |
| Convallaria majalis (G-cover) | Lily of the Valley | | | | | |
| Coronilla varia (G-cover) | Crown Vetch | | | | | |
| Eleutherococcus pentaphyllus | Five-leaf Aralia (shrub) | | | | | |
| Pachysandra terminalis (G-cover) | Japanese Pachysandra | | | | | |
| Philadelphus coronarius | Mock Orange (shrub) | | | | | |
| Pulmonaria officinalis (G-cover) | Lungwort | | | | | |
| Rubus phoenicolasius | Wineberry | | | | | |
| Iris pseudacorus (wetland) | Yellow Flag Iris | | | | | |
| Ornithogalum umbellatum | Star of Bethlehem | | | | | |
| Viburnum opulus var. opulus | European Cranberry (shrub) | | | | | |
| Viburnum plicatum | Doublefile Viburnum (shrub) | | | | | |
| Tier 4: Widespread and abundant | | Presence | | | | comments |
| | | NE | SE | SW | NW | |
| Alliaria petiolata | Garlic Mustard | | | | | |
| Ligustrum vulgare | Common Privet (shrub) | | | | | |
| L. morrowii, L. tatarica | Bush Honeysuckles (shrub) | | | | | |
| Phalaris arundinacea | Reed Canarygrass | | | | | |
| Phragmites australis (wetland) | Phragmites | | | | | |
| Polygonum cuspidatum | Japanese Knotweed | | | | | |
| Frangula alnus | Glossy Buckthorn (shrub) | | | | | |
| Rosa multiflora | Multiflora Rose (shrub) | | | | | |
| Typha angustifolia, T. x. glauca | Cattails (wetland) | | | | | |
| Cirsium arvense | Canada thistle | | | | | |
| Dipsacus fullonum | Common Teasel | | | | | |
| Hesperis matronalis | Dame's Rocket | | | | | |
| Vinca minor (G-cover) | Periwinkle | | | | | |

Presence
X: yes

of Plants
1: 1-10
2: 11-50.
3: 51-100
4: 101-1,000
5: >1,000

of Plants
1: 1-10
2: 11-50.
3: 51-100
4: 101-1,000
5: >1,000

of Plants
1: 1-10
2: 11-50.
3: 51-100
4: 101-1,000
5: >1,000

Note: For Ground-cover plants record "stem #" but in comment field describe # of colonies and patch size (S,M, L)

Project Label: PCAP

Project Name: 02567015

Plot No.: 1013

Page: 1 of 1

Explain subsample (additional room on back):

| mod # | species | voucher# | % sub or super sample | # shrub clumps | size class (cm) woody stems > 1m | | | | | | | | | | |
|-------|---------|----------|-----------------------|----------------|----------------------------------|-------------|-------------|------------|---------------|---------------|---------------|---------------|---------------|----------------|------------------------------|
| | | | | | 1 0-<1 | 2 1-<2.5 | 3 2.5-<5 | 4 5-<10 | 5 10 - <15 | 6 15 - <20 | 7 20 - <25 | 8 25 - <30 | 9 30 - <35 | 10 35 - <40 | 11 >40 (record each tree) |
| 1 | | | | | | | | | | | | | | | |
| 2 | | | | | | | | | | | | | | | |
| 3 | | | | | | | | | | | | | | | |
| 4 | | | | | | | | | | | | | | | |
| 5 | | | | | | | | | | | | | | | |
| 6 | | | | | | | | | | | | | | | |
| 7 | | | | | | | | | | | | | | | |
| 8 | | | | | | | | | | | | | | | |
| 9 | | | | | | | | | | | | | | | |
| 10 | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | |

NONE PRESENT

| Strata | Total % Cover |
|------------|---------------|
| Tree | |
| Shrub | |
| Herbaceous | |

| * Write None Present if no evidence: | |
|--------------------------------------|--------------------------------|
| NONE -Beech (Fungus) | -Asian Longhorned Beetle |
| -Hemlock (HWA) | -Other Forest Pest or Pathogen |
| -Walnut (Thousand Canker) | |

Slope 3 = maximum steepness that can be safely sampled -45°

| c.w.d. - count for pieces with minimum 1m length | | | | | |
|--|-------------------------------|-------------------------------|------------------------------|--------------------|--|
| c.w.d p-12 cm | c.w.d (12-40cm) | c.w.d >40 cm | microhab. interspers. | microhab. SLOPE | |
| depth 1 10x10cm (count) | depth 1 10x10cm (count) | depth 1 10x10cm (count) | depth 1 10x10cm (rank) | 10x10cm (rank) | |
| 18 | 5 | 0 | 2 | 1 | |
| 16 | 5 | 0 | 2 | 1 | |
| 22 | 1 | 0 | 2 | 1 | |
| 20 | 0 | 0 | 2 | 1 | |
| | | | | | |
| | | | | | |
| | | | | | |

| | | | | |
|----|----|----|----|----|
| 10 | 25 | 40 | 77 | 11 |
|----|----|----|----|----|

| | | | | | |
|----|----|-------|----|----|----|
| 21 | 11 | 0.077 | 11 | 11 | 11 |
|----|----|-------|----|----|----|

L

LFI is angle of plot to the horizon. TSI is angles formed by local slopes. For TSI measure angle from recorder eye to eye of person standing - 0 m away.

9

COVER BY STRATA

| STRATUM | GENERAL FORM |
|------------------------------|---|
| Tree (generally >5 m) | Tree (overstory), very tall shrubs*, liana, epiphyte) |
| Shrub (generally 0.5 to 5 m) | Tree (sapling), shrub, liana, epiphyte) |
| Herb (Field) | Herb, dwarf-shrub**, tree (seedling***) |
| Floating | Floating |
| Aquatic (submerged) | Submerged |

*Very tall shrubs are sometimes included in the tree stratum
 **Can also include seedlings of shrubs, i.e. all shrubs <0.5m
 ***Tree seedlings are often defined as up to 1.4 m height or as <2.5 cm DBH in which case they would span the herb and shrub layers.





FIGURE 3-20.—Generalized section of Upper Devonian, Mississippian, and Lower Pennsylvanian formations in northeastern Ohio. Asterisks indicate units that are fossiliferous. This composite section represents about 400 meters of rock exposed across the area. The section is not to scale, but the thicknesses indicated are proportional. The term "Waverly" is used in the older literature to refer to Mississippian rocks in Ohio. Some geologists use the European term "Carboniferous," which encompasses the Mississippian and Pennsylvanian Periods of the U.S. Many units have been named within the Cuyahoga Formation, but most units are local and cannot be traced over great distances. The Black Hand Member is a spectacular massive sandstone that is fairly widespread but discontinuous. See Hyde (1953), Hoover (1960), and Collins (1979) for more information on Mississippian rocks in Ohio. See Figure 3-18 for explanation of rock types.

Project label: PCAP

Project Name: 02B 2015

Plot No.: 1013

SOIL PIT DESCRIPTION: Excavate 20 cm plug with shovel. Describe using Munsell chart, visual exam, texture, and odor.

SOIL SAMPLES Standard procedure: collect a soil sample of the top 10 cm of soil from center of each intensive module and composite the sample

Soil pit module # _____ (one per entire plot)

| | | |
|-------|------------------|---------|
| 5 cm | matrix color | |
| | mottle color | |
| | %mottle | |
| | oxid roots | Y N |
| | texture* | |
| | redox features** | Y N |
| | hydr. cond.*** | I S M D |
| 20 cm | matrix color | |
| | mottle color | |
| | %mottle | |
| | oxid roots | Y N |
| | texture* | |
| | redox features** | Y N |
| | hydr. cond.*** | I S M D |

* refer to texture classes on reverse side
 ** e.g. hydrogen sulfide odor, gleying, etc.
 *** Circle one:
 I-indurated S-saturated M-moist D-dry
 Notes: include evidence of earthworms (worms, castings, middens)

1 - Worms present
 2 - Worms present
 3 - Worms present
 4 - Worms present

| | |
|---|-------------------|
| Soil Collection Method | Horizon (A, B, C) |
| 2.3.9 completed | A |
| Valid Soil Survey Information | |
| Soil Series/Type: | |
| Soil Series Source: | Ohio Soil Survey |
| Landform Type: | |
| Depth to root layer: | |
| Parent Material: | |
| DETAILED* | |
| <input type="checkbox"/> Excessively dr. <input type="checkbox"/> Somewhat excessively <input type="checkbox"/> Well drained <input type="checkbox"/> Moderately well dr. <input type="checkbox"/> Somewhat poorly dr. <input type="checkbox"/> Very poorly dr. <input type="checkbox"/> Impermeable surface | |

SOIL DEPTH MEASUREMENT: Measure to the nearest 0.1 cm in center of intensive modules. If >30.5 cm, record as >30

| 1 litter+ organic depth (cm) | 2 litter depth (cm) | water depth (cm) | depth sat soil (cm) |
|------------------------------|---------------------|------------------|---------------------|
| 1 2.4 | 2.4 | 0 | 0 |
| 2 3.0 | 3.0 | 0 | 0 |
| 3 2.0 | 2.0 | 0 | 0 |
| 4 2.3 | 2.3 | 0 | 0 |

| EARTH SURFACE & GROUND COVER | | | |
|------------------------------|------------------------|---------------|--|
| Underlying Earth Surface* | Ground Cover | percent | |
| Bare = 100% | percent | (Each ≤ 100%) | |
| Hiscod | Coarse Woody Debris*** | 15% | |
| Mineral Soil | Fine Woody Debris**** | 5% | |
| Gravel-Cobble* | Litter | 75% | |
| Boulder** | Duff (Ferm + Humus) | 1% | |
| Bedrock | Bryophyte-Lichen | 2% | |
| * Gravel-Cobble = 1/16-10" | Water | 10% | |
| ** Boulder = > 10 in | Bare Soil | | |
| *** > 5 cm in diameter | Dead/Tail | | |
| **** < 5 cm in diameter | Other | | |

| TRAIL INFORMATION: | |
|---|--------|
| record type and cover for each | |
| Type | %Cover |
| <input type="checkbox"/> All Purpose | |
| <input type="checkbox"/> Bridle | |
| <input type="checkbox"/> Hiking sanctioned | |
| <input type="checkbox"/> Bootleg unsanctioned | |
| <input type="checkbox"/> Gravel | |
| <input type="checkbox"/> Deer | |

| COVER BY STRATA | | | % |
|---|------------------|-----------------|---|
| estimate using midpoints of 5, 9, 13, 8, 13 | | | |
| Strata | Height Range (m) | Total Cover (%) | |
| Tree | 5 - | 93% | |
| Shrub | 0.5 - 5 | 13% | |
| Herb | 0 - 0.5 | 28% | |
| (Floating)* | - | | |
| (Aquatic)* | - | | |


* rooted and floating or slightly emergent
 ** submerged, most plant mass below surface

SEE BACK OF PAGE FOR TYPICAL STRATA DESCRIPTIONS. STRATA CAN VARY BY COVER TYPE


| STAND SIZE | |
|---|--|
| <input type="checkbox"/> > 600 x plot size | |
| <input type="checkbox"/> > 100 x plot size | |
| <input type="checkbox"/> 10-100 x plot size | |
| <input type="checkbox"/> 1-10 x plot size | |
| <input type="checkbox"/> 1-3 x plot size | |
| <input type="checkbox"/> < plot size | |

PERCENT MOTTLES (USE CLASS CODES):

| Class | Conv. | Code NASIS | Criteria: % of Surface Area Covered |
|--------|-------|------------|-------------------------------------|
| Few | f | # | < 2 |
| Common | c | # | 2 to < 20 |
| Many | m | # | ≥ 20 |



2%




20%

SOIL TEXTURE: Record the code for the soil texture of the 5 cm and 20 cm layers. To estimate texture, collect a soil sample from the appropriate layer and moisten it with water to the consistency of modeling clay/wet newspaper; the sample should be wet enough that all of the particles are saturated but excess water does not freely flow from the sample when squeezed. Attempt to roll the sample into a ball. If the soil will not stay in a ball and has a grainy texture, the texture is either sandy or coarse sandy. If the soil does form a ball, squeeze the sample between your fingers and attempt to form a self-supporting ribbon. Samples which form both a ball and a ribbon should be coded as clayey; samples which form a ball but not a ribbon should be coded as loamy.

- 0= Organic
- 1= Loamy
- 2= Clayey
- 3= Sandy
- 4= Coarse Sand
- 9= Not measured - make plot note


Geomorphic Component - Three-dimensional descriptors of parts of landforms or microfeatures that are best applied to areas. Unique descriptors are available for Hills, Terraces, Mountains, and Flat Plains; e.g., (for Hills) nose slope or NS.

| Hills | PDP | Code NASIS |
|--------------|-----|------------|
| interfluvial | IF | IF |
| head slope | HS | HS |
| nose slope | NS | NS |
| side slope | SS | SS |
| base slope | — | BS |



Hillslope - Profile Position (Hillslope Position in PDP) - Two-dimensional descriptors of parts of line segments (i.e., slope position) along a transect that runs up and down the slope; e.g., backslope or BS. This is best applied to transects or points, not areas.

| Position | Code |
|-----------|------|
| summit | SU |
| shoulder | SH |
| backslope | BS |
| footslope | FS |
| toeslope | TS |



HYDROLOGIC REGIME Modified from Grossman et al 1998. (Frequency and duration of flooding.)

UPLAND: Not a wetland. Very rarely flooded.

INTERMITTENTLY/SEASONALLY SATURATED: Dry at least once per year. Surface water is seldom present, but substrate is saturated to surface for extended periods during the growing season.

PERMANENTLY/SEMPERMANENTLY SATURATED: Dry less than once per year. Surface water is seldom present, but substrate is saturated to surface for extended periods during the growing season. Equivalent to Cowardin's Saturated modifier.

OCCASIONALLY FLOODED: Surface water can be present for brief periods during growing season, but not in most years. Often characterizes flood-plain upper terraces.

TEMPORARILY FLOODED: Surface water present for brief periods during growing season, but water table usually lies well below soil surface. Often characterizes flood-plain levees and lower terraces. Equivalent to Cowardin's Temporary modifier.

INTERMITTENTLY FLOODED: Substrate is usually exposed, but surface water can be present for variable periods without detectable seasonal periodicity. Inundation is not predictable to a given season and is dependent upon highly localized rain storms. This modifier was developed for use in the arid West for water regimes of Playa lakes, intermittent streams, and dry washes but can be used in other parts of the U.S. where appropriate. This modifier can be applied to both wetland and non-wetland situations. Equivalent to Cowardin's Intermittently Flooded modifier.

SEMPERMANENTLY FLOODED (exposed <1/year): Surface water persists throughout the growing season in most years. Land surface is normally saturated when water level drops below soil surface. Includes Cowardin's Intermittently Exposed and Semipermanently Flooded modifiers.

PERMANENTLY FLOODED: Water covers the land surface at all times of the year in all years. Equivalent to Cowardin's "permanently flooded".

UNKNOWN: The hydrologic regime cannot be determined from the available information.