CLEVELAND METROPARKS Plant Community Assessment Program: Quality Control Form Cleveland Metroparks

Project Label: PCAP **Plot No:** 1389 **Date Sampled:** 8/19/13 **Lead:** SJC

			Comment required if item answer is NO
Parking/Access outside of Park Boundaries:		Y (N)	If yes, write details in Comments section below
Field journals completed		(Y) N	
Site sketch made on 1:3000 map?		(Y) N	
Check cover page	X-axis Bearing of plot recorded	(Y) N	
	GPS coords. Recorded	(Y) N	
	North direction recorded	(Y) N	
	Photographs taken?	(Y) N	
Plot No., Date agreement on all pages?		(Y) N	
Header data completed all pages?		(Y) N	
Cover classes recorded in all Intensive modules		(Y) N	
Browse Level By Species		(Y) N	
Woody stem quality control check		(Y) N	
Invasive plant quality control check		(Y) N	
Ash trees mapped		(Y) (N)	na
Cover by Strata? (confirm cover type)		(Y) N	
Soil samples collected with matching plot #		(Y) N	
Vouchers labeled on datasheet with initials and number		(Y) N	
Vouchers labeled on collection bag		(Y) N	
Pink flags removed		(Y) N	
Data sheet QA before leaving site?		(Y) N	
Common equipment returned to tub.		(Y) N	
Data sheets scanned?		8/23/13	Enter date to left AB
Final data sheets scanned?			Enter date to left
Buffer Widths measured?		(Y) N	CL 6-28
Web Soil Survey		(Y) N	RSE 23Aug2013
Voucher Location	Refrigerator	Y N	
(# vouchers collected)	Press (#)		Enter number to left
SJC-233-34	Drier	Y N	
	Identified	Y N	
	Mounted	Y N	
	Thrown away	Y N	

GRTS point verification: Is plot sampleable?	
(Y) Yes	Original GRTS point is sampleable
☐ No	Original GRTS point lands in a non-sampleable area (fill in category below)
	☐ Point falls in a water (i.e. river, lake)
	☐ Managed mowed area (i.e. golf course, picnic area, right-of-way)
	☐ Paved area (i.e. parkinglot, road)
	☐ Unsafe to sample (i.e. steep slope)
	☐ Other

Additional Comments:

CLEVELAND METROPARKS Plant Community Assessment Program - Background Data Sheet

GENERAL INFORMATION

Project Label: PCAP

Project Name: 01Hi2013

Plot Name: Fantasy Forest I: The Dek u Tree

Plot No.: 1389

☐ Level 4 (no nested corners sampled)

■ Level 5 (nested corners sampled)

Date (mm/dd/yyyy): 08/19/2013

End date (if > 1 day): / /

Party	Role**
S. Catella	Plot leader
C. Lemmo	woody
C. Devono	woody

** Roles: Co-leader, Asst., Guide, Owner, Taxonomist, etc.

PLOT NOT SAMPLED: ☐ Other

☐ Perm. water ☐ Paved ☐ Slope ☐ Safety

SAMPLING QUALITY*

Effort Level:

☒ Very thorough

☐ Accurate

☐ Hurried

subjective evaluation of how much effort put into sampling. Hurried plots may still provide good data

TAXONOMIC ACCURACY

	high	modera.	low	not smpl
vascul.	☒			n/a
bryo			☒	
lichen				☒

TAXONOMIC STANDARD

Authority: G&C Pub Date: 1998

LOCATION

State: OH **County:** Medina

Quadrangle:

Local Place Names: State Rd. between East Dr. & Bellus

Landowner: CMP

Data Confidentiality:

Check one: ☒ Public data ☐ Private Data

☐ Fuzz 100m ☐ Fuzz 250m ☐ Fuzz 500m

Reason:

If data not public why?

Source of coordinates ☐ MAP ■ GPS

Coordinate system: **Coord. Units**

■ Lat/Long ☐ UTM ☐ StatePlane ■ deg ☐ deg min

☐ Other (*specify*) ■ m ☐ ft ☐ ___

Datum: ■ NAD83/WGS84 ☐ NAD27

GPS location in plot x=0 to 5, y=-1,0,+1):

x = 0 y = 0 (base of plot x=0, y=0)

Latitude: 41.27529

Longitude: 81.70518

Coord. Accuracy: ☐ m ☐ ft 100' +-

GPS File Name: 1389A

Plot size for cover data: .1 (hectares)

X-axis Bearing of plot: [177] °

Depth: (1-5): 4

Intensive modules: 2, 3, 8, 9 (EDIT IF MODIFIED)

Camera No.: C5

Photo Nos.: 2606

Plot placement: ☒ GRTS ☐ Representative

☐ Random ☐ Stratified Random ☐ Transect component

☐ Systematic (grid) ☐ Capture specific feature ☐ Other



NOTES: Include **Layout** (any unusual shape details), **Location** (directions and landscape content), **Rationale** (why here), and **Veg Characterization** (description of community, dominants, strata, BROWSE). Additional notes in space on back.

Layout: 2x5

Location: park on mowed burm on the E side of State Rd. about halfway between East Dr. & Bellus. Plot is ~50m east into the woods on the South side of the ravine. Runs perpendicular to the ravine.

Rationale: GRTS

Veg. Char. Canopy - dominated by Sugar Maple. Also had Quercus velutina & rubra, some huge Carya cordiformis and Ostrya, and some Tulip in the back with a Sassafras.

Shrub - 6-10 heavy with Acer Saccharum, some wide-canopied Ostrya, and Lindera & Berberis broke into the shrub layer occasionally.

Herb - for the most part depauperate. Mostly Carya cordiformis and Fraxinus seedlings. Noted a patch of Viburnum aceri folium, one Ribes sp., and scattered bunches of Berberis. Partheneocissus

OVER

Minimum required fields in Bold and Underlined

*Definitions and values in CM PCAP FOM v. 1.0 and CVS Field Guide

CLEVELAND METROPARKS Plant Community Assessment Program - Background Data Sheet

Project Label: PCAP Project Name: OHi2013 Plot No.: 1389

MODIFIED NATURESERVE CLASS*

CODE (on separate form): Fit=___ Conf=___

D

COMMUNITY NAME:

Mixed

DISTURBANCES

type*	severity**	yrs ago	% of plot	description
Human	M	0	100	herbicide use & trash
Natural				
Fire				
Cut				
Animal	M	0	100	deer browse
Other				

**L=low, ML=med low, M=med, MH=med high, H=high, VH=very high

Current Land Use: CMP

Former Land Use: UNK

HOMOGENEITY

- ☑ Homogeneous
- ☐ Compositional trend across the plot
- ☐ Conspicuous inclusions
- ☐ Irregular/pattern mosaic

HYDROLOGIC REGIME*

- ☒ Upland (seldom flooded)
- ☐ Intermittently/seasonally saturated (seldom flooded)
- ☐ Permanently/Semipermanent. saturated (dry <1/yr. seldom flooded)
- ☐ Occasionally flooded (<1/yr)
- ☐ Temporarily flooded
- ☐ Intermittently flooded
- ☐ Semipermanently flooded
- ☐ Permanently flooded
- ☐ Tidal/Seiche flooded daily
- ☐ Tidal/Seiche flooded monthly
- ☐ Tidal/Seiche flooded irregular (e.g. wind, storms)
- ☐ Unknown

SALINITY*

- ☐ Saltwater
- ☐ Brackish
- ☐ Fresh
- ☒ Upland (n/a)

(by default unless plot is a wetland)

Additional notes & diagrams: (Representativeness of plot to the stand, successional status, maturity, etc.)

The light gap in the corner of mod 6 was less than a frame's worth so I opted to call this plot homogenous. Herbicide use was not ~~appare~~ apparent but we found CWD with blue dye. Browse was not heavy on any one particular plant but nearly everything that could be reached was browsed at least once. Decided on mixed community type because of the absense of Beech in the shrubs & canopy. Couldve easily been a beech-maple because Sugar Maple was the dominant tree. Overall depauperate herb layer, 6-10 heavy with sugar in the shrub layer. Some of the BIGGEST astrya Ive ever seen!

CLEVELAND METROPARKS Plant Community Assessment Program Species Cover Data Sheet 2a Page 1 of 2

Project Label: PCAP Project name: 01Hi2013 Plot no.: 1389

Total modules: 10 Intensive modules: 4 Plot configuration: 2x5 Plot area (ha): .1

Cleveland Metroparks

Br = Browse Level. Use cover classes to describe amount of browse per species over entire plot

Strata - Cov. entire plot

Estimate for each intensive module:

	mod	corner	mod	corner	mod	corner	mod	corner	mod	corner	mod	corner	mod	corner	mod	corner	mod	corner
	2	4	2	2	3	4	3	2	8	4	8	2	9	4	9	2	R	R
	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov
%open water	1	0			1	0			1	0			1	0				
%unvegetated open water	1	0			1	0			1	0			1	0				
%unveg. ground (bare soil)	1	3			1	4			1	2			1	2				
%unveg. litter (bare litter)	1	9			1	9			1	10			1	10				

T	S	H	(F)	(A)	Br	Species	c	Voucher #	mod depth	mod cov	corner depth	corner cov	mod depth	mod cov	corner depth	corner cov	mod depth	mod cov	corner depth	corner cov	mod depth	mod cov	corner depth	corner cov	mod depth	mod cov	corner depth	corner cov	mod depth	mod cov	corner depth	corner cov
8	8	2			3	Acer saccharum			4	9	3		4	9	4		4	9	4		4	9	4									
6						Quercus velutina			4	5							4	7	2		4	3	4									
6		4			7	Carya cordiformis			3	5	4		3	4	2		4	3	4		4	8	4									
		2			2	Acer sp. (seedling)			4	2	2		2	2			2	2	2		1	2										
6	5	4				Ostrya virginiana			3	6	4		1	4					4	7	4	8	4									
	2	2			8	Lindera benzoin			2	2	2		1	4																		
		2			4	Parthenocissus quinquefolia			2	2	3		3	2					4	2	1	2										
2		2			6	Prunus serotina			2	2			1	2					3	2			3	2								
		2				Arisaema triphyllum			2	2	2		3	2			2	2	2		2	2	2									
		2			5	Actea alba			2	2																						
	2	3			8	Fraxinus pennsylvanica			2	4	3		3	3			3	2			1	2										
		2				Podophyllum peltatum			2	1			1	2																		
		1				Vitis sp. (seedling)			2	1			3	1	2		1	1														
		2				Fraxinus americana					3	2			4	2			3	2	1	2	2									
		2			6	Euonymous obovatus			1	2																						
		2			6	Cornus sp.			1	2					2	2																
		1				Aster sp. #1			1	1			1	1																		
6						Quercus rubra			4	2			2	7	4		2	6	2													
	2	2			2	Berberis thunbergii			1	2			1	2																		
		2				Fraxinus sp. (seedling)							4	2					3	2			2	2								
		2				Ulmus sp. (seedling)			1	1			4	2	2																	
		1				Carya sp. (seedling)							3	1																		
		1			10	Unknown dicot #1		CS-2607					3	1																		
		2				Moss sp.							3	1							1	2										
		1				Quercus sp. (seedling)							2	1																		

ugh w/ - nceolate 's.

Solidago - caesia?

EXAMPLES OF PERCENT OF AREA COVERED

The following graphic can be used for various data elements to convey "Amount" or "Quantity" *NOTE*: Within any given box, each quadrant contains the same total area covered, just different sized objects.



cover class	% cover	midpoint
1	solitary or few	0.0001
2	0-1%	0.005
3	1-2%	0.015
4	2-5%	0.035
5	5-10%	0.075
6	10-25%	0.175
7	25-50%	0.375
8	50-75%	0.625
9	75-95%	0.850
10	95-100%	0.975







BROWSE RATING NARRATIVE DESCRIPTION

LOW OR NONE: there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.

MEDIUM LOW values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

MEDIUM: browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.

MEDIUM HIGH values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.

HIGH: greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.

VERY HIGH values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.



CLEVELAND METROPARKS Plant Community Assessment Program Species Cover Data Sheet 2a Page 2 of 2

Project Label: PCAP Project name: DITH2013 Plot no.: 1389

Total modules: 10 Intensive modules: 4 Plot configuration: 2x5 Plot area (ha): .1

Cleveland Metroparks

Br = Browse Level. Use cover classes to describe amount of browse per species over entire plot

Strata - Cov. entire plot

Estimate for each intensive module:

	mod 2	corner 4	mod 2	corner 2	mod 3	corner 4	mod 3	corner 2	mod 8	corner 4	mod 8	corner 2	mod 9	corner 4	mod 9	corner 2	mod R	corner R
	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov
%open water					1				1				1					
%unvegetated open water					1				1				1					
%unveg. ground (bare soil)					1				1				1					
%unveg. litter (bare litter)					1				1				1					

T	S	H	(F)	(A)	Br	Species	c	Voucher #	mod 2 depth	mod 2 cov	corner 4 depth	corner 4 cov	mod 2 depth	mod 2 cov	corner 2 depth	corner 2 cov	mod 3 depth	mod 3 cov	corner 4 depth	corner 4 cov	mod 3 depth	mod 3 cov	corner 2 depth	corner 2 cov	mod 8 depth	mod 8 cov	corner 4 depth	corner 4 cov	mod 8 depth	mod 8 cov	corner 2 depth	corner 2 cov	mod 9 depth	mod 9 cov	corner 4 depth	corner 4 cov	mod 9 depth	mod 9 cov	corner 2 depth	corner 2 cov	mod R depth	mod R cov	corner R depth	corner R cov
		2				Fagus grandifolia											1	2																										
		1				~~Unknown dicot #2~~ Hackelia virginiana		CS-2608									1	1																										
		2				Solidago caesia		10-30-13									2	1								2	1																	
4		2			4	Sassafras albidum											1	2																										
		2				Viburnum acerifolium											1	2																										
6		1			10	Liriodendron tulipifera											1	1																										
		2				Crataegus sp.			1	2							1	1																										
		2				Toxicodendron radicans											1	1																										
		2				Ribes sp.	X	SC-233									1	2																										
		1				Monotropa uniflora											1	1																										
		1				Prenanthes sp.																	Ø			2	1																
		2			10	Rosa multiflora																	1	2																		
		1				Circaea lutetiana																	1										1	1								
		1				Geum sp.																																			R	1	
		2				Alliaria petiolata																																			R	2	
4						Tilia americana																																			R	4	
						~~Liriodendron tulipifera~~																																			R		
4						Carya sp.																																			R	4	
	2					Rubus occidentalis																																			R	2	
5	5					Vitis ~~sp.~~ aestivalis	X	SC-234																																	R	6	
						SRE 11-6-13																																						

nid nuts

may flr

2aCM PCAP Species Cover Data sheet Page 1 of x_ver 3.xls last revised 5/29/2012 ceh

Natural Resource Management FORM NR/2010-02a

EXAMPLES OF PERCENT OF AREA COVERED

The following graphic can be used for various data elements to convey "Amount" or "Quantity". *NOTE*: Within any given box, each quadrant contains the same total area covered, just different sized objects.



cover class	% cover	midpoint
1	solitary or few	0.0001
2	0-1%	0.005
3	1-2%	0.015
4	2-5%	0.035
5	5-10%	0.075
6	10-25%	0.175
7	25-50%	0.375
8	50-75%	0.625
9	75-95%	0.850
10	95-100%	0.975

Nested Corners





BROWSE RATING NARRATIVE DESCRIPTION

LOW OR NONE: there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.

MEDIUM LOW values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

MEDIUM: browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.

MEDIUM HIGH values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.

HIGH: greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.

VERY HIGH values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.



CLEVELAND METROPARKS Plant Community Assessment Program Natural Woody Stem Data Sheet

Project Label: PCAP Project Name: 01H: 2013 Plot No.: 1389 Page: 1 of 3

Cleveland Metroparks

Explain subsample (additional room on back):

mod #	species	c	voucher#	# stems 0-1.4m browsed	% sub or super sample	# shrub clumps	size class (cm) woody stems >1.4m 1 0-<1	2 1-<2.5	3 2.5-<5	4 5-<10	5 10 - <15	6 15 - <20	7 20 - <25	8 25 - <30	9 30 - <35	10 35 - <40	11 >40 (record each tree)
1	Standing dead							1		1	1						
1	Acer saccharum			2			2	4	2	2		1		1			
1	Quercus velutina																54.8
1	BERBERIS THUNBERGII			1		2											
1	Carya cordiformis														1		
1	ROSA MULTIFLORA			1													
1	Parthenocissus quinquefolia			1													
2	Ostrya virginiana								2								
2	Acer saccharum			2			3	3					1				43.9
2	Standing dead						5	1		1							
2	Prunus serotina			1													
2	Lindera benzoin			1													
2	Euonymus obovatus			2													
2	BERBERIS THUNBERGII					1											
3	Acer saccharum			1			2	3					2				68.7
3	Standing dead								2	1							
3	Ostrya virginiana						1										
3	Lindera benzoin					1											
3	Parthenocissus quinquefolia			2		3											
3	BERBERIS THUNBERGII			1		2											
3	Fraxinus pennsylvanica			1													
3	Cornus sp.			1													
3	Prunus serotina			1													
4	Quercus rubra																116.9

DBH Measurement Rules



Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.

Record using the tally system from 1 to 10

1 2 3 4 5

ASH CANOPY CONDITION

1. **Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
2. **Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
3. **Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
4. **>50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
5. **Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.

A B C D E

ASH CANOPY BREAKUP CONDITION (for dead trees):
(If an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)

A: All main branches contain fine twigs (newly dead).
B: Over 50% of main branches have fine twigs.
C: Less than 50% of main branches have fine twigs.
D: Stem still standing and tertiary main branches present.
E: Central stem still standing.

CLEVELAND METROPARKS Plant Community Assessment Program Natural Woody Stem Data Sheet

Project Label: PCAP Project Name: 01Hi2013 Plot No.: 1389 Page: 2 of 3

Explain subsample (additional room on back):

mod #	species	c	voucher#	# stems 0-1.4m browsed	% sub or super sample	# shrub clumps	size class (cm) woody stems >1.4m 1 0-<1	2 1-<2.5	3 2.5-<5	4 5-<10	5 10 - <15	6 15 - <20	7 20 - <25	8 25 - <30	9 30 - <35	10 35 - <40	11 >40 (record each tree)
4	Acer saccharum							••	••			•	•				
4	Standing dead							•			•						
4	Ostrya virginiana									••							
4	Fraxinus sp.			•													
5	Acer saccharum			•			••	⊠	⁝	•	•						
5	Standing dead									•							
5	Ostrya virginiana									•							
5	Liriodendron tulipifera			•						•		•				•	62.2
6	Acer saccharum						•X	⁝	⊠	••		••					
6	Standing dead							••	••	••							
6	Ostrya virginiana									•							
6	Sassafras albidum														•		
6	Carya sp.									•							
6	Carya cordiformis											••					
7	Acer saccharum						••	⁝	⁝	••	••	••					
7	Standing dead						••		⁝								
7	Tilia americana										•						
8	Acer saccharum			•			⁝	⊠	⁝	•	•						
8	Standing dead						•	⁝									
8	ROSA MULTIFLORA			•													
9	Acer saccharum			•			⊠	⊠	⁝								
9	Standing dead						⊠	⁝									
9	Ostrya virginiana											•			•		
9	Carya cordiformis																57.3, 44.6, 40.1

DBH Measurement Rules



Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.

Record using the tally system from 1 to 10

1

2

3

4

5

ASH CANOPY CONDITION

1. **Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
2. **Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
3. **Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches not exposed to sunlight, die naturally and are not considered.
4. **>50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
5. **Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.

A

B

C

D

E

ASH CANOPY BREAKUP CONDITION (for dead trees):
(If an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)

A: All main branches contain fine twigs (newly dead).
B: Over 50% of main branches have fine twigs.
C: Less than 50% of main branches have fine twigs.
D: Stem still standing and tertiary main branches present.
E: Central stem still standing.

CLEVELAND METROPARKS Plant Community Assessment Program Natural Woody Stem Data Sheet

Project Label: PCAP Project Name: OIH:2013 Plot No.: 1389 Page: 3 of 3

Cleveland Metroparks

Explain subsample (additional room on back):

mod #	species	c	voucher#	# stems 0-1.4m browsed	% sub or super sample	# shrub clumps	size class (cm) woody stems >1.4m 1 0-<1	2 1-<2.5	3 2.5-<5	4 5-<10	5 10 - <15	6 15 - <20	7 20 - <25	8 25 - <30	9 30 - <35	10 35 - <40	11 >40 (record each tree)
10	Acer saccharum			••			5	5	3	••	••						
10	Standing dead						•	1	•								
10	Carya cordiformis												•				
10	Ostrya virginiana									•							
10	Euonymus obovatus			••													

DBH Measurement Rules



Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.

Record using the tally system from 1 to 10

1 2 3 4 5

ASH CANOPY CONDITION

1. **Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
2. **Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
3. **Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
4. **>50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
5. **Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.

A B C D E

ASH CANOPY BREAKUP CONDITION (for dead trees):
(if an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)

A: All main branches contain fine twigs (newly dead).
B: Over 50% of main branches have fine twigs.
C: Less than 50% of main branches have fine twigs.
D: Stem still standing and tertiary main branches present.
E: Central stem still standing.

CLEVELAND METROPARKS Emerald Ash Borer - Fraxinus Sheet *INTENSIVE MODULES ONLY* *TREES ≥ 10CM ONLY*

Project Label: PCAP Project Name: 01Hi2013 Plot No.: 1384 Date: 8-19-13

Module	Tree ID.	Species	Dead	c	Voucher #	DBH (cm)	Ht @ DBH	ASH Only: Ash condition	*Dead condition	# Exit holes	Epicormic present	Woodpecker holes
	1	No Ash										
	2											
	3											
	4											
	5											
	6											
	7											
	8											
	9											
	10											
	11											
	12											
	13											
	14											
	15											
	16											
	17											
	18											
	19											
	20											
	21											
	22											
	23											
	24											
	25											



*** Change intensive module numbers when necessary



Map all ash trees ≥10cm in each module using Tree ID number

* If Ash Condition scores 5 (dead) provide breakup score (A-E)
Count EAB exit holes 1.25m≥ x ≥1.5m
Woodpecker and epicormic marked present (1) or absent (0)

CLEVELAND METROPARKS Plant Community Assessment Program: Invasive Species Survey Cleveland Metroparks

Tier 1: Early detection/ Rapid response		Presence				GPS
		NE	SE	SW	NW	
Microstegium vimineum	Japanese stiltgrass					
Ranunculus ficaria	Lesser Celandine					
Cynanchum louiseae (vine)	Black Swallow-wort					
Butomus umbellatus (wetland)	Flowering Rush					
Heracleum mantegazzianum	Giant Hogweed					

Tier 2: Assess as Needed		# of Plants				comments
		NE	SE	SW	NW	
Acer platanoides	Norway Maple					
Ailanthus altissima	Tree of Heaven					
Lonicera japonica (vine)	Japanese Honeysuckle		1			
Lythrum salicaria (wetland)	Purple Loosestrife					
Aegopodium podagraria (G-cover)	Bishop's Goutweed					
Celastrus orbiculatus (vine)	Asian Bittersweet					
Torilis sp.	Hedgeparsley					
Conium maculatum	Poison Hemlock					
Rhamnus cathartica	Common Buckthorn (shrub)					
Berberis thunbergii	Japanese Barberry (shrub)	2	2	2	1	
Alnus glutinosa	European Alder					
Dipsacus laciniatus	Cut-leaf Teasel					
Elaeagnus umbellata	Autumn Olive (shrub)					
Lonicera maackii	Amur Honeysuckle (shrub)		2		1	
Euonymus fortunei	Wintercreeper					

Tier 3: Presence is of Interest		# of Plants				comments
		NE	SE	SW	NW	
Convallaria majalis (G-cover)	Lily of the Valley					
Coronilla varia (G-cover)	Crown Vetch					
Eleutherococcus pentaphyllus	Five-leaf Aralia (shrub)					
Pachysandra terminalis (G-cover)	Japanese Pachysandra					
Philadelphus coronarius	Mock Orange (shrub)					
Pulmonaria officinalis (G-cover)	Lungwort					
Rubus phoenicolasius	Wineberry					
Iris pseudacorus (wetland)	Yellow Flag Iris					
Ornithogalum umbellatum	Star of Bethlehem					
Viburnum opulus var. opulus	European Cranberry (shrub)					
Viburnum plicatum	Doublefile Viburnum (shrub)					

Tier 4: Widespread and abundant		Presence				comments
		NE	SE	SW	NW	
Alliaria petiolata	Garlic Mustard	2	2	2	1	
Ligustrum vulgare	Common Privet (shrub)	1	2		1	
L. morrowii, L. tatarica	Bush Honeysuckles (shrub)				1	
Phalaris arundinacea	Reed Canarygrass					
Phragmites australis (wetland)	Phragmites					
Polygonum cuspidatum	Japanese Knotweed					
Frangula alnus	Glossy Buckthorn (shrub)	1				
Rosa multiflora	Multiflora Rose (shrub)	2	2	1	4	
Typha angustifolia, T. x.glauca	Cattails (wetland)					
Cirsium arvense	Canada thistle					
Dipsacus fullonum	Common Teasel		1			
Hesperis matronalis	Dame's Rocket					
Vinca minor (G-cover)	Periwinkle					

Presence
X: yes

of Plants
1: 1-10
2: 11-50.
3: 51-100
4: 101-1,000
5: >1,000

of Plants
1: 1-10
2: 11-50.
3: 51-100
4: 101-1,000
5: >1,000

of Plants
1: 1-10
2: 11-50.
3: 51-100
4: 101-1,000
5: >1,000

Note: For Ground-cover plants record "stem #" but in comment field describe # of colonies and patch size (S,M, L)

CLEVELAND METROPARKS Plant Community Assessment Program - Plant Cover and Earth Surface

Project Label: PCAP Project Name: 01 Hi 2013 Plot No.: 1389

Cleveland Metroparks

STANDING BIOMASS (required for emergent wetlands): collected in 0.1m clip plots (32x32 cm) from corners 1 and 3 in each intensive module. Required for VIBI-E score calculation. C?=check when collected

Module #	C?	Corner	Corner

CLASSIFICATION

(FIT = excellent, g Fit and Confidence

Hydrogeomorphic class (WETLANDS ONLY):

- ◻ DEPRESSION Fit=___ Conf=___
- ◻ IMPOUNDMENT ◻ Beaver ◻ Human Fit=___ Conf=___
- ◻ RIVERINE ◻ Headwater ◻ Mainstem ◻ Channel Fit=___ Conf=___
- ◻ SLOPE (ground water hydrology or on a physical slope) Fit=___ Conf=___
- ◻ FRINGING ◻ Reservoir ◻ Natural Lake Fit=___ Conf=___
- ◻ COASTAL (specify subclass) Fit=___ Conf=___
- ◻ BOG (strongly, moderately, weekly ombrotrophic) Fit=___ Conf=___

Ohio EPA VIBI Plant Community Class (WETLANDS ONLY):

- ◻ FOREST ◻ swamp forest ◻ bog forest ◻ forest seep Fit=___ Conf=___
- ◻ EMERGENT ◻ marsh ◻ wet meadow ◻ open bog Fit=___ Conf=___
- ◻ SHRUB ◻ shrub swamp ◻ tall sh. bog ◻ tall sh. fen Fit=___ Conf=___

McNAB INDICES (degrees) + for up - for down

[FILLED OUT USING GIS PROGRAM - DO NOT FILL OUT IN FIELD]

		LFI*	TSI**
At aspect	N		
+45 degrees	NE		
+90 degrees	E		
+135 degrees	SE		
+180 degrees	S		
+225 degrees	SW		
+270 degrees	W		
+315 degrees	NW		

LFI is angle of plot to the horizon. TSI is angles formed by local slopes. For TSI measure angle from recorders eye to eye of person standing ~10 m away.

* Landform Index (position within landscape)

** Terrain Shape Index (site microtopographic shape)

MICROTOPOGRAPHIC FEATURE COUNTS - Intensive modules only

Ranks for microhabitat features. Select one or select two and average the score. **NOTE:** If mod falls on a slope automatically gets ranked based on steepness (1-3) to begin + any features present

Slope 1 = slight elevational grade across module (hill) **Slope 2** = falls on slope ~20° **Slope 3** = maximum steepness that can be safely sampled ~45°

0 feature is absent or functionally absent from the wetland

3 feature is present in the wetland in very small amounts or if more common, of low quality

7 feature is present in moderate amounts, but not of highest quality, or in small amounts of highest quality

10 feature is present in moderate or greater amounts and of highest quality

c.w.d. - count for pieces with minimum 1m length

mod#	corner	no. of tussocks depth 3 1x1m (count)	no. of hummocks uplands (Tip-Ups) depth 2 3.16x3.16m (count)	no. macro. depressions depth 1 10x10m (count)	c.w.d (2-12 cm) depth 1 10x10m (count)	c.w.d (12-40cm) depth 1 10x10m (count)	c.w.d >40 cm depth 1 10x10m (count)	microhab. interspers. depth 1 10x10m (rank)	microhab. SLOPE 10x10m (rank)
2		0	0	2	36	1	0	4	1
3		0	0	1	26	1	0	3	1
8		0	0	0	23	1	0	3	2
9		0	0	1	22	0	0	3	2

NOTE: tussock and hummocks are counted in BOTH nested quadrat corners but counts are aggregated.

CROWN COVER (DENSIOMETER): Make 4 readings per module facing N, S, E, W. Place dot count in corresonding space. (4 dots per grid square)

Module	N	S	E	W
2	4	2	3	3
3	5	12	9	4
8	5	3	2	4
9	2	1	2	3

COVER BY STRATA

STRATUM	GENERAL FORM
Tree (generally >5 m)	Tree (overstory), very tall shrubs*, liana, epiphyte)
Shrub (generally 0.5 to 5 m)	Tree (sapling), shrub, liana, epiphyte)
Herb (Field)	Herb, dwarf-shrub**, tree (seedling***)
Floating	Floating
Aquatic (submerged)	Submerged

*Very tall shrubs are sometimes included in the tree stratum
**Can also include seedlings of shrubs, i.e. all shrubs <0.5m
***Tree seedlings are often defined as up to 1.4 m height or as <2.5 cm DBH in which case they would span the herb and shrub layers.





FIGURE 3-20.—Generalized section of Upper Devonian, Mississippian, and Lower Pennsylvanian formations in northeastern Ohio. Asterisks indicate units that are fossiliferous. This composite section represents about 400 meters of rock exposed across the area. The section is not to scale, but the thicknesses indicated are proportional. The term "Waverly" is used in the older literature to refer to Mississippian rocks in Ohio. Some geologists use the European term "Carboniferous," which encompasses the Mississippian and Pennsylvanian Periods of the U.S. Many units have been named within the Cuyahoga Formation, but most units are local and cannot be traced over great distances. The Black Hand Member is a spectacular massive sandstone that is fairly widespread but discontinuous. See Hyde (1953), Hoover (1960), and Colbns 19~9) for more information on Mississippian rocks in Ohio. See figure 3-18 for explanation of rock types.

CLEVELAND METROPARKS Plant Community Assessment Program - Soils, Crown Cover, Standing Biomass Data Sheet 6a

Project label: PCAP Project Name: 01 Hi 2013 Plot No.: 1389

Cleveland Metroparks

SOIL PIT DESCRIPTION: Excavate 20 cm plug wih shovel. Describe using Munsell chart, visual exam, texture, and odor.

Soil pit module # 3 (one per entire plot)

5 cm	matrix color	2.5Y3/3
	mottle color	—
	%mottle	0
	oxid roots	Y (N)
	texture*	2
	redox features**	Y (N)
	hydr. cond.***	I S M (D)
20 cm	matrix color	2.5Y4/2
	mottle color	—
	%mottle	0
	oxid roots	Y (N)
	texture*	2
	redox features**	Y (N)
	hydro. cond.***	I S M (D)

* refer to texture classes on reverse side

** e.g. hydrogen sulfide odor, gleying, etc.

*** Circle one: I=indundated S=saturated M=moist D=dry

Notes: include evidence of earthworms (worms, castings, middens)

2 worms found. No castings or middens.

SOIL SAMPLES Standard procedure: collect a soil sample of the top 10 cm of soil from center of each intensive module and composite the sample

Soil Collection Module	Horizon (A, B, C)
2,3,8,9 composited	A

RSE 23 Aug 2013

Web Soil Survey Information:

Soil Series/Type: ESC2/Ellsworth silt loam... Sandstone substratum, mildly eroded

Soil Series Source: Ohio Soil Survey

Landform type: Till Plains

Depth to rest. Layer: 40"-60" to paralithic Bedrock 76.2

Parent Material: Till

DRAINAGE*

- ☐ Excessively dr.
- ☐ Somewhat excessively
- ☐ Well drained
- ☒ Moderately well dr.
- ☐ Somewhat poorly dr.
- ☐ Very poorly dr.
- ☐ Impermeable surface

SOIL DEPTH MEASUREMENT: Measure to the nearest 0.1 cm in center of intensive modules. If >30.5 cm, record as >30

mod#	1 litter+ organic depth (cm)	2 litter depth (cm)	water depth (cm)	depth sat soil (cm)
2	1.0	1.0	0	730.0
3	1.0	1.0	0	730.0
8	2.0	2.0	0	730.0
9	0.3	0.3	0	730.0

EARTH SURFACE & GROUND COVER

Underlying Earth Surface* (Sum = 100%)	percent	Ground Cover (Each ≤ 100%)	percent
Histosol	0	Coarse Woody Debris***	15
Mineral Soil	97	Fine Woody Debris****	15
Gravel-Cobble*	3	Litter	95
Boulder**	0	Duff (Ferm.+ Humus)	0
Bedrock	0	Bryophyte- Lichen	2
* Gravel-Cobble = 1/16-10"		Water	0
**Boulder = > 10 in		Bare Soil	5
*** >5 cm in diameter		Road/Trail	0
**** <5 cm in diameter		Other	na

TRAIL INFORMATION:

record type and cover for each

Type	%Cover
☐ All Purpose	
☐ Bridle	NO
☐ Hiking sanctioned	TRAILS
☐ Bootleg unsanctioned	
☐ Gravel	
☐ Deer	

COVER BY STRATA %

estimate using midpoints of 5,ex:3, 8, 13

Strata	Height Range (m)	Total Cover (%)
Tree	5 -	93
Shrub	.5 - 5	58
Herb	0 - .5	8
(Floating)*	-	na
(Aquatic)*	-	na

* rooted and floating or slightly emersed

** submersed, most plant mass below surface

SEE BACK OF PAGE FOR "TYPICAL"STRATA DESCRIPTIONS. STRATA CAN VARY BY COVER TYPE.

STAND SIZE

- ☐ >600 x plot size
- ☐ > 100 x plot size
- ☐ 10-100 x plot size
- ☒ 3-10 x plot size
- ☐ 1-3 x plot size
- ☐ < plot size

PERCENT MOTTLES (USE CLASS CODES):

Class	Code		Criteria: % of Surface Area Covered
	Conv.	NASIS	
Few	f	#	< 2
Common	c	#	2 to < 20
Many	m	#	≥ 20



SOIL TEXTURE: Record the code for the soil texture of the 5 cm and 20 cm layers. To estimate texture, collect a soil sample from the appropriate layer and moisten it with water to the consistency of modeling clay/wet newspaper; the sample should be wet enough that all of the particles are saturated but excess water does not freely flow from the sample when squeezed. Attempt to roll the sample into a ball. If the soil will not stay in a ball and has a grainy texture, the texture is either sandy or coarse sandy. If the soil does form a ball, squeeze the sample between your fingers and attempt to form a self-supporting ribbon. Samples which form both a ball and a ribbon should be coded as clayey; samples which form a ball but not a ribbon should be coded as loamy.

0= Organic
1= Loamy
2= Clayey
3= Sandy
4= Coarse Sand
9= Not measured - make plot note

Geomorphic Component - Three-dimensional descriptors of parts of landforms or microfeatures that are best applied to areas. Unique descriptors are available for Hills, Terraces, Mountains, and Flat Plains; e.g., (for Hills) *nose slope* or *NS*.

Hills	Code	
	PDP	NASIS
interfluve	IF	IF
head slope	HS	HS
nose slope	NS	NS
side slope	SS	SS
base slope	---	BS



Terraces	Code
riser	RI
tread	TR



Hillslope - Profile Position (Hillslope Position in PDP) - Two-dimensional descriptors of parts of line segments (i.e., slope position) along a transect that runs up and down the slope; e.g., *backslope* or *BS*. This is best applied to transects or points, not areas.

Position	Code
summit	SU
shoulder	SH
backslope	BS
footslope	FS
toeslope	TS



HYDROLOGIC REGIME Modified from Grossman et al 1998. (Frequency and duration of flooding.)

UPLAND: Not a wetland. Very rarely flooded.

INTERMITTENTLY/SEASONALLY SATURATED: Dry at least once per year. Surface water is seldom present, but substrate is saturated to surface for extended periods during the growing season.

PERMANENTLY/SEMIPERMANENTLY SATURATED: Dry less than once per year. Surface water is seldom present, but substrate is saturated to surface for extended periods during the growing season. Equivalent to Cowardin's Saturated modifier.

OCCASIONALLY FLOODED: Surface water can be present for brief periods during growing season, but not in most years. Often characterizes flood-plain upper terraces.

TEMPORARILY FLOODED: Surface water present for brief periods during growing season, but water table usually lies well below soil surface. Often characterizes flood-plain levees and lower terraces. Equivalent to Cowardin's Temporary modifier.

INTERMITTENTLY FLOODED : Substrate is usually exposed, but surface water can be present for variable periods without detectable seasonal periodicity. Inundation is not predictable to a given season and is dependent upon highly localized rain storms. This modifier was developed for use in the arid West for water regimes of Playa lakes, intermittent streams, and dry washes but can be used in other parts of the U.S. where appropriate. This modifier can be applied to both wetland and non-wetland situations. Equivalent to Cowardin's Intermittently Flooded modifier.

SEMIPERMANENTLY FLOODED (exposed <1/year): Surface water persists throughout the growing season in most years. Land surface is normally saturated when water level drops below soil surface. Includes Cowardin's Intermittently Exposed and Semipermanently Flooded modifiers.

PERMANENTLY FLOODED: Water covers the land surface at all times of the year in all years. Equivalent to Cowardin's "permanently flooded".

UNKNOWN: The hydrologic regime cannot be determined from the available information.

FORM B-1: BUFFER SAMPLE PLOTS (Front)

Reviewed by (initial): _____

Site ID: PCAP Hi1389 DATE: 08 / 19 / 2013

Location: ● AA Center ○ N ○ S ○ E ○ W

Fill in bubble(s) if plot(s) could not be sampled and flag → ○ Plot 1 ○ Plot 2 ○ Plot 3

Buffer Natural Cover Strata

Fill in bubbles for all that apply: Canopy Type: D = Deciduous; E = Evergreen. Leaf Type: B = Broadleaf; N = Needle Leaf. Absent: No tree canopy.
Strata Section: Fill in appropriate cover class bubble for each strata type for each plot. 0 = Absent; 1 = Sparse(<10%); 2=Moderate(10-40%); 3 = Heavy (40-75%); 4 = Very Heavy (>75%)

Buffer Plot 1	Canopy Type: ● E	Leaf Type: ● N	Absent: ○	Flag
Big Trees (>0.3m DBH)	0	1	2	● 3	4	
Small Trees (<0.3m DBH)	0	1	2	● 3	4	
Woody Shrubs, Saplings (0.5m-5m HIGH)	0	1	2	● 3	4	
Woody Shrubs, Saplings (<0.5m HIGH)	0	● 1	2	3	4	
Herbs, Forbs and Grasses	0	● 1	2	3	4	
Bare ground	0	● 1	2	3	4	
Litter, duff	0	1	2	3	● 4	
Rock	0	● 1	2	3	4	
Water	● 0	1	2	3	4	
Submerged Vegetation	● 0	1	2	3	4	

Buffer Plot 2	Canopy Type: ○ D ○ E	Leaf Type: ○ B ○ N	Absent: ○	Flag
Big Trees (>0.3m DBH)	0	1	2	3	4	
Small Trees (<0.3m DBH)	0	1	2	3	4	
Woody Shrubs, Saplings (0.5m-5m HIGH)	0	1	2	3	4	
Woody Shrubs, Saplings (<0.5m HIGH)	0	1	2	3	4	
Herbs, Forbs and Grasses	0	1	2	3	4	
Bare ground	0	1	2	3	4	
Litter, duff	0	1	2	3	4	
Rock	0	1	2	3	4	
Water	0	1	2	3	4	
Submerged Vegetation	0	1	2	3	4	

Buffer Plot 3	Canopy Type: ○ D ○ E	Leaf Type: ○ B ○ N	Absent: ○	Flag
Big Trees (>0.3m DBH)	0	1	2	3	4	
Small Trees (<0.3m DBH)	0	1	2	3	4	
Woody Shrubs, Saplings (0.5m-5m HIGH)	0	1	2	3	4	
Woody Shrubs, Saplings (<0.5m HIGH)	0	1	2	3	4	
Herbs, Forbs and Grasses	0	1	2	3	4	
Bare ground	0	1	2	3	4	
Litter, duff	0	1	2	3	4	
Rock	0	1	2	3	4	
Water	0	1	2	3	4	
Submerged Vegetation	0	1	2	3	4	

Stressor Presence/Absence - Confirm that a filled data bubble indicates presence and an unfilled bubble indicates absence by filling this bubble. ●

Residential and Urban Stressors

Fill bubble if present - Plot	1	2	3	Flag
Road - gravel	○	○	○	
Road - two lane	○	○	○	
Road - four lane	○	○	○	
Parking Lot/Pavement	○	○	○	
Golf Course	○	○	○	
Lawn/Park	○	○	○	
Suburban Residential	○	○	○	
Urban/Multifamily	○	○	○	
Landfill	○	○	○	
Dumping	○	○	○	
Trash	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Hydrology Stressors

Fill bubble if present - Plot	1	2	3	Flag
Ditches, Channelization	○	○	○	
Dike/Dam/Road/RR Bed (IMPEDE FLOW)	○	○	○	
Water Level Control Structure	○	○	○	
Excavation, Dredging	○	○	○	
Fill/Spoil Banks	○	○	○	
Freshly Deposited Sediment (UNVEGETATED)	○	○	○	
Soil Loss/Root Exposure	○	○	○	
Wall/Riprap	○	○	○	
Inlets, Outlets	○	○	○	
Point Source/Pipe (EFFLUENT OR STORMWATER)	○	○	○	
Impervious surface input (SHEETFLOW)	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Agricultural & Rural Stressors

Fill bubble if present - Plot	1	2	3	Flag
Pasture/Hay	○	○	○	
Range	○	○	○	
Row Crops	○	○	○	
Fallow Field (RECENT-RESTING ROW CROP FIELD)	○	○	○	
Fallow Field (OLD - GRASS, SHRUBS, TREES)	○	○	○	
Nursery	○	○	○	
Dairy	○	○	○	
Orchard	○	○	○	
Confined Animal Feeding	○	○	○	
Rural Residential	○	○	○	
Gravel Pit	○	○	○	
Irrigation	○	○	○	
Other: _____	○	○	○	

Industrial Development Stressors

Fill bubble if present - Plot	1	2	3	Flag
Oil Drilling	○	○	○	
Gas Wells	○	○	○	
Mine (surface)	○	○	○	
Mine (underground)	○	○	○	
Military	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Habitat/Vegetation Stressors

Fill bubble if present - Plot	1	2	3	Flag
Forest Clear Cut	○	○	○	
Forest Selective Cut	○	○	○	
Tree Plantation	○	○	○	
Tree Canopy Herbivory (INSECT)	○	○	○	
Shrub Layer Browsed (WILD OR DOMESTIC)	●	○	○	
Highly Grazed Grasses (OVERALL <3" HIGH)	○	○	○	
Recently Burned Forest Canopy	○	○	○	
Recently Burned Grassland (BLACKENED)	○	○	○	

Fill bubble if present - Plot	1	2	3	Flag
Herbicide Use	●	○	○	i
Mowing/Shrub Cutting	○	○	○	
Trails	○	○	○	
Soil Compaction (ANIMAL OR HUMAN)	○	○	○	
Offroad vehicle damage	○	○	○	
Soil erosion (FROM WIND, WATER, OR OVERUSE)	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Flag codes: K = No measurement made, U = Suspect measurement., F1,F2, etc. = misc. flags assigned by each field crew.
Explain all flags in comment section on the back of this form

2428168304

FORM B-1: BUFFER SAMPLE PLOTS - TARGETED ALIEN SPECIES (Back)

Reviewed by (initial): _____

Site ID: PIAP Hi1389 **DATE:** 08 / 19 / 2013

● Confirm a filled data bubble indicates presence and an unfilled bubble indicates absence by filling in this bubble

Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag
Eurasian Watermilfoil	O	O	O		Purple Loosestrife	O	O	O		Johnson Grass	O	O	O	
Water hyacinth	O	O	O		Knotweed	O	O	O		Kudzu	O	O	O	
Yellow Floating Heart	O	O	O		Japanese Knotweed	O	O	O		Multiflora Rose	O	O	O	
Giant Salvinia	O	O	O		Perennial Pepperweed	O	O	O		Common Buckthorn	O	O	O	
Garlic Mustard	O	O	O		Giant Reed	O	O	O		Himalayan Blackberry	O	O	O	
Poison Hemlock	O	O	O		Cheatgrass	O	O	O		Tamarisk	O	O	O	
Mile-A-Minute Weed	O	O	O		Reed Canary Grass	O	O	O		Other: _____	O	O	O	
Birdsfoot Trefoil	O	O	O		Common Reed	O	O	O		Other: _____	O	O	O	
Canada Thistle	O	O	O		Leafy Spurge	O	O	O		Other: _____	O	O	O	
										Other: _____	O	O	O	

PLOT COORDINATES

Provide GPS coordinates at the center of the Buffer Plot (#3) at the far end of each Buffer Transect and for the Buffer Plot at the AA CENTER. Indicate the location of the plot coordinates by filling in the appropriate bubble.

If Buffer Plot 3 can not be accessed, take the coordinates at the nearest practicable location ALONG THE TRANSECT. This is important because all Buffer Plots are centered on the Buffer Transects and the coordinates will indicate the location of the transect. Fill in the "nearest practicable location" bubble, fill in the flag box, and describe where the coordinates were taken and why in the comment section below. The coordinates of the nearest practicable location can be either placed as close to the center of Plot 3 as possible or at the center of the last accessible Buffer Plot.

Location of coordinates (choose one):

● AA CENTER O N3 O S3 O E3 O W3 O Nearest practicable location (flag and comment below)

Flag: 2

Latitude North 41.22520 Longitude West 081.70567

Use Decimal Degrees; NAD83

Flag	Comments
1	Found very old evidence of herbicide use at far S side of plot, assume target area included AA center plot.
2	Took point at 50% accuracy due to bad sat. reception

7966623548

FORM B-1: BUFFER SAMPLE PLOTS (Front)

Reviewed by (initial): _____

Site ID: PCAPHi 1389 DATE: 08 / 19 / 2013

Location: ○ AA Center ● N ○ S ○ E ○ W

Fill in bubble(s) if plot(s) could not be sampled and flag → ○ Plot 1 ○ Plot 2 ○ Plot 3

Buffer Natural Cover Strata

Fill in bubbles for all that apply: Canopy Type: D = Deciduous; E = Evergreen. Leaf Type: B = Broadleaf; N = Needle Leaf. Absent: No tree canopy.
Strata Section: Fill in appropriate cover class bubble for each strata type for each plot. 0 = Absent; 1 = Sparse(<10%); 2=Moderate(10-40%); 3 = Heavy (40-75%); 4 = Very Heavy (>75%)

Buffer Plot 1	Canopy Type: ● E	Leaf Type: ● N	Absent: ○	Flag
Big Trees (>0.3m DBH)	0 1 ● 3 4			
Small Trees (<0.3m DBH)	0 1 2 3 ●			
Woody Shrubs, Saplings (0.5m-5m HIGH)	0 1 2 3 ●			
Woody Shrubs, Saplings (<0.5m HIGH)	0 ● 2 3 4			
Herbs, Forbs and Grasses	0 ● 2 3 4			
Bare ground	0 ● 2 3 4			
Litter, duff	0 1 2 3 ●			
Rock	0 ● 2 3 4			
Water	● 1 2 3 4			
Submerged Vegetation	● 1 2 3 4			

Buffer Plot 2	Canopy Type: ● E	Leaf Type: ● N	Absent: ○	Flag
Big Trees (>0.3m DBH)	● 1 2 3 4			
Small Trees (<0.3m DBH)	0 1 ● 3 4			
Woody Shrubs, Saplings (0.5m-5m HIGH)	0 1 2 3 ●			
Woody Shrubs, Saplings (<0.5m HIGH)	0 ● 2 3 4			
Herbs, Forbs and Grasses	0 ● 2 3 4			
Bare ground	0 ● 2 3 4			
Litter, duff	0 1 2 3 ●			
Rock	0 ● 2 3 4			
Water	● 1 2 3 4			
Submerged Vegetation	● 1 2 3 4			

Buffer Plot 3	Canopy Type: ○ D ● E	Leaf Type: ○ B ● N	Absent: ○	Flag
Big Trees (>0.3m DBH)	● 1 2 3 4			
Small Trees (<0.3m DBH)	0 1 2 3 ●			
Woody Shrubs, Saplings (0.5m-5m HIGH)	0 1 ● 3 4			
Woody Shrubs, Saplings (<0.5m HIGH)	0 ● 2 3 4			
Herbs, Forbs and Grasses	0 ● 2 3 4			
Bare ground	0 ● 2 3 4			
Litter, duff	0 1 2 3 ●			
Rock	0 ● 2 3 4			
Water	● 1 2 3 4			
Submerged Vegetation	● 1 2 3 4			

Stressor Presence/Absence - Confirm that a filled data bubble indicates presence and an unfilled bubble indicates absence by filling this bubble. ●

Residential and Urban Stressors

Fill bubble if present - Plot	1	2	3	Flag
Road - gravel	○	○	○	
Road - two lane	○	○	○	
Road - four lane	○	○	○	
Parking Lot/Pavement	○	○	○	
Golf Course	○	○	○	
Lawn/Park	○	○	○	
Suburban Residential	○	○	○	
Urban/Multifamily	○	○	○	
Landfill	○	○	○	
Dumping	○	○	○	
Trash	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Hydrology Stressors

Fill bubble if present - Plot	1	2	3	Flag
Ditches, Channelization	○	○	○	
Dike/Dam/Road/RR Bed (IMPEDE FLOW)	○	○	○	
Water Level Control Structure	○	○	○	
Excavation, Dredging	○	○	○	
Fill/Spoil Banks	○	○	○	
Freshly Deposited Sediment (UNVEGETATED)	○	○	○	
Soil Loss/Root Exposure	○	○	○	
Wall/Riprap	○	○	○	
Inlets, Outlets	○	○	○	
Point Source/Pipe (EFFLUENT OR STORMWATER)	○	○	○	
Impervious surface input (SHEETFLOW)	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Agricultural & Rural Stressors

Fill bubble if present - Plot	1	2	3	Flag
Pasture/Hay	○	○	○	
Range	○	○	○	
Row Crops	○	○	○	
Fallow Field (RECENT-RESTING ROW CROP FIELD)	○	○	○	
Fallow Field (OLD - GRASS, SHRUBS, TREES)	○	○	○	
Nursery	○	○	○	
Dairy	○	○	○	
Orchard	○	○	○	
Confined Animal Feeding	○	○	○	
Rural Residential	○	○	○	
Gravel Pit	○	○	○	
Irrigation	○	○	○	
Other: _____	○	○	○	

Industrial Development Stressors

Fill bubble if present - Plot	1	2	3	Flag
Oil Drilling	○	○	○	
Gas Wells	○	○	○	
Mine (surface)	○	○	○	
Mine (underground)	○	○	○	
Military	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Habitat/Vegetation Stressors

Fill bubble if present - Plot	1	2	3	Flag
Forest Clear Cut	○	○	○	
Forest Selective Cut	○	○	○	
Tree Plantation	○	○	○	
Tree Canopy Herbivory (INSECT)	○	○	○	
Shrub Layer Browsed (WILD OR DOMESTIC)	●	●	●	
Highly Grazed Grasses (OVERALL <3" HIGH)	○	○	○	
Recently Burned Forest Canopy	○	○	○	
Recently Burned Grassland (BLACKENED)	○	○	○	

Fill bubble if present - Plot	1	2	3	Flag
Herbicide Use	○	○	○	
Mowing/Shrub Cutting	○	○	○	
Trails	○	○	○	
Soil Compaction (ANIMAL OR HUMAN)	○	○	○	
Offroad vehicle damage	○	○	○	
Soil erosion (FROM WIND, WATER, OR OVERUSE)	○	○	○	
Other: tree fall	○	●	○	1
Other: _____	○	○	○	

Flag codes: K = No measurement made, U = Suspect measurement., F1,F2, etc. = misc. flags assigned by each field crew. Explain all flags in comment section on the back of this form

2428168304

MR lig GM
Ber Rham

FORM B-1: BUFFER SAMPLE PLOTS - TARGETED ALIEN SPECIES (Back)

Reviewed by (initial): _____

Site ID: PCAP Hi 1389 **DATE:** 08 / 19 / 2013

● Confirm a filled data bubble indicates presence and an unfilled bubble indicates absence by filling in this bubble

Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag
Eurasian Watermilfoil	○	○	○		Purple Loosestrife	○	○	○		Johnson Grass	○	○	○	
Water hyacinth	○	○	○		Knotweed	○	○	○		Kudzu	○	○	○	
Yellow Floating Heart	○	○	○		Japanese Knotweed	○	○	○		Multiflora Rose	○	○	●	
Giant Salvinia	○	○	○		Perennial Pepperweed	○	○	○		Common Buckthorn	○	○	○	
Garlic Mustard	○	○	○		Giant Reed	○	○	○		Himalayan Blackberry	○	○	○	
Poison Hemlock	○	○	○		Cheatgrass	○	○	○		Tamarisk	○	○	○	
Mile-A-Minute Weed	○	○	○		Reed Canary Grass	○	○	○		Other: _____	○	○	○	
Birdsfoot Trefoil	○	○	○		Common Reed	○	○	○		Other: _____	○	○	○	
Canada Thistle	○	○	○		Leafy Spurge	○	○	○		Other: _____	○	○	○	
										Other: _____	○	○	○	

PLOT COORDINATES

Provide GPS coordinates at the center of the Buffer Plot (#3) at the far end of each Buffer Transect and for the Buffer Plot at the AA CENTER. Indicate the location of the plot coordinates by filling in the appropriate bubble.

If Buffer Plot 3 can not be accessed, take the coordinates at the nearest practicable location ALONG THE TRANSECT. This is important because all Buffer Plots are centered on the Buffer Transects and the coordinates will indicate the location of the transect. Fill in the "nearest practicable location" bubble, fill in the flag box, and describe where the coordinates were taken and why in the comment section below. The coordinates of the nearest practicable location can be either placed as close to the center of Plot 3 as possible or at the center of the last accessible Buffer Plot.

Location of coordinates (choose one):

○ AA CENTER ● N3 ○ S3 ○ E3 ○ W3 ○ Nearest practicable location (flag and comment below)

Flag: ☐

Latitude North 41.22639 Longitude West 081.70522

Use Decimal Degrees; NAD83

Flag	Comments
1	Enormous red oak recently fallen through center of plot 2 creating a decent sized light gap.

7966623548

FORM B-1: BUFFER SAMPLE PLOTS (Front)

Reviewed by (initial): _____

Site ID: PCAP Hi 1389 DATE: 08 / 19 / 2013

Location: ○ AA Center ○ N ○ S ● E ○ W

Fill in bubble(s) if plot(s) could not be sampled and flag → ○ Plot 1 ○ Plot 2 ○ Plot 3

Buffer Natural Cover Strata

Fill in bubbles for all that apply: Canopy Type: D = Deciduous; E = Evergreen. Leaf Type: B = Broadleaf; N = Needle Leaf. Absent: No tree canopy.
Strata Section: Fill in appropriate cover class bubble for each strata type for each plot. 0 = Absent; 1 = Sparse(<10%); 2=Moderate(10-40%); 3 = Heavy (40-75%); 4 = Very Heavy (>75%)

Buffer Plot 1	Canopy Type: ● (L) Leaf Type: ● (N)	Absent: ○	Flag
Big Trees (>0.3m DBH)	⊗ 1 ● 3 4		
Small Trees (<0.3m DBH)	0 1 2 3 ●		
Woody Shrubs, Saplings (0.5m-5m HIGH)	0 1 2 3 ●		
Woody Shrubs, Saplings (<0.5m HIGH)	0 1 2 3 4		
Herbs, Forbs and Grasses	0 1 ● 3 4		
Bare ground	0 ● 2 3 4		
Litter, duff	0 1 2 3 ●		
Rock	0 ● 2 3 4		
Water	● 1 2 3 4		
Submerged Vegetation	● 1 2 3 4		

Buffer Plot 2	Canopy Type: ● (E) Leaf Type: ● (N)	Absent: ○	Flag
Big Trees (>0.3m DBH)	● 1 2 3 4		
Small Trees (<0.3m DBH)	0 1 2 3 ●		
Woody Shrubs, Saplings (0.5m-5m HIGH)	0 1 2 ● 4		
Woody Shrubs, Saplings (<0.5m HIGH)	0 ⊗ ● 3 4		
Herbs, Forbs and Grasses	0 ● 2 3 4		
Bare ground	0 1 ● 3 4		
Litter, duff	0 1 2 3 ●		
Rock	0 ● 2 3 4		
Water	● 1 2 3 4		
Submerged Vegetation	● 1 2 3 4		

Buffer Plot 3	Canopy Type: ○ (D) ● (E) Leaf Type: ○ (B) ○ (N)	Absent: ●	Flag
Big Trees (>0.3m DBH)	● 1 2 3 4		
Small Trees (<0.3m DBH)	● 1 2 3 4		
Woody Shrubs, Saplings (0.5m-5m HIGH)	0 1 2 ● 4		
Woody Shrubs, Saplings (<0.5m HIGH)	0 ● 2 3 4		
Herbs, Forbs and Grasses	0 1 2 3 ●		
Bare ground	● 1 2 3 4		
Litter, duff	0 ● 2 3 4		
Rock	● 1 2 3 4		
Water	● 1 2 3 4		
Submerged Vegetation	● 1 2 3 4		

Stressor Presence/Absence - Confirm that a filled data bubble indicates presence and an unfilled bubble indicates absence by filling this bubble. ●

Residential and Urban Stressors

Fill bubble if present - Plot	1	2	3	Flag
Road - gravel	○	○	○	
Road - two lane	○	○	○	
Road - four lane	○	○	○	
Parking Lot/Pavement	○	○	○	
Golf Course	○	○	○	
Lawn/Park	○	○	○	
Suburban Residential	○	○	○	
Urban/Multifamily	○	○	○	
Landfill	○	○	○	
Dumping	○	○	○	
Trash	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Hydrology Stressors

Fill bubble if present - Plot	1	2	3	Flag
Ditches, Channelization	○	○	○	
Dike/Dam/Road/RR Bed (IMPEDE FLOW)	○	○	○	
Water Level Control Structure	○	○	○	
Excavation, Dredging	○	○	○	
Fill/Spoil Banks	○	○	○	
Freshly Deposited Sediment (UNVEGETATED)	○	○	○	
Soil Loss/Root Exposure	○	○	○	
Wall/Riprap	○	○	○	
Inlets, Outlets	○	○	○	
Point Source/Pipe (EFFLUENT OR STORMWATER)	○	○	○	
Impervious surface input (SHEETFLOW)	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Agricultural & Rural Stressors

Fill bubble if present - Plot	1	2	3	Flag
Pasture/Hay	○	○	○	
Range	○	○	○	
Row Crops	○	○	○	
Fallow Field (RECENT-RESTING ROW CROP FIELD)	○	○	○	
Fallow Field (OLD - GRASS, SHRUBS, TREES)	○	○	○	
Nursery	○	○	○	
Dairy	○	○	○	
Orchard	○	○	○	
Confined Animal Feeding	○	○	○	
Rural Residential	○	○	○	
Gravel Pit	○	○	○	
Irrigation	○	○	○	
Other: _____	○	○	○	

Industrial Development Stressors

Fill bubble if present - Plot	1	2	3	Flag
Oil Drilling	○	○	○	
Gas Wells	○	○	○	
Mine (surface)	○	○	○	
Mine (underground)	○	○	○	
Military	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Habitat/Vegetation Stressors

Fill bubble if present - Plot	1	2	3	Flag
Forest Clear Cut	○	○	○	
Forest Selective Cut	○	○	○	
Tree Plantation	○	○	○	
Tree Canopy Herbivory (INSECT)	○	○	○	
Shrub Layer Browsed (WILD OR DOMESTIC)	●	●	●	
Highly Grazed Grasses (OVERALL <3" HIGH)	○	○	○	
Recently Burned Forest Canopy	○	○	○	
Recently Burned Grassland (BLACKENED)	○	○	○	

Fill bubble if present - Plot	1	2	3	Flag
Herbicide Use	○	●	○	i
Mowing/Shrub Cutting	○	○	○	
Trails	○	○	○	
Soil Compaction (ANIMAL OR HUMAN)	○	○	○	
Offroad vehicle damage	○	○	○	
Soil erosion (FROM WIND, WATER, OR OVERUSE)	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Flag codes: K = No measurement made, U = Suspect measurement., F1,F2, etc. = misc. flags assigned by each field crew.
Explain all flags in comment section on the back of this form

2428168304

Lonicera all MR bert, Hop hack berb teasel

FORM B-1: BUFFER SAMPLE PLOTS - TARGETED ALIEN SPECIES (Back)

Reviewed by (initial): _____

Site ID: PCAPHi1389 **DATE:** 08 / 19 / 2013

● Confirm a filled data bubble indicates presence and an unfilled bubble indicates absence by filling in this bubble

Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag
Eurasian Watermilfoil	○	○	○		Purple Loosestrife	○	○	○		Johnson Grass	○	○	○	
Water hyacinth	○	○	○		Knotweed	○	○	○		Kudzu	○	○	○	
Yellow Floating Heart	○	○	○		Japanese Knotweed	○	○	○		*Multiflora Rose*	○	●	○	
Giant Salvinia	○	○	○		Perennial Pepperweed	○	○	○		Common Buckthorn	○	○	○	
Garlic Mustard	○	●	○		Giant Reed	○	○	○		Himalayan Blackberry	○	○	○	
Poison Hemlock	○	○	○		Cheatgrass	○	○	○		Tamarisk	○	○	○	
Mile-A-Minute Weed	○	○	○		Reed Canary Grass	○	○	○		Other: _____	○	○	○	
Birdsfoot Trefoil	○	○	○		Common Reed	○	○	○		Other: _____	○	○	○	
Canada Thistle	○	○	○		Leafy Spurge	○	○	○		Other: _____	○	○	○	
										Other: _____	○	○	○	

PLOT COORDINATES

Provide GPS coordinates at the center of the Buffer Plot (#3) at the far end of each Buffer Transect and for the Buffer Plot at the AA CENTER. Indicate the location of the plot coordinates by filling in the appropriate bubble.

If Buffer Plot 3 can not be accessed, take the coordinates at the nearest practicable location ALONG THE TRANSECT. This is important because all Buffer Plots are centered on the Buffer Transects and the coordinates will indicate the location of the transect. Fill in the "nearest practicable location" bubble, fill in the flag box, and describe where the coordinates were taken and why in the comment section below. The coordinates of the nearest practicable location can be either placed as close to the center of Plot 3 as possible or at the center of the last accessible Buffer Plot.

Location of coordinates (choose one): Flag

○ AA CENTER ○ N3 ○ S3 ● E3 ○ W3 ○ Nearest practicable location (flag and comment below)

Latitude North 41.22547 Longitude West 081.70390

Use Decimal Degrees; NAD83

Flag	Comments
1	Clear herbicide use, possibly more than a year old on Lonicera morrowii.

FORM B-1: BUFFER SAMPLE PLOTS (Front)

Reviewed by (initial): _____

Site ID: PCAP Hi 1389 DATE: 08 / 19 / 2015

Location: ○ AA Center ○ N ● S ○ E ○ W

Fill in bubble(s) if plot(s) could not be sampled and flag → ○ Plot 1 ○ Plot 2 ○ Plot 3

Buffer Natural Cover Strata

Fill in bubbles for all that apply: Canopy Type: D = Deciduous; E = Evergreen. Leaf Type: B = Broadleaf; N = Needle Leaf. Absent: No tree canopy.
Strata Section: Fill in appropriate cover class bubble for each strata type for each plot. 0 = Absent; 1 = Sparse(<10%); 2=Moderate(10-40%); 3 = Heavy (40-75%); 4 = Very Heavy (>75%)

Buffer Plot 1 (Canopy Type: D L; Leaf Type: B N; Absent: ●; Flag)		Buffer Plot 2 (Canopy Type: ● E; Leaf Type: ● N; Absent: ○; Flag)		Buffer Plot 3 (Canopy Type: ● E; Leaf Type: ● N; Absent: ○; Flag)	
Big Trees (>0.3m DBH)	●	Big Trees (>0.3m DBH)	●	Big Trees (>0.3m DBH)	●
Small Trees (<0.3m DBH)	●	Small Trees (<0.3m DBH)	●	Small Trees (<0.3m DBH)	●
Woody Shrubs, Saplings (0.5m-5m HIGH)	●	Woody Shrubs, Saplings (0.5m-5m HIGH)	●	Woody Shrubs, Saplings (0.5m-5m HIGH)	●
Woody Shrubs, Saplings (<0.5m HIGH)	●	Woody Shrubs, Saplings (<0.5m HIGH)	●	Woody Shrubs, Saplings (<0.5m HIGH)	●
Herbs, Forbs and Grasses	●	Herbs, Forbs and Grasses	●	Herbs, Forbs and Grasses	●
Bare ground	●	Bare ground	●	Bare ground	●
Litter, duff	●	Litter, duff	●	Litter, duff	●
Rock	●	Rock	●	Rock	●
Water	●	Water	●	Water	●
Submerged Vegetation	●	Submerged Vegetation	●	Submerged Vegetation	●

Stressor Presence/Absence - Confirm that a filled data bubble indicates presence and an unfilled bubble indicates absence by filling this bubble. ●

Residential and Urban Stressors

Fill bubble if present - Plot	1	2	3	Flag
Road - gravel	○	○	○	
Road - two lane	○	○	○	
Road - four lane	○	○	○	
Parking Lot/Pavement	○	○	○	
Golf Course	○	○	○	
Lawn/Park	○	○	○	
Suburban Residential	○	○	○	
Urban/Multifamily	○	○	○	
Landfill	○	○	○	
Dumping	○	○	○	
Trash	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Hydrology Stressors

Fill bubble if present - Plot	1	2	3	Flag
Ditches, Channelization	○	○	○	
Dike/Dam/Road/RR Bed (IMPEDE FLOW)	○	○	○	
Water Level Control Structure	○	○	○	
Excavation, Dredging	○	○	○	
Fill/Spoil Banks	○	○	○	
Freshly Deposited Sediment (UNVEGETATED)	○	○	○	
Soil Loss/Root Exposure	○	○	○	
Wall/Riprap	○	○	○	
Inlets, Outlets	○	○	○	
Point Source/Pipe (EFFLUENT OR STORMWATER)	○	○	○	
Impervious surface input (SHEETFLOW)	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Agricultural & Rural Stressors

Fill bubble if present - Plot	1	2	3	Flag
Pasture/Hay	○	○	○	
Range	○	○	○	
Row Crops	○	○	○	
Fallow Field (RECENT-RESTING ROW CROP FIELD)	○	○	○	
Fallow Field (OLD - GRASS, SHRUBS, TREES)	○	○	○	
Nursery	○	○	○	
Dairy	○	○	○	
Orchard	○	○	○	
Confined Animal Feeding	○	○	○	
Rural Residential	○	○	○	
Gravel Pit	○	○	○	
Irrigation	○	○	○	
Other: _____	○	○	○	

Industrial Development Stressors

Fill bubble if present - Plot	1	2	3	Flag
Oil Drilling	○	○	○	
Gas Wells	○	○	○	
Mine (surface)	○	○	○	
Mine (underground)	○	○	○	
Military	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Habitat/Vegetation Stressors

Fill bubble if present - Plot	1	2	3	Flag
Forest Clear Cut	○	○	○	
Forest Selective Cut	○	○	○	
Tree Plantation	○	○	○	
Tree Canopy Herbivory (INSECT)	○	○	○	
Shrub Layer Browsed (WILD OR DOMESTIC)	●	●	●	
Highly Grazed Grasses (OVERALL <3" HIGH)	○	○	○	
Recently Burned Forest Canopy	○	○	○	
Recently Burned Grassland (BLACKENED)	○	○	○	

Fill bubble if present - Plot	1	2	3	Flag
Herbicide Use	○	○	○	
Mowing/Shrub Cutting	○	○	○	
Trails	○	○	○	
Soil Compaction (ANIMAL OR HUMAN)	○	○	○	
Offroad vehicle damage	○	○	○	
Soil erosion (FROM WIND, WATER, OR OVERUSE)	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Flag codes: K = No measurement made, U = Suspect measurement., F1,F2, etc. = misc. flags assigned by each field crew. Explain all flags in comment section on the back of this form

2428168304

FORM B-1: BUFFER SAMPLE PLOTS - TARGETED ALIEN SPECIES (Back)

Reviewed by (initial): _____

Site ID: PCAP Hi 1389 **DATE:** 08 / 19 / 2013

Confirm a filled data bubble indicates presence and an unfilled bubble indicates absence by filling in this bubble

Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag
Eurasian Watermilfoil	○	○	○		Purple Loosestrife	○	○	○		Johnson Grass	○	○	○	
Water hyacinth	○	○	○		Knotweed	○	○	○		Kudzu	○	○	○	
Yellow Floating Heart	○	○	○		Japanese Knotweed	○	○	○		Multiflora Rose	○	○	○	
Giant Salvinia	○	○	○		Perennial Pepperweed	○	○	○		Common Buckthorn	○	○	○	
Garlic Mustard	○	●	○		Giant Reed	○	○	○		Himalayan Blackberry	○	○	○	
Poison Hemlock	○	○	○		Cheatgrass	○	○	○		Tamarisk	○	○	○	
Mile-A-Minute Weed	○	○	○		Reed Canary Grass	○	○	○		Other: _____	○	○	○	
Birdsfoot Trefoil	○	○	○		Common Reed	○	○	○		Other: _____	○	○	○	
Canada Thistle	○	○	○		Leafy Spurge	○	○	○		Other: _____	○	○	○	
										Other: _____	○	○	○	

PLOT COORDINATES

Provide GPS coordinates at the center of the Buffer Plot (#3) at the far end of each Buffer Transect and for the Buffer Plot at the AA CENTER. Indicate the location of the plot coordinates by filling in the appropriate bubble.

If Buffer Plot 3 can not be accessed, take the coordinates at the nearest practicable location ALONG THE TRANSECT. This is important because all Buffer Plots are centered on the Buffer Transects and the coordinates will indicate the location of the transect. Fill in the "nearest practicable location" bubble, fill in the flag box, and describe where the coordinates were taken and why in the comment section below. The coordinates of the nearest practicable location can be either placed as close to the center of Plot 3 as possible or at the center of the last accessible Buffer Plot.

Location of coordinates (choose one):

○ AA CENTER ○ N3 ● S3 ○ E3 ○ W3 ○ Nearest practicable location (flag and comment below)

Flag: ☐

Latitude North 41 . 22430 Longitude West 081 . 70470

Use Decimal Degrees; NAD83

Flag	Comments

FORM B-1: BUFFER SAMPLE PLOTS (Front)

Reviewed by (initial): _____

Site ID: PCAP Hi[illegible] 1389

DATE: 08/19/2013

Location: ○ AA Center ○ N ○ S ○ E ● W

Fill in bubble(s) if plot(s) could not be sampled and flag → ○ Plot 1 ○ Plot 2 ○ Plot 3

Buffer Natural Cover Strata

Fill in bubbles for all that apply: Canopy Type: D = Deciduous; E = Evergreen. Leaf Type: B = Broadleaf; N = Needle Leaf. Absent: No tree canopy.
Strata Section: Fill in appropriate cover class bubble for each strata type for each plot. 0 = Absent; 1 = Sparse(<10%); 2=Moderate(10-40%); 3 = Heavy (40-75%); 4 = Very Heavy (>75%)

Buffer Plot 1 (Canopy Type: D ● E; Leaf Type: B ● N; Absent: ○; Flag)	0	1	2	3	4	Flag
Big Trees (>0.3m DBH)	○	○	○	●	○	
Small Trees (<0.3m DBH)	○	○	●	○	○	
Woody Shrubs, Saplings (0.5m-5m HIGH)	○	○	○	●	○	
Woody Shrubs, Saplings (<0.5m HIGH)	○	●	○	○	○	
Herbs, Forbs and Grasses	○	●	○	○	○	
Bare ground	○	●	○	○	○	
Litter, duff	○	○	○	○	●	
Rock	●	●	○	○	○	
Water	●	○	○	○	○	
Submerged Vegetation	●	○	○	○	○	

Buffer Plot 2 (Canopy Type: D ● E; Leaf Type: B ● N; Absent: ○; Flag)	0	1	2	3	4	Flag
Big Trees (>0.3m DBH)	○	○	●	○	○	
Small Trees (<0.3m DBH)	○	○	●	○	○	
Woody Shrubs, Saplings (0.5m-5m HIGH)	○	○	○	●	○	
Woody Shrubs, Saplings (<0.5m HIGH)	○	○	○	●	○	
Herbs, Forbs and Grasses	○	○	●	○	○	
Bare ground	●	○	○	○	○	
Litter, duff	○	○	○	●	○	
Rock	○	●	○	○	○	
Water	●	○	○	○	○	
Submerged Vegetation	●	○	○	○	○	

Buffer Plot 3 (Canopy Type: D ● E; Leaf Type: B ● N; Absent: ○; Flag)	0	1	2	3	4	Flag
Big Trees (>0.3m DBH)	●	○	○	○	○	
Small Trees (<0.3m DBH)	○	○	○	●	○	
Woody Shrubs, Saplings (0.5m-5m HIGH)	○	○	●	○	○	
Woody Shrubs, Saplings (<0.5m HIGH)	○	○	○	●	○	
Herbs, Forbs and Grasses	○	○	●	○	○	
Bare ground	●	●	○	○	○	
Litter, duff	○	●	●	○	○	
Rock	●	○	○	○	○	
Water	●	○	○	○	○	
Submerged Vegetation	●	○	○	○	○	

Stressor Presence/Absence - Confirm that a filled data bubble indicates presence and an unfilled bubble indicates absence by filling this bubble. ●

Residential and Urban Stressors

Fill bubble if present - Plot	1	2	3	Flag
Road - gravel	○	○	○	
Road - two lane	○	○	○	
Road - four lane	○	○	○	
Parking Lot/Pavement	○	○	○	
Golf Course	○	○	○	
Lawn/Park	○	○	○	
Suburban Residential	○	○	○	
Urban/Multifamily	○	○	○	
Landfill	○	○	○	
Dumping	○	○	○	
Trash	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Hydrology Stressors

Fill bubble if present - Plot	1	2	3	Flag
Ditches, Channelization	●	○	○	\
Dike/Dam/Road/RR Bed (IMPEDE FLOW)	○	○	○	
Water Level Control Structure	○	○	○	
Excavation, Dredging	○	○	○	
Fill/Spoil Banks	○	○	○	
Freshly Deposited Sediment (UNVEGETATED)	○	○	○	
Soil Loss/Root Exposure	○	○	○	
Wall/Riprap	○	○	○	
Inlets, Outlets	○	○	○	
Point Source/Pipe (EFFLUENT OR STORMWATER)	○	○	○	
Impervious surface input (SHEETFLOW)	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Agricultural & Rural Stressors

Fill bubble if present - Plot	1	2	3	Flag
Pasture/Hay	○	○	○	
Range	○	○	○	
Row Crops	○	○	○	
Fallow Field (RECENT-RESTING ROW CROP FIELD)	○	○	○	
Fallow Field (OLD - GRASS, SHRUBS, TREES)	○	○	○	
Nursery	○	○	○	
Dairy	○	○	○	
Orchard	○	○	○	
Confined Animal Feeding	○	○	○	
Rural Residential	○	○	○	
Gravel Pit	○	○	○	
Irrigation	○	○	○	
Other: _____	○	○	○	

Industrial Development Stressors

Fill bubble if present - Plot	1	2	3	Flag
Oil Drilling	○	○	○	
Gas Wells	○	○	○	
Mine (surface)	○	○	○	
Mine (underground)	○	○	○	
Military	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Habitat/Vegetation Stressors

Fill bubble if present - Plot	1	2	3	Flag
Forest Clear Cut	○	○	○	
Forest Selective Cut	○	○	○	
Tree Plantation	○	○	○	
Tree Canopy Herbivory (INSECT)	○	○	○	
Shrub Layer Browsed (WILD OR DOMESTIC)	●	●	●	
Highly Grazed Grasses (OVERALL <3" HIGH)	○	○	○	
Recently Burned Forest Canopy	○	○	○	
Recently Burned Grassland (BLACKENED)	○	○	○	

Fill bubble if present - Plot	1	2	3	Flag
Herbicide Use	○	●	○	
Mowing/Shrub Cutting	○	○	○	
Trails	○	○	○	
Soil Compaction (ANIMAL OR HUMAN)	○	○	○	
Offroad vehicle damage	○	○	○	
Soil erosion (FROM WIND, WATER, OR OVERUSE)	●	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Flag codes: K = No measurement made, U = Suspect measurement., F1,F2, etc. = misc. flags assigned by each field crew.
Explain all flags in comment section on the back of this form

2428168304

rose, morrowii, privet, maackii, garlic, berberis

FORM B-1: BUFFER SAMPLE PLOTS - TARGETED ALIEN SPECIES (Back)

Reviewed by (initial): _____

Site ID: PCAP Hi 1389 DATE: 08/19/2013

● Confirm a filled data bubble indicates presence and an unfilled bubble indicates absence by filling in this bubble

Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag
Eurasian Watermilfoil	○	○	○		Purple Loosestrife	○	○	○		Johnson Grass	○	○	○	
Water hyacinth	○	○	○		Knotweed	○	○	○		Kudzu	○	○	○	
Yellow Floating Heart	○	○	○		Japanese Knotweed	○	○	○		Multiflora Rose	○	●	●	
Giant Salvinia	○	○	○		Perennial Pepperweed	○	○	○		Common Buckthorn	○	○	○	
Garlic Mustard	○	○	○		Giant Reed	○	○	○		Himalayan Blackberry	○	○	○	
Poison Hemlock	○	○	○		Cheatgrass	○	○	○		Tamarisk	○	○	○	
Mile-A-Minute Weed	○	○	○		Reed Canary Grass	○	○	○		Other: _____	○	○	○	
Birdsfoot Trefoil	○	○	○		Common Reed	○	○	○		Other: _____	○	○	○	
Canada Thistle	○	○	○		Leafy Spurge	○	○	○		Other: _____	○	○	○	
										Other: _____	○	○	○	

PLOT COORDINATES

Provide GPS coordinates at the center of the Buffer Plot (#3) at the far end of each Buffer Transect and for the Buffer Plot at the AA CENTER. Indicate the location of the plot coordinates by filling in the appropriate bubble.

If Buffer Plot 3 can not be accessed, take the coordinates at the nearest practicable location ALONG THE TRANSECT. This is important because all Buffer Plots are centered on the Buffer Transects and the coordinates will indicate the location of the transect. Fill in the "nearest practicable location" bubble, fill in the flag box, and describe where the coordinates were taken and why in the comment section below. The coordinates of the nearest practicable location can be either placed as close to the center of Plot 3 as possible or at the center of the last accessible Buffer Plot.

Location of coordinates (choose one):

○ AA CENTER ○ N3 ○ S3 ○ E3 ● W3 ○ Nearest practicable location (flag and comment below)

Flag: ☐

Latitude North 41.22496 Longitude West 081.70698

Use Decimal Degrees; NAD83

Flag	Comments
1	Ditch runs along road.