CLEVELAND METROPARKS Plant Community Assessment Program: Quality Control Form Cleveland Metroparks

Project Label: PCAP **Plot No:** 1201 **Date Sampled:** 6-20-2012 **Lead:** Eyzenbach

			Comment required if item answer is NO
Parking/Access outside of Park Boundaries:		Y (N)	If yes, write details in Comments section below
Field journals completed		(Y) N	
Site sketch made on 1:3000 map?		(Y) N	
Check cover page	X-axis Bearing of plot recorded	(Y) N	
Check Ash -	GPS coords. Recorded	(Y) N	
	North direction recorded	(Y) N	
	Photographs taken?	(Y) N	
Plot No., Date agreement on all pages?		(Y) N	
Header data completed all pages?		(Y) N	
Cover classes recorded in all Intensive modules		(Y) N	
Browse Level By Species		(Y) N	
Woody stem quality control check		(Y) N	
Invasive plant quality control check		(Y) N	
Ash trees mapped		(Y) N	
Cover by Strata? (confirm cover type)		(Y) N	
Soil samples collected with matching plot #.		(Y) N	
Vouchers labeled on datasheet with initials and number		(Y) N	
Vouchers labeled on collection bag		(Y) N	
Pink flags removed		(Y) N	
Data sheet QA before leaving site?		(Y) N	
Common equipment returned to tub.		Y N	
Data sheets scanned?		6/21/12	Enter date to left NB
Final data sheets scanned?			Enter date to left
Buffer Widths measured?		(Y) N	JTP 6-22-12
Web Soil Survey		(Y) N	By 6/21/2012
Voucher Location	Refrigerator	(Y) N	6-20-2012
(# vouchers collected)	Press (#)		Enter number to left
SRE 511-519	Drier	Y N	
	Identified	(Y) N	
	Mounted	Y N	
	Thrown away	Y N	

GRTS point verification: Is plot sampleable?	
☑ Yes	Original GRTS point is sampleable
☐ No	Original GRTS point lands in a non-sampleable area (fill in category below)
	☐ Point falls in a water (i.e. river, lake)
	☐ Managed mowed area (i.e. golf course, picnic area, right-of-way)
	☐ Paved area (i.e. parkinglot, road)
	☐ Unsafe to sample (i.e. steep slope)
	☐ Other

Additional Comments:

CLEVELAND METROPARKS Plant Community Assessment Program - Background Data Sheet

GENERAL INFORMATION

Project Label: PCAP

Project Name: 01Br2012

Plot Name: Where is the pine Stand?

Plot No.: 1201

☐ Level 4 (no nested corners sampled)

■ Level 5 (nested corners sampled)

Date (mm/dd/yyyy): 6/20/2012

End date (if > 1 day): / /

Party	Role**
S. Eysenbach	Plot leader
K. Lewis	Asst
A. Young	Asst

** Roles: Co-leader, Asst., Guide, Owner, Taxonomist, etc.

PLOT NOT SAMPLED: ☐ Other

☐ Perm. water ☐ Paved ☐ Slope ☐ Safety

SAMPLING QUALITY*

Effort Level:

✓ Very thorough

☐ Accurate

☐ Hurried

subjective evaluation of how much effort put into sampling. Hurried plots may still provide good data

TAXONOMIC ACCURACY

	high	modera.	low	not smpl
vascul.	✓			n/a
bryo			✓	
lichen				✓

TAXONOMIC STANDARD

Authority: G&C Pub Date: 1998

LOCATION

State: OH County: Cuyahoga

Quadrangle: *Not on map Northfield SE

Local Place Names: Meadows Picnic Area (near ballfield)

Landowner: CM

Data Confidentiality:

Check one: ☐ Public data ☐ Private Data

☐ Fuzz 100m ☐ Fuzz 250m ☐ Fuzz 500m

Reason:

If data not public why?

Source of coordinates ☐ MAP ■ GPS

Coordinate system:	Coord. Units
■ Lat/Long ☐ UTM ☐ StatePlane	■ deg ☐ deg min
☐ Other (*specify*)	■ m ☐ ft ☐

Datum: ■ NAD83/WGS84 ☐ NAD27

GPS location in plot x=0 to 5, y=-1,0,+1):

x = 0 y = 0 (base of plot x=0, y=0)

Latitude: 41.31461

Longitude: 081.61924

Coord. Accuracy: ✓ m ☐ ft +- 15

GPS File Name: 1201A

Plot size for cover data: 0.04 (hectares)

X-axis Bearing of plot: [108] °

Depth: (1-5): 4

Intensive modules: 2, 3, 8, 9 (EDIT IF MODIFIED)

Camera No.: 2

Photo Nos.: C2-1688-1689

Plot placement: ~~GRTS~~ ☐ Representative

☐ Random ☐ Stratified Random ☐ Transect component

☐ Systematic (grid) ☐ Capture specific feature ☐ Other



NOTES: Include **Layout** (any unusual shape details), **Location** (directions and landscape context), **Rationale** (why here), and **Veg Characterization** (description of community, dominants, strata; BROWSE). Additional notes in space on back.

Meadows Drive

Layout: 1 x 4 Narrow disturbed area w/pine mixed in

Location: Park at Meadows Picnic Area near the 1st ball field. Plot is located south of the ball field and east of the entrance drive

Rationale: GRTS pt. GPS was acting goofy (had pt in mowed area) used map to determine pt.

Veg Char:

Canopy: Variable - Red Maple, Sugar Maple, Red Oak, American Elm, Ash White Oak

Shrub: Bush honeysuckle, Privet, Spice Bush, Hawthorn

Herb: Cinquefoil, Virginia Creeper, Ash seedlings Poison ivy, Orchard Grass

OVER

Minimum required fields in Bold and Underlined

*Definitions and values in CM PCAP FOM v. 1.0 and CVS Field Guide

CLEVELAND METROPARKS Plant Community Assessment Program - Background Data Sheet

Project Label: PCAP Project Name: 01Br2012 Plot No.: 1201

MODIFIED NATURESERVE CLASS*

CODE (on separate form): O Fit= fair Conf= Low

COMMUNITY NAME: Mixed

*~~Mag~~ Very strange mix in canopy but not enough pines to call it atypical planted

HOMOGENEITY

- ☑ Homogeneous
- ☐ Compositional trend across the plot
- ☐ Conspicuous inclusions
- ☐ Irregular/pattern mosaic

DISTURBANCES

type*	severity**	yrs ago	% of plot	description
Human	L	±5	100	trash
Natural				
Fire				
Cut				
Animal	MH	0	100	Deer Browse
Other				

**L=low, ML=med low, M=med, MH=med high, H=high, VH=very high

Current Land Use: Park

Former Land Use: UNK

HYDROLOGIC REGIME*

- ☑ Upland (seldom flooded)
- ☐ Intermittently/seasonally saturated (seldom flooded)
- ☐ Permanently/Semipermanent. saturated (dry <1/yr, seldom flooded)
- ☐ Occasionally flooded (<1/yr)
- ☐ Temporarily flooded
- ☐ Intermittently flooded
- ☐ Semipermanently flooded
- ☐ Permanently flooded
- ☐ Tidal/Seiche flooded daily
- ☐ Tidal/Seiche flooded monthly
- ☐ Tidal/Seiche flooded irregular (e.g. wind, storms)
- ☐ Unknown

SALINITY*

- ☐ Saltwater
- ☐ Brackish
- ☐ Fresh
- ☑ Upland (n/a)

(by default unless plot is a wetland)

Additional notes & diagrams: (Representativeness of plot to the stand, successional status, maturity, etc.)

Plot was set up because only 10 m would stretch across the narrow disturbed forest. We could only go 40 meters before a more mature less disturbed forest emerged. This is a highly disturbed edge plot with a variable canopy.

There was lots of evidence of browse on the Asters, Fraxinus seedlings, Lonicera. No evident deer trail.

Tons of wood chips in Mod 3

Lots of invasives and ruderals or disturbance species (prunella, taraxacum, dactylus, holcus) Ash appear to be declining. Mod 4 was more open due to dead Juglans and failing Crateagus

CLEVELAND METROPARKS Plant Community Assessment Program Species Cover Data Sheet 2a Page 1 of 4

Project Label: PCAP Project name: 01Br2012 Plot no.: 1201

Total modules: 4 Intensive modules: 4 Plot configuration: 1 x 4 Plot area (ha): 0.1

Cleveland Metroparks

Br = Browse Level. Use cover classes to describe amount of browse per species over entire plot

Strata - Cov. entire plot

Estimate for each intensive module:

	mod 1	corner 4	mod 1	corner 2	mod 2	corner 4	mod 2	corner 2	mod 3	corner 4	mod 3	corner 2	mod 4	corner 4	mod 4	corner 2	mod R	corner R
	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov
%open water	1	0			1	0			1	0			1	0				
%unvegetated open water	1	0			1	0			1	0			1	0				
%unveg. ground (bare soil)	1	5			1	3			1	2			1	2				
%unveg. litter (bare litter)	1	5			1	2			1	3			1	4				

T	S	H	(F)	(A)	Br	Species	c	Voucher #	mod 1 depth	mod 1 cov	corner 4 depth	corner 4 cov	mod 2 depth	mod 2 cov	corner 2 depth	corner 2 cov	mod 3 depth	mod 3 cov	corner 4 depth	corner 4 cov	mod 3 depth	mod 3 cov	corner 2 depth	corner 2 cov	mod 4 depth	mod 4 cov	corner 4 depth	corner 4 cov	mod R depth	mod R cov	corner R depth	corner R cov
6	5					Acer saccharum			2	6			4	4	4	+1	4								1	4						
6	5					Acer rubrum					3	6			4	6				4	4	6			1	2						
	2	1				Cornus florida			1	3					2	1																
	3	2			8	Lonicera maackii			1	2			X	X			1	1							1	2						
	5	2			8	Ligustrum vulgare			4	6			1	3			1	2					-1	4								
	5	2			9	Hamamelis virginiana			3	5	3				2	2	3	1														
	6	2			8	Rosa multiflora			2	4	2		3	6			4	6	4				1	3								
		4			9	Fraxinus sp. (seedlings)			4	2	3		3	2	3		2	2	2		2	3	4									
		2			8	Ulmus sp. (seedlings)			3	2	2				3	2	1	2			2	2	2									
	6	2			7	Crataegus sp.			1	2			1	2			2	4			2	5	4									
		2				Prunus serotina			1	2			2	5																		
		2				Taraxacum officinale			4	2	2		3	2	2		1	2			1	1										
		2				Dactylus glomerata			1	2			1	1													2	3				
	1	6			3	Parthenocissus quinquefolia		see 8-23-12	3	2	3		3	2	4		3	3	4		2	2	4									
		3			9	Aster ~~Solidago~~ lateriflorus		C2-1691	4	2	3		4	2	3		2	2	2		2	2	2									
	6	2			5	Lindera benzoin			1	1			1	2		-1	6		-				2	2								
	6	2			6	Lonicera morrowii			2	3	4		3	5		-			2	4	2	2										
		1				Berberis thunbergii			1	2																						
		2				Solidago caesia					2	2	2	2			1	1														
		2				Moss sp.			2	2	2		1	2			1	2					1	2								
4		4				Pinus nigra			1	4			5		5																	
2						Quercus palustris			4	2																						
6						Quercus rubra			4	3	4		4	7	2	-1	3															
4	2	2			6	Prunus cerasus			2	4	2		4	2			3	3					-1	3								
		2				Circaea lutetiana			4	2							-1	2														

EXAMPLES OF PERCENT OF AREA COVERED

The following graphic can be used for various data elements to convey "Amount" or "Quantity". *NOTE*. Within any given box, each quadrant contains the same total area covered, just different sized objects.



cover class	% cover	midpoint
1	solitary or few	0.0001
2	0-1%	0.005
3	1-2%	0.015
4	2-5%	0.035
5	5-10%	0.075
6	10-25%	0.175
7	25-50%	0.375
8	50-75%	0.625
9	75-95%	0.850
10	95-100%	0.975







BROWSE RATING NARRATIVE DESCRIPTION

LOW OR NONE: there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.

MEDIUM LOW values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

MEDIUM: browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.

MEDIUM HIGH values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.

HIGH: greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.

VERY HIGH values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.

CLEVELAND METROPARKS Plant Community Assessment Program Species Cover Data Sheet 2a Page 2 of 4

Project Label: PCAP Project name: 01Br2012 Plot no.: 1201

Total modules: 4 Intensive modules: 4 Plot configuration: 1x4 Plot area (ha): 0.1

Cleveland Metroparks

Br = Browse Level. Use cover classes to describe amount of browse per species over entire plot

Strata - Cov. entire plot

Estimate for each intensive module:

	mod 1	corner 4	mod 1	corner 2	mod 2	corner 4	mod 2	corner 2	mod 3	corner 4	mod 3	corner 2	mod 4	corner 4	mod 4	corner 2	mod R	corner R
	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov
%open water	1				1				1				1					
%unvegetated open water	1				1				1				1					
%unveg. ground (bare soil)	1				1				1				1					
%unveg. litter (bare litter)	1				1				1				1					

T	S	H	(F)	(A)	Br	Species	c	Voucher #	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov
		2				Prunella vulgaris			4	2	2				4	2							1	2				
		4				Juncus tenuis	X	SRE 511	3	2			4	2			-1	3					4	3				
		1				Antroxanthum odoratum	X	SRE 512	3	2																		
2		2				Toxicodendron radicans			3	2			4	2			4	2	4	2								
		3				Carex sp. blanda ~~laxiculmis~~	X	see 520 C2-1690	3	2							1	2			-1	2						
		1				Veronica serphyllifolia			3	2																		
		2				Oxalis stricta			2	1	1	2																
		2			3	Cornus sp			3	2													1	2				
		1				Plantago sp			2	1	~~1~~		4	1	1	1							1	1				
		2				Arisaema triphyllum ssp triphyllum			2	2	3		1	2			3	2	2				3	2				
		1				Daucus carota			2	1							1	1										
		3			6	Rhamnus frangula			1	2					2	2	-1	2					2	2				
		2				Rubus sp. (no fruit or flower)					3	2	3	2			2	2							4			
		6				Carex rosea	X	SRE 513			2	5	1	2			1	3					4	3	3			
		2				Rubus sp. (no repro)					2	2	2	2			2	2			-				4	2		
	6	2				Carpinus caroliana					2	2							4	5	-				2	5		
		2			6	Carya seedling (hairy)					2	2			2	2	2	2			-1	2						
		6				Potentilla simplex			1	2			3	7	4		2	2					3	3	3			
		1				Vitis sp (seedlings)			1	2					3	2	1	2										
		2				Viburnum opulus var. opulus			1	1	2	2							3	2	1	2						
		2				Elymus hystrix			1	2																		
		2				Quercus seedling			1	1					3	2	1	1			-1	2						
		2				Carex cristatella SRE 6-22-2012	X	SRE 514	1	2	1	2																
6						Carya glabra			1	3									3	4	~~4~~	5						
6						Carya ovata			1	3											4	5						

EXAMPLES OF PERCENT OF AREA COVERED

The following graphic can be used for various data elements to convey "Amount" or "Quantity". *NOTE*: Within any given box, each quadrant contains the same total area covered, just different sized objects.



cover class	% cover	midpoint
1	solitary or few	0.0001
2	0-1%	0.005
3	1-2%	0.015
4	2-5%	0.035
5	5-10%	0.075
6	10-25%	0.175
7	25-50%	0.375
8	50-75%	0.625
9	75-95%	0.850
10	95-100%	0.975







BROWSE RATING NARRATIVE DESCRIPTION

LOW OR NONE: there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.

MEDIUM LOW values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

MEDIUM: browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.

MEDIUM HIGH values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.

HIGH: greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.

VERY HIGH values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.



CLEVELAND METROPARKS Plant Community Assessment Program Species Cover Data Sheet 2a Page 3 of 4

Project Label: PCAP Project name: 013r 2012 Plot no.: 1201

Total modules: 4 Intensive modules: 4 Plot configuration: 1x4 Plot area (ha): 0.1

Cleveland Metroparks

Br = Browse Level. Use cover classes to describe amount of browse per species over entire plot

Strata - Cov. entire plot

Estimate for each intensive module:
%open water
%unvegetated open water
%unveg. ground (bare soil)
%unveg. litter (bare litter)

T	S	H	(F)	(A)	Br	Species	c	Voucher #	mod 1 depth	mod 1 cov	corner 4 depth	corner 4 cov	mod 1 depth	mod 1 cov	corner 2 depth	corner 2 cov	mod 2 depth	mod 2 cov	corner 4 depth	corner 4 cov	mod 2 depth	mod 2 cov	corner 2 depth	corner 2 cov	mod 3 depth	mod 3 cov	corner 4 depth	corner 4 cov	mod 3 depth	mod 3 cov	corner 4 depth	corner 4 cov	mod R depth	mod R cov	corner R depth	corner R cov
		1				Acer seedling											3	1																		
5						~~Robinia pseudoacacia~~		Gleditsia triacanthos										4	6																		
		1				Veronica officinalis											2	1							1	2										
		2				Geum ~~sp.~~ canadense															3	2			1	2					1	2				
		6				Fragaria vesca var vesca NMZ 6/22/12															3	2			1	2					4	6	4			
7		1				Quercus alba											1	5							4	5					4	6				
		6				Poa trivialis	X	SRE 515									1	1							1	1					1	6				
		2				Barberea sp.											1	1							2	1					2	1				
		4				Rumex obtusifolia											1	1							1	2					1	2				
Poaceae		1				Danthonia spicata SRE 513 ~~SRE 513~~	X	SRE 516									1	1																		
3	3				8	Fraxinus pennsylvanica			1	1							1	2															4	4		
		2				Podophyllum peltatum		(sheesing)									1	2																		
	2					Fagus grandifolia											1	2																		
5						Pinus strobus											3	6	3												1	1				
6	4				9	Ulmus americana																			1	7					2	5				
6						Fraxinus sp. (tree)																			2	5					4	6				
		3				Carex gracillima	X	SRE 517																	1	1					1	3				
whitened par fuzzy		1				Scirpus atrovirens																			1	1										
		2			8	~~Unknown dicot~~ Epilobium parviflorum		SRE 1695-1697																	1	1					1	2				
		1				Dryopteris carthusiana																			1	1										
		2			8	Polygonum virginianum																			1	1					1	2				
		2				Leersia virginica																			1	1					4	2	2			
		1				Carex complanata	X	SRE 518																	1	1										
paryanoides		2				Carex cephalophora SRE 6-7-12	X	SRE 519																	1	1					2	2				
		1				Solidago flexicaulis																			1	1										

EXAMPLES OF PERCENT OF AREA COVERED

The following graphic can be used for various data elements to convey "Amount" or "Quantity". *NOTE*: Within any given box, each quadrant contains the same total area covered, just different sized objects.



cover class	% cover	midpoint
1	solitary or few	0.0001
2	0-1%	0.005
3	1-2%	0.015
4	2-5%	0.035
5	5-10%	0.075
6	10-25%	0.175
7	25-50%	0.375
8	50-75%	0.625
9	75-95%	0.850
10	95-100%	0.975







BROWSE RATING NARRATIVE DESCRIPTION

LOW OR NONE: there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.

MEDIUM LOW values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

MEDIUM: browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.

MEDIUM HIGH values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.

HIGH: greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.

VERY HIGH values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.



CLEVELAND METROPARKS Plant Community Assessment Program Species Cover Data Sheet 2a Page 4 of 4

Project Label: PCAP Project name: 01Br2012 Plot no.: 1201

Total modules: 4 Intensive modules: 4 Plot configuration: 1X4 Plot area (ha): 0.1

Cleveland Metroparks

Br = Browse Level. Use cover classes to describe amount of browse per species over entire plot

Strata - Cov. entire plot

Estimate for each intensive module:

	mod 1	corner 4	mod 1	corner 2	mod 2	corner 4	mod 2	corner 2	mod 3	corner 4	mod 3	corner 2	mod 4	corner 4	mod 4	corner 2	mod R	corner R
	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov
%open water	1				1				1				1					
%unvegetated open water	1				1				1				1					
%unveg. ground (bare soil)	1				1				1				1					
%unveg. litter (bare litter)	1				1				1				1					

T	S	H	(F)	(A)	Br	Species	c	Voucher #	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov	depth	cov
		1				Solanum dulcamara											1	2										
	2	2				Vitis sp.											*				-1	2						
		1				Viburnum dentatum															1	1						
		1				Epilobium coloratum															1	1						
		1				Rubus occidentalis															1	1						
		1			10	Juglans nigra															1	2						
		2				Hotrus lanatus															1	2						
5						Tilia americana															1	2						
		1				Sisrinchyium angustifolium SRE 8 22-12	X	SRE 521	1	1											X							

EXAMPLES OF PERCENT OF AREA COVERED

The following graphic can be used for various data elements to convey "Amount" or "Quantity". *NOTE*: Within any given box, each quadrant contains the same total area covered, just different sized objects.



cover class	% cover	midpoint
1	solitary or few	0.0001
2	0-1%	0.005
3	1-2%	0.015
4	2-5%	0.035
5	5-10%	0.075
6	10-25%	0.175
7	25-50%	0.375
8	50-75%	0.625
9	75-95%	0.850
10	95-100%	0.975







BROWSE RATING NARRATIVE DESCRIPTION

LOW OR NONE: there is no measurable browse line AND there are very few or no plants 1-m nested quadrat and intensive module. In general, low values relate to less than 10 percent, by numbers of stems browsed.

MEDIUM LOW values include evidence of browse at about 10 percent of the stems with no significant impact to plant reproduction evident. In this rating, plants are browsed but preferential species are observed to be reproducing in numbers that appear normal or near-normal in comparison to low browse areas. For example, trilliums may flower and fruit, but jewelweed and arrowwood viburnum exhibit browse.

MEDIUM: browse affects greater than 10 percent and less than 25 percent of stems in the 1 m2 nested quadrat and intensive module. A browse line is usually not evident or obvious for all classes and species of vegetation, but careful examination may show preferential browse and/or browse lines for some species of plants.

MEDIUM HIGH values include evidence of a browse line and 25 percent of stems browsed with very little vegetation regeneration evident. In this rating, for some species of plants, reproduction does not appear to occur or it is very severely limited.

HIGH: greater than 25 percent of the stems of plants in the 1 m2 nested quadrat and intensive module AND a browse line is evident.

VERY HIGH values include extensive browse conditions, where the browse line is very evident AND almost all seedlings and herbs are severely browsed or missing. Browse line may be 5 to 6 feet in height with no or little green growth beneath.



CLEVELAND METROPARKS Plant Community Assessment Program Natural Woody Stem Data Sheet

Project Label: PCAP Project Name: OIBR 2012 Plot No.: 1201 Page: 1 of 3

Cleveland Metroparks

Explain subsample (additional room on back):

mod #	species	c	voucher#	# stems 0-1.4m browsed	% sub or super sample	# shrub clumps	size class (cm) woody stems >1.4m 1 0-<1	2 1-<2.5	3 2.5-<5	4 5-<10	5 10 - <15	6 15 - <20	7 20 - <25	8 25 - <30	9 30 - <35	10 35 - <40	11 >40 (record each tree)
~~1~~	~~Acer nigrum~~							~~•~~									
1	Carya ovata						•										
1	Acer saccharum							••	•	••							
1	Cornus florida									•							
1	Pinus nigra														•		
1	Acer rubrum										•	••	•				
1	Standing dead							•			•		••				
1	Fraxinus pennsylvanica			••			•										
1	Prunus cerasus			••								•					
1	Hamamelis virginiana			••			•		••								
1	Lonicera morrowii					18											
1	Rosa multiflora					16											
1	Lonicera maackii					2											
1	Ligustrum vulgare					29											
1	Berberis thunbergi					3											
1	Crateagus sp.			•													
2	Acer saccharum							•	•	•							
2	Acer rubrum							•		•	••	••	••				
2	Pinus strobus											•	•			•	
2	Fraxinus pennsylvanica						•										
2	Quercus rubra									•							46.9
2	Gleditsia triacanthos														•		
~~2~~	~~Parthenoci~~																
2	Toxicodendron radicans						•										

DBH Measurement Rules



Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.

Record using the tally system from 1 to 10

1

2

3

4

5

ASH CANOPY CONDITION

1. **Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
2. **Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
3. **Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
4. **>50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
5. **Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.

A B C D E

ASH CANOPY BREAKUP CONDITION (for dead trees):
(if an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)

A: All main branches contain fine twigs (newly dead).
B: Over 50% of main branches have fine twigs.
C: Less than 50% of main branches have fine twigs.
D: Stem still standing and tertiary main branches present.
E: Central stem still standing.

CLEVELAND METROPARKS Plant Community Assessment Program Natural Woody Stem Data Sheet

Project Label: PCAP Project Name: 01BR2012 Plot No.: 1201 Page: 2 of 3

Cleveland Metroparks

Explain subsample (additional room on back):

mod #	species	c	voucher#	# stems 0-1.4m browsed	% sub or super sample	# shrub clumps	size class (cm) woody stems >1.4m 1 0-<1	2 1-<2.5	3 2.5-<5	4 5-<10	5 10 - <15	6 15 - <20	7 20 - <25	8 25 - <30	9 30 - <35	10 35 - <40	11 >40 (record each tree)
2	Quercus alba													••			
2	Crataegus sp.							•									
2	Prunus serotina											•	•				
2	Rosa multiflora					12											
2	Lonicera maackii					7											
2	Lonicera morrowii					7											
2	Ligustrum vulgare					2											
2	Fraxinus sp			3													
2	Carya sp.			32													
2	Lindera benzoin			1													
3	Parthenocissus quinquefolia						•• •										
3	Acer rubrum											• •			• •		
3	Quercus rubra									•							
3	Acer saccharum							•		•							
3	Fraxinus pensylvanica								•								
3	Prunus cerasus								•								
3	Ulmus americana								• •	• •	• •	•					
3	Carya glabra									•							
3	Crataegus sp.			•				• •	• • •								
3	Ligustrum vulgare					2											
3	Fraxinus sp			! •							• •						
3	Standing dead											•					
3	~~Multiflora~~ Rosa multiflora rose					18											
3	Lonicera ~~maackii~~ morrowii			•		9											

DBH Measurement Rules



Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.

Record using the tally system from 1 to 10

1 2 3 4 5

ASH CANOPY CONDITION

1. **Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
2. **Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
3. **Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
4. **>50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
5. **Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.

A B C D E

ASH CANOPY BREAKUP CONDITION (for dead trees):
(if an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)

A: All main branches contain fine twigs (newly dead).
B: Over 50% of main branches have fine twigs.
C: Less than 50% of main branches have fine twigs.
D: Stem still standing and tertiary main branches present.
E: Central stem still standing.

CLEVELAND METROPARKS Plant Community Assessment Program Natural Woody Stem Data Sheet

Project Label: PCAP Project Name: 01BR2012 Plot No.: 1201 Page: 3 of 3

Explain subsample (additional room on back):

mod #	species	c	voucher#	# stems 0-1.4m browsed	% sub or super sample	# shrub clumps	size class (cm) woody stems >1.4m 1 0-<1	2 1-<2.5	3 2.5-<5	4 5-<10	5 10 - <15	6 15 - <20	7 20 - <25	8 25 - <30	9 30 - <35	10 35 - <40	11 >40 (record each tree)
3	Lindera benzoin			••		5											
3	~~Frangula alnus~~ Rhamnus ~~Rhamnus~~ Frangula			•						X							
4	Ulmus americana			•				•		•							
4	Crateagus sp.						•••	•• •	••			•					
4	Parthenocissus quinquefolia						••										
4	Vitis sp.						•										
4	Standing dead						•	••	•				••				
4	Carya ovata									•							
4	Carpinus caroliniana						•	••		••							
4	Carya glabra							•	•	•							
4	Prunus serasus									•							
4	Quercus alba														•		
4	Tilia americana											•		•			
4	Fraxinus sp.			dx!											•		
4	Acer saccharum								•								
4	Lonicera morrowii			••		7											
4	Lonicera maackii			•		3											
4	Lindera benzoin			•		3											
4	Ligustrum vulgare					10											
4	Rosa multiflora					11											
4	Juglans nigra			•													

DBH Measurement Rules



Woody Stem Deer Browse

Record the number of stems/plants between 0.5-1.0 meters tall that exhibit evidence of this years deer browse.

Record using the tally system from 1 to 10

1 2 3 4 5

ASH CANOPY CONDITION

1. **Healthy, full canopy:** A healthy ash canopy is normally thinner than many other trees such as maple.
2. **Thinning canopy:** There aren't as many leaves as there ought to be, but all top branches exposed to sunlight have leaves.
3. **Dieback:** Canopy is thinning and some top branches exposed to sunlight are dead (have no leaves). Lower branches, not exposed to sunlight, die naturally and are not considered.
4. **>50% Dieback:** The canopy has less than half of the leaves that should be there and/or half of the top branches are dead.
5. **Dead canopy:** No leaves remain in the canopy portion of the tree. It still counts as a 5 even if there are epicormic sprouts below the canopy (lowest branch) on the trunk.

A B C D E

ASH CANOPY BREAKUP CONDITION (for dead trees):
(if an ash receives a score of 5 (dead) under canopy condition it must also receive a breakup condition rank as described below)

A: All main branches contain fine twigs (newly dead).
B: Over 50% of main branches have fine twigs.
C: Less than 50% of main branches have fine twigs.
D: Stem still standing and tertiary main branches present.
E: Central stem still standing.

CLEVELAND METROPARKS Emerald Ash Borer - Fraxinus Sheet — *INTENSIVE MODULES ONLY* — *TREES ≥ 10CM ONLY*

Project Label: PCAP Project Name: 01B2012 Plot No.: 1201 Date: 6/20/2012 Page: 1 of 2

Module	Tree ID.	Species	Dead	c	Voucher #	DBH (cm)	Ht @ DBH	ASH Only: Ash condition	ASH Only: *Dead condition	ASH Only: # Exit holes	ASH Only: Epicormic present	ASH Only: Woodpecker holes
3	1	Fraxinus sp.				15.4		2		0	1	0
3	2	Fraxinus sp.				11.8		4		0	1	1
4	3	Fraxinus sp.				31.1		2		0	0	1
	4											
	5											
	6											
	7											
	8											
	9											
	10											
	11											
	12											
	13											
	14											
	15											
	16											
	17											
	18											
	19											
	20											
	21											
	22											
	23											
	24											
	25											



*** Change Intensive module numbers when necessary



Map all ash trees ≥10cm in each module using Tree ID number



* If Ash Condition scores 5 (dead) provide breakup score (A-E)
Count EAB exit holes 1.25m2 x ≥1.5m
Woodpecker and epicormic marked present (1) or absent (0)

CLEVELAND METROPARKS Plant Community Assessment Program: Invasive Species Survey Cleveland Metroparks

Tier 1: Early detection/ Rapid response		Presence				GPS
		NE	SE	SW	NW	
Microstegium vimineum	Japanese stiltgrass					
Ranunculus ficaria	Lesser Celandine					
Cynanchum louiseae (vine)	Black Swallow-wort					
Butomus umbellatus (wetland)	Flowering Rush					
Heracleum mantegazzianum	Giant Hogweed					
Tier 2: Assess as Needed		**# of Plants**				**comments**
		NE	SE	SW	NW	
Acer platanoides	Norway Maple					
Ailanthus altissima	Tree of Heaven					
Lonicera japonica (vine)	Japanese Honeysuckle					
Lythrum salicaria (wetland)	Purple Loosestrife					
Aegopodium podagraria (G-cover)	Bishop's Goutweed					
Celastrus orbiculatus (vine)	Asian Bittersweet					
Torilis sp.	Hedgeparsley					
Conium maculatum	Poison Hemlock					
Rhamnus cathartica	Common Buckthorn (shrub)		1	x		
Berberis thunbergii	Japanese Barberry (shrub)	1	1	1	2	
Alnus glutinosa	European Alder					
Dipsacus laciniatus	Cut-leaf Teasel					
Elaeagnus umbellata	Autumn Olive (shrub)					
Lonicera maackii	Amur Honeysuckle (shrub)	2	1	1	2	
Euonymus fortunei	Wintercreeper					
Tier 3: Presence is of Interest		**# of Plants**				**comments**
		NE	SE	SW	NW	
Convallaria majalis (G-cover)	Lily of the Valley					
Coronilla varia (G-cover)	Crown Vetch					
Eleutherococcus pentaphyllus	Five-leaf Aralia (shrub)					
Pachysandra terminalis (G-cover)	Japanese Pachysandra					
Philadelphus coronarius	Mock Orange (shrub)					
Pulmonaria officinalis (G-cover)	Lungwort					
Rubus phoenicolasius	Wineberry					
Iris pseudacorus (wetland)	Yellow Flag Iris					
Ornithogalum umbellatum	Star of Bethlehem					
Viburnum opulus var. opulus	European Cranberry (shrub)					
Viburnum plicatum	Doublefile Viburnum (shrub)					
Tier 4: Widespread and abundant		**Presence**				**comments**
		NE	SE	SW	NW	
Alliaria petiolata	Garlic Mustard	3	2	2		
Ligustrum vulgare	Common Privet (shrub)		1	2°	3°	
L. morrowii, L. tatarica	Bush Honeysuckles (shrub)	2	1	1°	2°	
Phalaris arundinacea	Reed Canarygrass					
Phragmites australis (wetland)	Phragmites					
Polygonum cuspidatum	Japanese Knotweed					
Frangula alnus	Glossy Buckthorn (shrub)			1	3°	
Rosa multiflora	Multiflora Rose (shrub)	3	3	2	3°	
Typha angustifolia, T. x.glauca	Cattails (wetland)					
Cirsium arvense	Canada thistle					
Dipsacus fullonum	Common Teasel					
Hesperis matronalis	Dame's Rocket					
Vinca minor (G-cover)	Periwinkle					

Presence
X: yes

of Plants
1: 1-10
2: 11-50.
3: 51-100
4: 101-1,000
5: >1,000

of Plants
1: 1-10
2: 11-50.
3: 51-100
4: 101-1,000
5: >1,000

Presence
X: yes

Note: For Ground-cover plants record "stem #" but in comment field describe # of colonies and patch size (S,M, L)

CLEVELAND METROPARKS Plant Community Assessment Program - Soils, Crown Cover, Standing Biomass Data Sheet 6a

Project label: PCAP Project Name: 01BR2012 Plot No.: 1201

Cleveland Metroparks

SOIL PIT DESCRIPTION: Excavate 20 cm plug wih shovel. Describe using Munsell chart, visual exam, texture, and odor

Soil pit module # 3 (one per entire plot)

5 cm	matrix color	10YR 3/2		
	mottle color	n/a		
	%mottle	0		
	oxid roots	(Y)	N	
	texture*	2		
	redox features**	(Y)	N	
	hydr cond ***	I S (M)	D	
20 cm	matrix color	10YR 4/3		
	mottle color	n/a		
	%mottle	0		
	oxid roots	(Y)	N	
	texture*	2		
	redox features**	(Y)	N	
	hydro cond ***	I S M (D)		

* refer to texture classes on reverse side

** e.g. hydrogen sulfide odor, gleying, etc.

*** Circle one:
I=indundated S=saturated M=moist D=dry

Notes: include evidence of earthworms (worms, castings, middens)

earthworms within soil pit, castings in proximity to pit soil pit

SOIL SAMPLES Standard procedure: collect a soil sample of the top 10 cm of soil from center of each intensive module and composite the sample

Soil Collection Module	Horizon (A, B, C)
2,3,8,9 composited	A

Soil Series/Type: Mahoning silt loam

Soil Series Source: Ohio Soil Survey

Landform type: Till plains

Depth to rest. Layer: > 80 inches

Parent Material: Till

DRAINAGE

- □ Excessively dr.
- □ Somewhat excessively
- □ Well drained
- □ Moderately well dr.
- X Somewhat poorly dr.
- □ Very poorly dr.
- □ Impermeable surface

A. yarw 6/28/2012

SOIL DEPTH MEASUREMENT: Measure to the nearest 0.1 cm in center of intensive modules. If >30.5 cm, record as >30

mod#	1 litter+ organic depth (cm)	2 litter depth (cm)	water depth (cm)	depth sat soil (cm)
1	0.3	0.3	0	>30
2	1.8	1.8	0	>30
3	0.2	0.2	0	>30
4	0.1	0.1	0	>30

EARTH SURFACE & GROUND COVER

Underlying Earth Surface*	percent	Ground Cover	percent
(Sum = 100%)		(Each ≤ 100%)	
Histosol	Ø	Coarse Woody Debris***	15
Mineral Soil	97	Fine Woody Debris****	5
Gravel-Cobble*	Ø	Litter	55
Boulder**	3	Duff (Ferm.+ Humus)	Ø
Bedrock	Ø	Bryophyte- Lichen	1
* Gravel-Cobble = 1/16-10"		Water	Ø
**Boulder = > 10 in		Bare Soil	5
*** >5 cm in diameter		Road/Trail	Ø
**** <5 cm in diameter		Other	

TRAIL INFORMATION:

record type and cover for each

Type	%Cover
□ All Purpose	
□ Bridle	
□ Hiking sanctioned	
□ Bootleg unsanctioned	
□ Gravel	
□ Deer	

No trails

COVER BY STRATA %

estimate using midpoints of 5,ex:3, 8, 13

Strata	Height Range (m)	Total Cover (%)
Tree	>5	88
Shrub	.5 - 5	63
Herb	0 - 5	48
(Floating)*	-	
(Aquatic)*	-	

* rooted and floating or slightly emersed

** submersed, most plant mass below surface

SEE BACK OF PAGE FOR "TYPICAL"STRATA DESCRIPTIONS. STRATA CAN VARY BY COVER TYPE.

STAND SIZE

- □ >600 x plot size
- □ > 100 x plot size
- □ 10-100 x plot size
- □ 3-10 x plot size
- ☑ 1-3 x plot size
- □ < plot size

PERCENT MOTTLES (USE CLASS CODES):

Class	Code Conv.	Code NASIS	Criteria: % of Surface Area Covered
Few	f	#	< 2
Common	c	#	2 to < 20
Many	m	#	≥ 20



SOIL TEXTURE: Record the code for the soil texture of the 5 cm and 20 cm layers. To estimate texture, collect a soil sample from the appropriate layer and moisten it with water to the consistency of modeling clay/wet newspaper; the sample should be wet enough that all of the particles are saturated but excess water does not freely flow from the sample when squeezed. Attempt to roll the sample into a ball. If the soil will not stay in a ball and has a grainy texture, the texture is either sandy or coarse sandy. If the soil does form a ball, squeeze the sample between your fingers and attempt to form a self-supporting ribbon. Samples which form both a ball and a ribbon should be coded as clayey; samples which form a ball but not a ribbon should be coded as loamy.

0= Organic
1= Loamy
2= Clayey
3= Sandy
4= Coarse Sand
9= Not measured - make plot note

Geomorphic Component - Three-dimensional descriptors of parts of landforms or microfeatures that are best applied to areas. Unique descriptors are available for Hills, Terraces, Mountains, and Flat Plains; e.g., (for Hills) *nose slope* or *NS*.

Hills	Code PDP	Code NASIS
interfluve	IF	IF
head slope	HS	HS
nose slope	NS	NS
side slope	SS	SS
base slope	---	BS



(FJS, 1998; adapted from Ruhe, 1975)

Terraces	Code
riser	RI
tread	TR



(S&W, 1998)

Hillslope - Profile Position (Hillslope Position in PDP) - Two-dimensional descriptors of parts of line segments (i.e., slope position) along a transect that runs up and down the slope; e.g., *backslope* or *BS*. This is best applied to transects or points, not areas.

Position	Code
summit	SU
shoulder	SH
backslope	BS
footslope	FS
toeslope	TS



(FJS, 1998; adapted from Ruhe, 1975)

HYDROLOGIC REGIME Modified from Grossman et al 1998. (Frequency and duration of flooding.)

UPLAND: Not a wetland. Very rarely flooded.

INTERMITTENTLY/SEASONALLY SATURATED: Dry at least once per year. Surface water is seldom present, but substrate is saturated to surface for extended periods during the growing season.

PERMANENTLY/SEMIPERMANENTLY SATURATED: Dry less than once per year. Surface water is seldom present, but substrate is saturated to surface for extended periods during the growing season. Equivalent to Cowardin's Saturated modifier.

OCCASIONALLY FLOODED: Surface water can be present for brief periods during growing season, but not in most years. Often characterizes flood-plain upper terraces.

TEMPORARILY FLOODED: Surface water present for brief periods during growing season, but water table usually lies well below soil surface. Often characterizes flood-plain levees and lower terraces. Equivalent to Cowardin's Temporary modifier.

INTERMITTENTLY FLOODED : Substrate is usually exposed, but surface water can be present for variable periods without detectable seasonal periodicity. Inundation is not predictable to a given season and is dependent upon highly localized rain storms. This modifier was developed for use in the arid West for water regimes of Playa lakes, intermittent streams, and dry washes but can be used in other parts of the U.S. where appropriate. This modifier can be applied to both wetland and non-wetland situations. Equivalent to Cowardin's Intermittently Flooded modifier.

SEMIPERMANENTLY FLOODED (exposed <1/year): Surface water persists throughout the growing season in most years. Land surface is normally saturated when water level drops below soil surface. Includes Cowardin's Intermittently Exposed and Semipermanently Flooded modifiers.

PERMANENTLY FLOODED: Water covers the land surface at all times of the year in all years. Equivalent to Cowardin's "permanently flooded".

UNKNOWN: The hydrologic regime cannot be determined from the available information.

CLEVELAND METROPARKS Plant Community Assessment Program - Plant Cover and Earth Surface

Project Label: PCAP Project Name: 01BR2002 Plot No.: 1201

Cleveland Metroparks

STANDING BIOMASS (required for emergent wetlands): collected in 0.1m clip plots (32x32 cm) from corners 1 and 3 in each intensive module. Required for VIBI-E score calculation. C?=check when collected

Module #	C?	Corner	Corner

CLASSIFICATION

(FIT = excellent, g Fit and Confidence

Hydrogeomorphic class (WETLANDS ONLY):

- ◻ DEPRESSION Fit=___ Conf=___
- ◻ IMPOUNDMENT ◻ Beaver ◻ Human Fit=___ Conf=___
- ◻ RIVERINE ◻ Headwater ◻ Mainstem ◻ Channel Fit=___ Conf=___
- ◻ SLOPE (ground water hydrology or on a physical slop) Fit=___ Conf=___
- ◻ FRINGING ◻ Reservoir ◻ Natural Lake Fit=___ Conf=___
- ◻ COASTAL (specify subclass) Fit=___ Conf=___
- ◻ BOG (strongly, moderately, weekly ombrotrophic) Fit=___ Conf=___

Ohio EPA VIBI Plant Community Class (WETLANDS ONLY):

- ◻ FOREST ◻ swamp forest ◻ bog forest ◻ forest seep Fit=___ Conf=___
- ◻ EMERGENT ◻ marsh ◻ wet meadow ◻ open bog Fit=___ Conf=___
- ◻ SHRUB ◻ shrub swamp ◻ tall sh. bog ◻ tall sh. fen Fit=___ Conf=___

McNAB INDICES (degrees) + for up - for down

(FILLED OUT USING GIS PROGRAM - DO NOT FILL OUT IN FIELD)

		LFI*	TSI**
At aspect	N		
+45 degrees	NE		
+90 degrees	E		
+135 degrees	SE		
+180 degrees	S		
+225 degrees	SW		
+270 degrees	W		
+315 degrees	NW		

LFI is angle of plot to the horizon. TSI is angles formed by local slopes. For TSI measure angle from recorders eye to eye of person standing ~10 m away.

* Landform Index (position within landscape)

** Terrain Shape Index (site microtopographic shape)

MICROTOPOGRAPHIC FEATURE COUNTS - Intensive modules only

Ranks for microhabitat features. Select one or select two and average the score.NOTE: If mod falls on a slope automatically gets ranked based on steepness (1-3) to begin + any features present

Slope 1 = slight elevational grade across module (hill) Slope 2 = falls on slope ~20° Slope 3 = maximum steepness that can be safely sampled ~45°

0 feature is absent or functionally absent from the wetland

3 feature is present in the wetland in very small amounts or if more common, of low quality

7 feature is present in moderate amounts, but not of highest quality, or in small amounts of highest quality

10 feature is present in moderate or greater amounts and of highest quality

c.w.d. - count for pieces with minimum 1m length

mod#	corner	no. of tussocks depth 3 1x1m (count)	no. of hummocks uplands (Tip-Ups) depth 2 3.16x3.16m (count)	no. macro. depressions depth 1 10x10m (count)	c.w.d (2-12 cm) depth 1 10x10m (count)	c.w.d (12-40cm) depth 1 10x10m (count)	c.w.d >40 cm depth 1 10x10m (count)	microhab. interspers. depth 1 10x10m (rank)	microhab. SLOPE 10x10m (rank)
1		0	0	1	15	0	0	1	0
2		0	0	0	17	+2	0	2	1
3		0	0	1	18	23	0	1	1
4		0	0	0	16	3	0	1	1

NOTE: tussock and hummocks are counted in BOTH nested quadrat corners but counts are aggregated.

17

14 5

CROWN COVER (DENSIOMETER) Make 4 readings per module facing N, S, E, W. Place dot count in corresponding space. (4 dots per grid square)

Module	N	S	E	W
1	7	5	6	3
2	5	7	5	3
3	6	5	5	5
4	4	15	15	2
	~~6~~ 8	~~10~~ 4	7 6	3 4
	5	9	4	3
	5	6	7	3
	5	5	5	6
	6	6	5	5
	1	16	13	3
	6	15	19	1

COVER BY STRATA

STRATUM	GENERAL FORM
Tree (generally >5 m)	Tree (overstory), very tall shrubs*, liana, epiphyte)
Shrub (generally 0.5 to 5 m)	Tree (sapling), shrub, liana, epiphyte)
Herb (Field)	Herb, dwarf-shrub**, tree (seedling***)
Floating	Floating
Aquatic (submerged)	Submerged

*Very tall shrubs are sometimes included in the tree stratum

**Can also include seedlings of shrubs, i.e. all shrubs <0.5m

***Tree seedlings are often defined as up to 1.4 m height or as <2.5 cm DBH in which case they would span the herb and shrub layers.





FIGURE 3-20.—Generalized section of Upper Devonian, Mississippian, and Lower Pennsylvanian formations in northeastern Ohio. Asterisks indicate units that are fossiliferous. This composite section represents about 400 meters of rock exposed across the area. The section is not to scale, but the thicknesses indicated are proportional. The term "Waverly" is used in the older literature to refer to Mississippian rocks in Ohio. Some geologists use the European term "Carboniferous," which encompasses the Mississippian and Pennsylvanian Periods of the U.S. Many units have been named within the Cuyahoga Formation, but most units are local and cannot be traced over great distances. The Black Hand Member is a spectacular massive sandstone that is fairly widespread but discontinuous. See Hyde (1953), Hoover (1960), and Collins (1979) for more information on Mississippian rocks in Ohio. See figure 3-18 for explanation of rock types.

FORM B-1: BUFFER SAMPLE PLOTS (Front)

Reviewed by (initial): _____

Site ID: PCAPBR1201

DATE: 06/20/2012

Location: ● AA Center ○ N ○ S ○ E ○ W

Fill in bubble(s) if plot(s) could not be sampled and flag → ○ Plot 1 ○ Plot 2 ○ Plot 3

Buffer Natural Cover Strata

Fill in bubbles for all that apply: Canopy Type: D = Deciduous; E = Evergreen. Leaf Type: B = Broadleaf; N = Needle Leaf. Absent: No tree canopy.
Strata Section: Fill in appropriate cover class bubble for each strata type for each plot. 0 = Absent; 1 = Sparse(<10%); 2=Moderate(10-40%); 3 = Heavy (40-75%); 4 = Very Heavy (>75%)

Buffer Plot 1	Canopy Type: ● E	Absent: ○	Flag	Buffer Plot 2	Canopy Type: D E	Absent: ○	Flag	Buffer Plot 3	Canopy Type: D E	Absent: ○	Flag
	Leaf Type: ● N				Leaf Type: B N				Leaf Type: B N		
Big Trees (>0.3m DBH)	3			Big Trees (>0.3m DBH)				Big Trees (>0.3m DBH)			
Small Trees (<0.3m DBH)	3			Small Trees (<0.3m DBH)				Small Trees (<0.3m DBH)			
Woody Shrubs, Saplings (0.5m-5m HIGH)	3			Woody Shrubs, Saplings (0.5m-5m HIGH)				Woody Shrubs, Saplings (0.5m-5m HIGH)			
Woody Shrubs, Saplings (<0.5m HIGH)	3			Woody Shrubs, Saplings (<0.5m HIGH)				Woody Shrubs, Saplings (<0.5m HIGH)			
Herbs, Forbs and Grasses	3			Herbs, Forbs and Grasses				Herbs, Forbs and Grasses			
Bare ground	2			Bare ground				Bare ground			
Litter, duff	3			Litter, duff				Litter, duff			
Rock	1			Rock				Rock			
Water	0			Water				Water			
Submerged Vegetation	0			Submerged Vegetation				Submerged Vegetation			

Stressor Presence/Absence - Confirm that a filled data bubble indicates presence and an unfilled bubble indicates absence by filling this bubble. ●

Residential and Urban Stressors					Hydrology Stressors					Agricultural & Rural Stressors				
Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag
Road - gravel	○	○	○		Ditches, Channelization	○	○	○		Pasture/Hay	○	○	○	
Road - two lane	○	○	○		Dike/Dam/Road/RR Bed (IMPEDE FLOW)	○	○	○		Range	○	○	○	
Road - four lane	○	○	○		Water Level Control Structure	○	○	○		Row Crops	○	○	○	
Parking Lot/Pavement	○	○	○		Excavation, Dredging	○	○	○		Fallow Field (RECENT-RESTING ROW CROP FIELD)	○	○	○	
Golf Course	○	○	○		Fill/Spoil Banks	○	○	○		Fallow Field (OLD - GRASS, SHRUBS, TREES)	○	○	○	
Lawn/Park	○	○	○		Freshly Deposited Sediment (UNVEGETATED)	○	○	○		Nursery	○	○	○	
Suburban Residential	○	○	○		Soil Loss/Root Exposure	○	○	○		Dairy	○	○	○	
Urban/Multifamily	○	○	○		Wall/Riprap	○	○	○		Orchard	○	○	○	
Landfill	○	○	○		Inlets, Outlets	○	○	○		Confined Animal Feeding	○	○	○	
Dumping	○	○	○		Point Source/Pipe (EFFLUENT OR STORMWATER)	○	○	○		Rural Residential	○	○	○	
Trash	○	○	○		Impervious surface input (SHEETFLOW)	●	○	○		Gravel Pit	○	○	○	
Other: _____	○	○	○		Other: _____	○	○	○		Irrigation	○	○	○	
Other: _____	○	○	○		Other: _____	○	○	○		Other: _____	○	○	○	

Industrial Development Stressors					Habitat/Vegetation Stressors									
Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag
Oil Drilling	○	○	○		Forest Clear Cut	○	○	○		Herbicide Use	○	○	○	
Gas Wells	○	○	○		Forest Selective Cut	●	○	○		Mowing/Shrub Cutting	○	○	○	
Mine (surface)	○	○	○		Tree Plantation	○	○	○		Trails	○	○	○	
Mine (underground)	○	○	○		Tree Canopy Herbivory (INSECT)	○	○	○		Soil Compaction (ANIMAL OR HUMAN)	○	○	○	
Military	○	○	○		Shrub Layer Browsed (WILD OR DOMESTIC)	●	○	○		Offroad vehicle damage	○	○	○	
Other: _____	○	○	○		Highly Grazed Grasses (OVERALL <3" HIGH)	○	○	○		Soil erosion (FROM WIND, WATER, OR OVERUSE)	○	○	○	
Other: _____	○	○	○		Recently Burned Forest Canopy	○	○	○		Other: _____	○	○	○	
Other: _____	○	○	○		Recently Burned Grassland (BLACKENED)	○	○	○		Other: _____	○	○	○	

Flag codes: K = No measurement made, U = Suspect measurement., F1,F2, etc. = misc. flags assigned by each field crew.
Explain all flags in comment section on the back of this form

2428168304

FORM B-1: BUFFER SAMPLE PLOTS - TARGETED ALIEN SPECIES (Back)

Reviewed by (initial): _____

Site ID: PCAPBR120 DATE: 06/20/2012

● Confirm a filled data bubble indicates presence and an unfilled bubble indicates absence by filling in this bubble

Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag
Eurasian Watermilfoil	○	○	○		Purple Loosestrife	○	○	○		Johnson Grass	○	○	○	
Water hyacinth	○	○	○		Knotweed	○	○	○		Kudzu	○	○	○	
Yellow Floating Heart	○	○	○		Japanese Knotweed	○	○	○		Multiflora Rose	●	○	○	
Giant Salvinia	○	○	○		Perennial Pepperweed	○	○	○		Common Buckthorn	○	○	○	
Garlic Mustard	○	○	○		Giant Reed	○	○	○		Himalayan Blackberry	○	○	○	
Poison Hemlock	○	○	○		Cheatgrass	○	○	○		Tamarisk	○	○	○	
Mile-A-Minute Weed	○	○	○		Reed Canary Grass	○	○	○		Other:	○	○	○	
Birdsfoot Trefoil	○	○	○		Common Reed	○	○	○		Other:	○	○	○	
Canada Thistle	○	○	○		Leafy Spurge	○	○	○		Other:	○	○	○	
										Other:	○	○	○	

PLOT COORDINATES

Provide GPS coordinates at the center of the Buffer Plot (#3) at the far end of each Buffer Transect and for the Buffer Plot at the AA CENTER. Indicate the location of the plot coordinates by filling in the appropriate bubble.

If Buffer Plot 3 can not be accessed, take the coordinates at the nearest practicable location ALONG THE TRANSECT. This is important because all Buffer Plots are centered on the Buffer Transects and the coordinates will indicate the location of the transect. Fill in the "nearest practicable location" bubble, fill in the flag box, and describe where the coordinates were taken and why in the comment section below. The coordinates of the nearest practicable location can be either placed as close to the center of Plot 3 as possible or at the center of the last accessible Buffer Plot.

Location of coordinates (choose one): Flag

● AA CENTER ○ N3 ○ S3 ○ E3 ○ W3 ○ Nearest practicable location (flag and comment below)

Latitude North 41.31459 Longitude West 081.61902

Use Decimal Degrees; NAD83

Flag	Comments
1	2 lane asphalt road proximal (< 10 m) from AA center

FORM B-1: BUFFER SAMPLE PLOTS (Front)

Reviewed by (initial): _____

Site ID: PCAPBR1201 DATE: 06 / 20 / 2012

Location: ○ AA Center ◉ N ○ S ○ E ○ W

Fill in bubble(s) if plot(s) could not be sampled and flag → ○ Plot 1 ○ Plot 2 ○ Plot 3

Buffer Natural Cover Strata

Fill in bubbles for all that apply: Canopy Type: D = Deciduous; E = Evergreen. Leaf Type: B = Broadleaf; N = Needle Leaf. Absent No tree canopy.
Strata Section: Fill in appropriate cover class bubble for each strata type for each plot. 0 = Absent; 1 = Sparse(<10%); 2=Moderate(10-40%); 3 = Heavy (40-75%); 4 = Very Heavy (>75%)

Buffer Plot 1 (Canopy Type: D E; Leaf Type: B N; Absent: ◉; Flag)		Buffer Plot 2 (Canopy Type: D E; Leaf Type: B N; Absent: ◉; Flag)		Buffer Plot 3 (Canopy Type: D E; Leaf Type: B N; Absent: ◉; Flag)	
Big Trees (>0.3m DBH)	◉	Big Trees (>0.3m DBH)	◉	Big Trees (>0.3m DBH)	◉
Small Trees (<0.3m DBH)	◉	Small Trees (<0.3m DBH)	◉	Small Trees (<0.3m DBH)	◉
Woody Shrubs, Saplings (0.5m-5m HIGH)	◉	Woody Shrubs, Saplings (0.5m-5m HIGH)	◉	Woody Shrubs, Saplings (0.5m-5m HIGH)	◉
Woody Shrubs, Saplings (<0.5m HIGH)	◉	Woody Shrubs, Saplings (<0.5m HIGH)	◉	Woody Shrubs, Saplings (<0.5m HIGH)	◉
Herbs, Forbs and Grasses	4	Herbs, Forbs and Grasses	4	Herbs, Forbs and Grasses	4
Bare ground	1	Bare ground	◉	Bare ground	◉
Litter, duff	◉	Litter, duff	◉	Litter, duff	◉
Rock	◉	Rock	◉	Rock	◉
Water	◉	Water	◉	Water	◉
Submerged Vegetation	◉	Submerged Vegetation	◉	Submerged Vegetation	◉

Stressor Presence/Absence - Confirm that a filled data bubble indicates presence and an unfilled bubble indicates absence by filling this bubble. ◉

Residential and Urban Stressors	1	2	3	Flag
Road - gravel	○	○	○	
Road - two lane	○	○	○	
Road - four lane	○	○	○	
Parking Lot/Pavement	◉	○	○	
Golf Course	○	○	○	
Lawn/Park	◉	◉	◉	
Suburban Residential	○	○	○	
Urban/Multifamily	○	○	○	
Landfill	○	○	○	
Dumping	○	○	○	
Trash	○	○	◉	
Other: _____	○	○	○	
Other: _____	○	○	○	

Hydrology Stressors	1	2	3	Flag
Ditches, Channelization	○	○	○	
Dike/Dam/Road/RR Bed (IMPEDE FLOW)	○	○	○	
Water Level Control Structure	○	○	○	
Excavation, Dredging	○	○	○	
Fill/Spoil Banks	○	○	○	
Freshly Deposited Sediment (UNVEGETATED)	○	○	○	
Soil Loss/Root Exposure	○	○	○	
Wall/Riprap	○	○	○	
Inlets, Outlets	○	○	○	
Point Source/Pipe (EFFLUENT OR STORMWATER)	○	○	○	
Impervious surface input (SHEETFLOW)	◉	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Agricultural & Rural Stressors	1	2	3	Flag
Pasture/Hay	○	○	○	
Range	○	○	○	
Row Crops	○	○	○	
Fallow Field (RECENT-RESTING ROW CROP FIELD)	○	○	○	
Fallow Field (OLD - GRASS, SHRUBS, TREES)	○	○	○	
Nursery	○	○	○	
Dairy	○	○	○	
Orchard	○	○	○	
Confined Animal Feeding	○	○	○	
Rural Residential	○	○	○	
Gravel Pit	○	○	○	
Irrigation	○	○	○	
Other: _____	○	○	○	

Industrial Development Stressors	1	2	3	Flag
Oil Drilling	○	○	○	
Gas Wells	○	○	○	
Mine (surface)	○	○	○	
Mine (underground)	○	○	○	
Military	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Habitat/Vegetation Stressors	1	2	3	Flag
Forest Clear Cut	○	○	○	
Forest Selective Cut	○	○	○	
Tree Plantation	○	○	○	
Tree Canopy Herbivory (INSECT)	○	○	○	
Shrub Layer Browsed (WILD OR DOMESTIC)	○	○	○	
Highly Grazed Grasses (OVERALL <3" HIGH)	○	○	○	
Recently Burned Forest Canopy	○	○	○	
Recently Burned Grassland (BLACKENED)	○	○	○	

Herbicide Use	1	2	3	Flag
Herbicide Use	○	○	○	
Mowing/Shrub Cutting	◉	◉	◉	
Trails	○	○	○	
Soil Compaction (ANIMAL OR HUMAN)	○	○	○	
Offroad vehicle damage	○	○	○	
Soil erosion (FROM WIND, WATER, OR OVERUSE)	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Flag codes: K = No measurement made, U = Suspect measurement., F1,F2, etc. = misc. flags assigned by each field crew.
Explain all flags in comment section on the back of this form

2428168304

FORM B-1: BUFFER SAMPLE PLOTS - TARGETED ALIEN SPECIES (Back)

Reviewed by (initial): _____

Site ID: PCAPBR1201 DATE: 06/20/2012

● Confirm a filled data bubble indicates presence and an unfilled bubble indicates absence by filling in this bubble

Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag
Eurasian Watermilfoil	○	○	○		Purple Loosestrife	○	○	○		Johnson Grass	○	○	○	
Water hyacinth	○	○	○		Knotweed	○	○	○		Kudzu	○	○	○	
Yellow Floating Heart	○	○	○		Japanese Knotweed	○	○	○		Multiflora Rose	○	○	●	
Giant Salvinia	○	○	○		Perennial Pepperweed	○	○	○		Common Buckthorn	○	○	○	
Garlic Mustard	○	○	○		Giant Reed	○	○	○		Himalayan Blackberry	○	○	○	
Poison Hemlock	○	○	○		Cheatgrass	○	○	○		Tamarisk	○	○	○	
Mile-A-Minute Weed	○	○	○		Reed Canary Grass	○	○	○		Other:	○	○	○	
Birdsfoot Trefoil	○	○	○		Common Reed	○	○	○		Other:	○	○	○	
Canada Thistle	○	○	○		Leafy Spurge	○	○	○		Other:	○	○	○	
										Other:	○	○	○	

PLOT COORDINATES

Provide GPS coordinates at the center of the Buffer Plot (#3) at the far end of each Buffer Transect and for the Buffer Plot at the AA CENTER. Indicate the location of the plot coordinates by filling in the appropriate bubble.

If Buffer Plot 3 can not be accessed, take the coordinates at the nearest practicable location ALONG THE TRANSECT. This is important because all Buffer Plots are centered on the Buffer Transects and the coordinates will indicate the location of the transect. Fill in the "nearest practicable location" bubble, fill in the flag box, and describe where the coordinates were taken and why in the comment section below. The coordinates of the nearest practicable location can be either placed as close to the center of Plot 3 as possible or at the center of the last accessible Buffer Plot.

Location of coordinates (choose one):

○ AA CENTER ● N3 ○ S3 ○ E3 ○ W3 ○ Nearest practicable location (flag and comment below)

Flag: ☐

Latitude North 41.31587 Longitude West 081.61906

Use Decimal Degrees; NAD83

Flag	Comments

FORM B-1: BUFFER SAMPLE PLOTS (Front)

Reviewed by (initial): _____

Site ID: PCAPBR1201 DATE: 06/20/2012

Location: ○ AA Center ○ N ● S ○ E ○ W

Fill in bubble(s) if plot(s) could not be sampled and flag → ○ Plot 1 ○ Plot 2 ○ Plot 3

Buffer Natural Cover Strata

Fill in bubbles for all that apply: Canopy Type: D = Deciduous; E = Evergreen. Leaf Type: B = Broadleaf; N = Needle Leaf. Absent: No tree canopy.
Strata Section: Fill in appropriate cover class bubble for each strata type for each plot. 0 = Absent; 1 = Sparse(<10%); 2=Moderate(10-40%); 3 = Heavy (40-75%); 4 = Very Heavy (>75%)

Buffer Plot 1	Canopy Type: D E / Leaf Type: B N	Absent: ○	Flag	Buffer Plot 2	Canopy Type: D E / Leaf Type: B N	Absent: ○	Flag	Buffer Plot 3	Canopy Type: D E / Leaf Type: B N	Absent: ○	Flag
Big Trees (>0.3m DBH)	0 1 2 **3** 4			Big Trees (>0.3m DBH)	0 1 2 **3** 4			Big Trees (>0.3m DBH)	0 1 2 **3** 4		
Small Trees (<0.3m DBH)	0 1 2 **3** 4			Small Trees (<0.3m DBH)	0 1 2 **3** 4			Small Trees (<0.3m DBH)	0 1 2 **3** 4		
Woody Shrubs, Saplings (0.5m-5m HIGH)	0 1 **2** 3 4			Woody Shrubs, Saplings (0.5m-5m HIGH)	0 1 **2** 3 4			Woody Shrubs, Saplings (0.5m-5m HIGH)	0 1 **2** 3 4		
Woody Shrubs, Saplings (<0.5m HIGH)	0 **1** 2 3 4			Woody Shrubs, Saplings (<0.5m HIGH)	0 **1** 2 3 4			Woody Shrubs, Saplings (<0.5m HIGH)	0 **1** 2 3 4		
Herbs, Forbs and Grasses	0 1 **2** 3 4			Herbs, Forbs and Grasses	0 1 **2** 3 4			Herbs, Forbs and Grasses	0 1 **2** 3 4		
Bare ground	0 **1** 2 3 4			Bare ground	0 **1** 2 3 4			Bare ground	0 **1** 2 3 4		
Litter, duff	0 1 2 **3** 4			Litter, duff	0 1 2 **3** 4			Litter, duff	0 1 2 3 **4**		
Rock	0 **1** 2 3 4			Rock	**0** 1 2 3 4			Rock	0 **1** 2 3 4		
Water	**0** 1 2 3 4			Water	**0** 1 2 3 4			Water	**0** 1 2 3 4		
Submerged Vegetation	**0** 1 2 3 4			Submerged Vegetation	**0** 1 2 3 4			Submerged Vegetation	**0** 1 2 3 4		

Stressor Presence/Absence - Confirm that a filled data bubble indicates presence and an unfilled bubble indicates absence by filling this bubble. ●

Residential and Urban Stressors

Fill bubble if present - Plot	1	2	3	Flag
Road - gravel	○	○	○	
Road - two lane	●	●	○	3
Road - four lane	○	○	○	
Parking Lot/Pavement	○	○	○	
Golf Course	○	○	○	
Lawn/Park	○	○	○	
Suburban Residential	○	○	○	
Urban/Multifamily	○	○	○	
Landfill	○	○	○	
Dumping	○	○	○	
Trash	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Hydrology Stressors

Fill bubble if present - Plot	1	2	3	Flag
Ditches, Channelization	○	○	○	
Dike/Dam/Road/RR Bed (IMPEDE FLOW)	○	○	○	
Water Level Control Structure	○	○	○	
Excavation, Dredging	○	○	○	
Fill/Spoil Banks	○	○	○	
Freshly Deposited Sediment (UNVEGETATED)	○	○	○	
Soil Loss/Root Exposure	●	○	○	1
Wall/Riprap	○	○	○	
Inlets, Outlets	○	○	○	
Point Source/Pipe (EFFLUENT OR STORMWATER)	○	●	○	34
Impervious surface input (SHEETFLOW)	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Agricultural & Rural Stressors

Fill bubble if present - Plot	1	2	3	Flag
Pasture/Hay	○	○	○	
Range	○	○	○	
Row Crops	○	○	○	
Fallow Field (RECENT-RESTING ROW CROP FIELD)	○	○	○	
Fallow Field (OLD - GRASS, SHRUBS, TREES)	○	○	○	
Nursery	○	○	○	
Dairy	○	○	○	
Orchard	○	○	○	
Confined Animal Feeding	○	○	○	
Rural Residential	○	○	○	
Gravel Pit	○	○	○	
Irrigation	○	○	○	
Other: _____	○	○	○	

Industrial Development Stressors

Fill bubble if present - Plot	1	2	3	Flag
Oil Drilling	○	○	○	
Gas Wells	○	○	○	
Mine (surface)	○	○	○	
Mine (underground)	○	○	○	
Military	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Habitat/Vegetation Stressors

Fill bubble if present - Plot	1	2	3	Flag
Forest Clear Cut	○	○	○	
Forest Selective Cut	○	○	○	
Tree Plantation	○	○	○	
Tree Canopy Herbivory (INSECT)	○	○	○	
Shrub Layer Browsed (WILD OR DOMESTIC)	●	●	○	
Highly Grazed Grasses (OVERALL <3" HIGH)	○	○	○	
Recently Burned Forest Canopy	○	○	○	
Recently Burned Grassland (BLACKENED)	○	○	○	

Fill bubble if present - Plot	1	2	3	Flag
Herbicide Use	○	○	○	
Mowing/Shrub Cutting	●	○	○	2
Trails	○	○	○	
Soil Compaction (ANIMAL OR HUMAN)	○	○	○	
Offroad vehicle damage	○	○	○	
Soil erosion (FROM WIND, WATER, OR OVERUSE)	○	●	●	4,5
Other: _____	○	○	○	
Other: _____	○	○	○	

Flag codes: K = No measurement made, U = Suspect measurement., F1,F2, etc. = misc. flags assigned by each field crew.
Explain all flags in comment section on the back of this form

2428168304

FORM B-1: BUFFER SAMPLE PLOTS - TARGETED ALIEN SPECIES (Back)

Reviewed by (initial): _____

Site ID: PCAPBR1201 DATE: 06/20/2012

● Confirm a filled data bubble indicates presence and an unfilled bubble indicates absence by filling in this bubble

Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag
Eurasian Watermilfoil	○	○	○		Purple Loosestrife	○	○	○		Johnson Grass	○	○	○	
Water hyacinth	○	○	○		Knotweed	○	○	○		Kudzu	○	○	○	
Yellow Floating Heart	○	○	○		Japanese Knotweed	○	○	○		Multiflora Rose	○	○	○	
Giant Salvinia	○	○	○		Perennial Pepperweed	○	○	○		Common Buckthorn	○	○	○	
Garlic Mustard	●	●	○		Giant Reed	○	○	○		Himalayan Blackberry	○	○	○	
Poison Hemlock	○	○	○		Cheatgrass	○	○	○		Tamarisk	○	○	○	
Mile-A-Minute Weed	○	○	○		Reed Canary Grass	○	○	○		Other:	○	○	○	
Birdsfoot Trefoil	○	○	○		Common Reed	○	○	○		Other:	○	○	○	
Canada Thistle	○	○	○		Leafy Spurge	○	○	○		Other:	○	○	○	
										Other:	○	○	○	

PLOT COORDINATES

Provide GPS coordinates at the center of the Buffer Plot (#3) at the far end of each Buffer Transect and for the Buffer Plot at the AA CENTER. Indicate the location of the plot coordinates by filling in the appropriate bubble.

If Buffer Plot 3 can not be accessed, take the coordinates at the nearest practicable location ALONG THE TRANSECT. This is important because all Buffer Plots are centered on the Buffer Transects and the coordinates will indicate the location of the transect. Fill in the "nearest practicable location" bubble, fill in the flag box, and describe where the coordinates were taken and why in the comment section below. The coordinates of the nearest practicable location can be either placed as close to the center of Plot 3 as possible or at the center of the last accessible Buffer Plot.

Location of coordinates (choose one): Flag

○ AA CENTER ○ N3 ● S3 ○ E3 ○ W3 ○ Nearest practicable location (flag and comment below)

Latitude North 41.31347 Longitude West 081.61908

Use Decimal Degrees; NAD83

Flag	Comments
1	plot #1 falls on a slope (20-30°)
2	plot #1 falls on margin of a road (2 lane, paved)
3	plot #2 falls 10m away from paved 2 lane road
4	plot #2 falls 15 m from culuert, gully runs into plot
5	plot #3 falls on a steep (≈45°) slope

FORM B-1: BUFFER SAMPLE PLOTS (Front)

Reviewed by (initial): _____

Site ID: PCAPBR1201 DATE: 06/20/2012

Location: ○ AA Center ○ N ○ S ● E ○ W

Fill in bubble(s) if plot(s) could not be sampled and flag → ○ Plot 1 ○ Plot 2 ○ Plot 3

Buffer Natural Cover Strata

Fill in bubbles for all that apply: Canopy Type: D = Deciduous; E = Evergreen. Leaf Type: B = Broadleaf; N = Needle Leaf. Absent: No tree canopy.
Strata Section. Fill in appropriate cover class bubble for each strata type for each plot. 0 = Absent; 1 = Sparse(<10%); 2=Moderate(10-40%); 3 = Heavy (40-75%); 4 = Very Heavy (>75%)

Buffer Plot 1	Canopy Type: D ● E	Absent: ○	Leaf Type: B ● N	Flag	Buffer Plot 2	Canopy Type: D ● E	Absent: ○	Leaf Type: B ● N	/ Flag	Buffer Plot 3	Canopy Type: D ● E	Absent: ○	Leaf Type: B ● N	Flag
Big Trees (>0.3m DBH)	0 ● 2 3 4				Big Trees (>0.3m DBH)	0 1 ● 3 4				Big Trees (>0.3m DBH)	0 ● 2 3 4			
Small Trees (<0.3m DBH)	● 1 2 3 4				Small Trees (<0.3m DBH)	● 1 2 3 4				Small Trees (<0.3m DBH)	● 1 2 3 4			
Woody Shrubs, Saplings (0.5m-5m HIGH)	● 1 2 3 4				Woody Shrubs, Saplings (0.5m-5m HIGH)	● 1 2 3 4				Woody Shrubs, Saplings (0.5m-5m HIGH)	● 1 2 3 4			
Woody Shrubs, Saplings (<0.5m HIGH)	● 1 2 3 4				Woody Shrubs, Saplings (<0.5m HIGH)	● 1 2 3 4				Woody Shrubs, Saplings (<0.5m HIGH)	● 1 2 3 4			
Herbs, Forbs and Grasses	0 1 2 3 ●				Herbs, Forbs and Grasses	0 1 2 ● 4				Herbs, Forbs and Grasses	0 ● 2 3 4			
Bare ground	0 ● 2 3 4				Bare ground	● 1 2 3 4				Bare ground	● 1 2 3 4			
Litter, duff	● 1 2 3 4				Litter, duff	● 1 2 3 4				Litter, duff	● 1 2 3 4			
Rock	● 1 2 3 4				Rock	● 1 2 3 4				Rock	● 1 2 3 4			
Water	● 1 2 3 4				Water	● 1 2 3 4				Water	● 1 2 3 4			
Submerged Vegetation	● 1 2 3 4				Submerged Vegetation	● 1 2 3 4				Submerged Vegetation	● 1 2 3 4			

Stressor Presence/Absence - Confirm that a filled data bubble indicates presence and an unfilled bubble indicates absence by filling this bubble. ☺

Residential and Urban Stressors

Fill bubble if present - Plot	1	2	3	Flag
Road - gravel	○	○	○	
Road - two lane	●	○	●	
Road - four lane	○	○	○	
Parking Lot/Pavement	○	●	●	
Golf Course	○	○	○	
Lawn/Park	●	●	●	
Suburban Residential	○	○	○	
Urban/Multifamily	○	○	○	
Landfill	○	○	○	
Dumping	○	○	○	
Trash	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Hydrology Stressors

Fill bubble if present - Plot	1	2	3	Flag
Ditches, Channelization	○	○	○	
Dike/Dam/Road/RR Bed (IMPEDE FLOW)	○	○	○	
Water Level Control Structure	○	○	○	
Excavation, Dredging	○	○	○	
Fill/Spoil Banks	○	○	○	
Freshly Deposited Sediment (UNVEGETATED)	○	○	○	
Soil Loss/Root Exposure	○	○	○	
Wall/Riprap	○	○	○	
Inlets, Outlets	○	○	○	
Point Source/Pipe (EFFLUENT OR STORMWATER)	○	○	○	
Impervious surface input (SHEETFLOW)	●	●	●	
Other: _____	○	○	○	
Other: _____	○	○	○	

Agricultural & Rural Stressors

Fill bubble if present - Plot	1	2	3	Flag
Pasture/Hay	○	○	○	
Range	○	○	○	
Row Crops	○	○	○	
Fallow Field (RECENT-RESTING ROW CROP FIELD)	○	○	○	
Fallow Field (OLD - GRASS, SHRUBS, TREES)	○	○	○	
Nursery	○	○	○	
Dairy	○	○	○	
Orchard	○	○	○	
Confined Animal Feeding	○	○	○	
Rural Residential	○	○	○	
Gravel Pit	○	○	○	
Irrigation	○	○	○	
Other: _____	○	○	○	

Industrial Development Stressors

Fill bubble if present - Plot	1	2	3	Flag
Oil Drilling	○	○	○	
Gas Wells	○	○	○	
Mine (surface)	○	○	○	
Mine (underground)	○	○	○	
Military	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Habitat/Vegetation Stressors

Fill bubble if present - Plot	1	2	3	Flag
Forest Clear Cut	○	○	○	
Forest Selective Cut	○	○	○	
Tree Plantation	○	○	○	
Tree Canopy Herbivory (INSECT)	○	○	○	
Shrub Layer Browsed (WILD OR DOMESTIC)	○	○	○	
Highly Grazed Grasses (OVERALL <3" HIGH)	○	○	○	
Recently Burned Forest Canopy	○	○	○	
Recently Burned Grassland (BLACKENED)	○	○	○	

Fill bubble if present - Plot	1	2	3	Flag
Herbicide Use	○	○	○	
Mowing/Shrub Cutting	●	●	●	
Trails	○	○	○	
Soil Compaction (ANIMAL OR HUMAN)	○	○	○	
Offroad vehicle damage	○	○	○	
Soil erosion (FROM WIND, WATER, OR OVERUSE)	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Flag codes: K = No measurement made, U = Suspect measurement., F1,F2, etc. = misc. flags assigned by each field crew.
Explain all flags in comment section on the back of this form

2428168304

FORM B-1: BUFFER SAMPLE PLOTS - TARGETED ALIEN SPECIES (Back)

Reviewed by (initial): _____

Site ID: PCAPBR1201 DATE: 06/20/2012

Confirm a filled data bubble indicates presence and an unfilled bubble indicates absence by filling in this bubble

Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag
Eurasian Watermilfoil	○	○	○		Purple Loosestrife	○	○	○		Johnson Grass	○	○	○	
Water hyacinth	○	○	○		Knotweed	○	○	○		Kudzu	○	○	○	
Yellow Floating Heart	○	○	○		Japanese Knotweed	○	○	○		Multiflora Rose	○	○	○	
Giant Salvinia	○	○	○		Perennial Pepperweed	○	○	○		Common Buckthorn	○	○	○	
Garlic Mustard	○	○	○		Giant Reed	○	○	○		Himalayan Blackberry	○	○	○	
Poison Hemlock	○	○	○		Cheatgrass	○	○	○		Tamarisk	○	○	○	
Mile-A-Minute Weed	○	○	○		Reed Canary Grass	○	○	○		Other:	○	○	○	
Birdsfoot Trefoil	○	○	○		Common Reed	○	○	○		Other:	○	○	○	
Canada Thistle	○	○	○		Leafy Spurge	○	○	○		Other:	○	○	○	
										Other:	○	○	○	

PLOT COORDINATES

Provide GPS coordinates at the center of the Buffer Plot (#3) at the far end of each Buffer Transect and for the Buffer Plot at the AA CENTER. Indicate the location of the plot coordinates by filling in the appropriate bubble.

If Buffer Plot 3 can not be accessed, take the coordinates at the nearest practicable location ALONG THE TRANSECT. This is important because all Buffer Plots are centered on the Buffer Transects and the coordinates will indicate the location of the transect. Fill in the "nearest practicable location" bubble, fill in the flag box, and describe where the coordinates were taken and why in the comment section below. The coordinates of the nearest practicable location can be either placed as close to the center of Plot 3 as possible or at the center of the last accessible Buffer Plot.

Location of coordinates (choose one):

○ AA CENTER ○ N3 ○ S3 ● E3 ○ W3 ○ Nearest practicable location (flag and comment below)

Flag: ____

Latitude North 41 . 31499 Longitude West 081 . 61739

Use Decimal Degrees; NAD83

Flag	Comments
1	~50% of plot is asphalt parking area

7966623548

FORM B-1: BUFFER SAMPLE PLOTS (Front)

Reviewed by (initial): _____

Site ID: PCAPBR1201 DATE: 06/20/2012

Location: ○ AA Center ○ N ○ S ○ E ● W

Fill in bubble(s) if plot(s) could not be sampled and flag → ○ Plot 1 ○ Plot 2 ○ Plot 3

Buffer Natural Cover Strata

Fill in bubbles for all that apply: Canopy Type: D = Deciduous; E = Evergreen. Leaf Type: B = Broadleaf; N = Needle Leaf. Absent: No tree canopy.
Strata Section: Fill in appropriate cover class bubble for each strata type for each plot. 0 = Absent; 1 = Sparse(<10%); 2=Moderate(10-40%); 3 = Heavy (40-75%); 4 = Very Heavy (>75%)

Buffer Plot 1	Canopy Type: D ● E ○ Absent: ○ Leaf Type: B ● N ○					
	0	1	2	3	4	Flag
Big Trees (>0.3m DBH)	○	●	○	○	○	
Small Trees (<0.3m DBH)	○	○	●	○	○	
Woody Shrubs, Saplings (0.5m-5m HIGH)	○	○	●	○	○	
Woody Shrubs, Saplings (<0.5m HIGH)	○	○	●	○	○	
Herbs, Forbs and Grasses	○	○	○	●	○	
Bare ground	○	●	○	○	○	
Litter, duff	○	○	●	○	○	
Rock	●	○	○	○	○	
Water	●	○	○	○	○	
Submerged Vegetation	●	○	○	○	○	

Buffer Plot 2	Canopy Type: D ● E ○ Absent: ○ Leaf Type: B ● N ○					
	0	1	2	3	4	Flag
Big Trees (>0.3m DBH)	○	●	○	○	○	
Small Trees (<0.3m DBH)	○	○	○	●	○	
Woody Shrubs, Saplings (0.5m-5m HIGH)	○	○	●	○	○	
Woody Shrubs, Saplings (<0.5m HIGH)	○	○	●	○	○	
Herbs, Forbs and Grasses	○	○	○	○	●	
Bare ground	○	●	○	○	○	
Litter, duff	○	○	●	○	○	
Rock	●	○	○	○	○	
Water	●	○	○	○	○	
Submerged Vegetation	●	○	○	○	○	

Buffer Plot 3	Canopy Type: D ● E ○ Absent: ○ Leaf Type: B ● N ○					
	0	1	2	3	4	Flag
Big Trees (>0.3m DBH)	○	○	○	●	○	
Small Trees (<0.3m DBH)	○	○	○	●	○	
Woody Shrubs, Saplings (0.5m-5m HIGH)	○	○	●	○	○	
Woody Shrubs, Saplings (<0.5m HIGH)	○	○	●	○	○	
Herbs, Forbs and Grasses	○	○	●	○	○	
Bare ground	○	○	●	○	○	
Litter, duff	○	○	○	●	○	
Rock	○	●	○	○	○	
Water	●	○	○	○	○	
Submerged Vegetation	●	○	○	○	○	

Stressor Presence/Absence - Confirm that a filled data bubble indicates presence and an unfilled bubble indicates absence by filling this bubble. ○

Residential and Urban Stressors

Fill bubble if present - Plot	1	2	3	Flag
Road - gravel	○	○	○	
Road - two lane	●	○	○	1
Road - four lane	○	○	○	
Parking Lot/Pavement	○	○	○	
Golf Course	○	○	○	
Lawn/Park	○	○	○	
Suburban Residential	○	○	○	
Urban/Multifamily	○	○	○	
Landfill	○	○	○	
Dumping	○	○	○	
Trash	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Hydrology Stressors

Fill bubble if present - Plot	1	2	3	Flag
Ditches, Channelization	○	○	○	
Dike/Dam/Road/RR Bed (IMPEDE FLOW)	○	○	○	
Water Level Control Structure	○	○	○	
Excavation, Dredging	○	○	○	
Fill/Spoil Banks	○	○	○	
Freshly Deposited Sediment (UNVEGETATED)	○	○	○	
Soil Loss/Root Exposure	○	○	●	3
Wall/Riprap	○	○	○	
Inlets, Outlets	○	○	○	
Point Source/Pipe (EFFLUENT OR STORMWATER)	○	○	○	
Impervious surface input (SHEETFLOW)	●	○	○	1
Other: _____	○	○	○	
Other: _____	○	○	○	

Agricultural & Rural Stressors

Fill bubble if present - Plot	1	2	3	Flag
Pasture/Hay	○	○	○	
Range	○	○	○	
Row Crops	○	○	○	
Fallow Field (RECENT-RESTING ROW CROP FIELD)	○	○	○	
Fallow Field (OLD - GRASS, SHRUBS, TREES)	○	○	○	
Nursery	○	○	○	
Dairy	○	○	○	
Orchard	○	○	○	
Confined Animal Feeding	○	○	○	
Rural Residential	○	○	○	
Gravel Pit	○	○	○	
Irrigation	○	○	○	
Other: _____	○	○	○	

Industrial Development Stressors

Fill bubble if present - Plot	1	2	3	Flag
Oil Drilling	○	○	○	
Gas Wells	○	○	○	
Mine (surface)	○	○	○	
Mine (underground)	○	○	○	
Military	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	
Other: _____	○	○	○	

Habitat/Vegetation Stressors

Fill bubble if present - Plot	1	2	3	Flag
Forest Clear Cut	○	○	○	
Forest Selective Cut	○	○	○	
Tree Plantation	○	○	○	
Tree Canopy Herbivory (INSECT)	○	○	○	
Shrub Layer Browsed (WILD OR DOMESTIC)	○	●	●	
Highly Grazed Grasses (OVERALL <3" HIGH)	○	○	○	
Recently Burned Forest Canopy	○	○	○	
Recently Burned Grassland (BLACKENED)	○	○	○	

Habitat/Vegetation Stressors

Fill bubble if present - Plot	1	2	3	Flag
Herbicide Use	○	○	○	
Mowing/Shrub Cutting	●	○	○	1
Trails	○	●	○	2
Soil Compaction (ANIMAL OR HUMAN)	○	●	○	2
Offroad vehicle damage	○	○	○	
Soil erosion (FROM WIND, WATER, OR OVERUSE)	○	○	●	3
Other: _____	○	○	○	
Other: _____	○	○	○	

Flag codes: K = No measurement made, U = Suspect measurement., F1,F2, etc. = misc. flags assigned by each field crew.
Explain all flags in comment section on the back of this form

2428168304

FORM B-1: BUFFER SAMPLE PLOTS - TARGETED ALIEN SPECIES (Back)

Reviewed by (initial): _____

Site ID: PCAPBR1201 DATE: 06/20/2012

● Confirm a filled data bubble indicates presence and an unfilled bubble indicates absence by filling in this bubble

Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag	Fill bubble if present - Plot	1	2	3	Flag
Eurasian Watermilfoil	○	○	○		Purple Loosestrife	○	○	○		Johnson Grass	○	○	○	
Water hyacinth	○	○	○		Knotweed	○	○	○		Kudzu	○	○	○	
Yellow Floating Heart	○	○	○		Japanese Knotweed	○	○	○		Multiflora Rose	○	●	○	
Giant Salvinia	○	○	○		Perennial Pepperweed	○	○	○		Common Buckthorn	○	○	○	
Garlic Mustard	○	○	○		Giant Reed	○	○	○		Himalayan Blackberry	○	○	○	
Poison Hemlock	○	○	○		Cheatgrass	○	○	○		Tamarisk	○	○	○	
Mile-A-Minute Weed	○	○	○		Reed Canary Grass	○	○	○		Other:	○	○	○	
Birdsfoot Trefoil	○	○	○		Common Reed	○	○	○		Other:	○	○	○	
Canada Thistle	○	○	○		Leafy Spurge	○	○	○		Other:	○	○	○	
										Other:	○	○	○	

PLOT COORDINATES

Provide GPS coordinates at the center of the Buffer Plot (#3) at the far end of each Buffer Transect and for the Buffer Plot at the AA CENTER. Indicate the location of the plot coordinates by filling in the appropriate bubble.

If Buffer Plot 3 can not be accessed, take the coordinates at the nearest practicable location ALONG THE TRANSECT. This is important because all Buffer Plots are centered on the Buffer Transects and the coordinates will indicate the location of the transect. Fill in the "nearest practicable location" bubble, fill in the flag box, and describe where the coordinates were taken and why in the comment section below. The coordinates of the nearest practicable location can be either placed as close to the center of Plot 3 as possible or at the center of the last accessible Buffer Plot.

Location of coordinates (choose one):

○ AA CENTER ○ N3 ○ S3 ○ E3 ● W3 ○ Nearest practicable location (flag and comment below)

Flag: ☐

Latitude North 41.31409 Longitude West 81.62025

Use Decimal Degrees; NAD83

Flag	Comments
1	2 lane paved road, 1 m from plot
2	deer trail falls in plot
3	plot # 3 falls on a 30° - 40° slope