

# Exploring druggability with the JEDI collective variable

Joan Clark-Nicolas  
PLUMED users meeting  
25<sup>th</sup> – 27<sup>th</sup> May 2017  
SISSA Trieste (Italy)

# Contents



- **Part 1: Protein Druggability**
- **Part 2: Dynamics**
- **Part 3: Biased MD**
- **Part 4: Performance**
- **Part 5: Conclusions and Outlook**

# Part 1: Protein Druggability






**What are proteins and diseases?  
And how are drugs related to them?**

# How are diseases related to proteins?



*"From a simplified perspective, proteins can be considered as very complex machines that perform a function inside a cell..."*




*"...and diseases can be considered the result of a protein not performing its function properly"*

# How do drugs cure diseases?




*"From a simplified perspective, proteins can be considered as very complex machines that perform a function inside a cell..."*



*"...and diseases can be considered the result of a protein not performing its function properly."*

*"A drug is a molecule whose role is **usually** to correct this malfunction by binding to a target protein and modulating its activity"*

# Not all proteins can bind drugs...




~10% human genome is involved in diseases

~20% - 50% of human genes involved in diseases code proteins targeted by drugs

Hopkins & Groom, Nat. Rev. Drug. Disc. 2002

**Druggability: the ability of a protein to have its activity modulated by the non-covalent binding of a small molecule**

# Can protein flexibility help drug design?





## Inhibitor SKF

Volume = 258 Å<sup>3</sup>

K<sub>i</sub> = 580 nM

Pocket Volume: 304 Å<sup>3</sup>




## Inhibitor F83

Volume = 422 Å<sup>3</sup>



K<sub>i</sub> = 63 nM

Pocket Volume: 545 Å<sup>3</sup>





# How to find more potent ligands?



**Druggability: *the ability of a protein to BIND a small molecule***

JCTC


Journal of Chemical Theory and Computation

Article

[pubs.acs.org/JCTC](https://pubs.acs.org/JCTC)

## A Collective Variable for the Rapid Exploration of Protein Druggability

Rémi Cuchillo, Kevin Pinto-Gil, and Julien Michel\*





# Quantification of Protein Druggability: JEDI



JCTC



Journal of Chemical Theory and Computation

Article

[pubs.acs.org/JCTC](https://pubs.acs.org/JCTC)


## A Collective Variable for the Rapid Exploration of Protein Druggability

Rémi Cuchillo, Kevin Pinto-Gil, and Julien Michel\*



$$a_i = S_{BS_i}^{off}(1.0, BS_{min}, \Delta BS) S_{mind_i}^{on}(1.0, CC_{mind}, \Delta CC) S_{exposure_i}^{on}(1.0, E_{min}, \Delta E)$$

Kernel Functions



Michel, J., JEDI Theory, Feb. 2016

Activity calculation

$$a_i = S_{BS_i}^{off}(1.0, BS_{min}, \Delta BS) S_{mind_i}^{on}(1.0, CC_{mind}, \Delta CC) S_{exposure_i}^{on}(1.0, E_{min}, \Delta E)$$

JEDI score

$$JEDI = V_{druglike}(\alpha V_a + \beta H_a + \gamma)$$

Exposure of grid point  $i$  to the

Active De

$$V = \sum_{i=1}^N a_i V_g$$

$$V_g = spacing^3 \longrightarrow$$

$$V_a = \frac{V}{V_{max}}$$

$$PAR V_{druglike} = S_V^{off}(1.0, V_{max}, \Delta V_{max}) S_V^{on}(1.0, V_{min}, \Delta V_{min})$$

at the same time as

Penalize pockets that are too big

Penalize pockets that are too small



# Quantification of Protein Druggability: JEDI



JCTC



Journal of Chemical Theory and Computation

Article

[pubs.acs.org/JCTC](https://pubs.acs.org/JCTC)

## A Collective Variable for the Rapid Exploration of Protein Druggability

Rémi Cuchillo, Kevin Pinto-Gil, and Julien Michel\*



$$a_i = S_{BS_i}^{off}(1.0, BS_{min}, \Delta BS) S_{mind_i}^{on}(1.0, CC_{mind}, \Delta CC) S_{exposure_i}^{on}(1.0, E_{min}, \Delta E)$$

$$JEDI = F(a_{gridpoints})$$

$$a_i = F(\vec{r}_{gridpoints}, \vec{r}_{atoms})$$

$$JEDI = F(\vec{r}_{gridpoints}, \vec{r}_{atoms})$$

CONTINUOUS FUNCTION

CONTINUOUS DERIVATIVES

# JEDI vs Experimental druggability



Non redundant dataset


Proteins chosen and scored manually

Discrete scores


1-4: Non druggable (few or no known ligands)

5-7: Difficult (some known ligands)


8-10: Druggable (many known ligands)



# Can JEDI distinguish hPNMT conformations?



Pocket	Small
LIGAND	SKF (580nM)
JEDI	7.3219 (8.17*)
Vd	1
Va	0.61497
Ha	0.71826



Pocket	Big
LIGAND	F83 (63 nM)
JEDI	9.8087 (8.98*)
Vd	1
Va	0.82106
Ha	0.77558



## Part 2: Dynamics



**Can we predict the binding mode of the big ligand (F83) if knowing only the structure bound to the small one (SKF)?**

# Druggability vs time



JCTC

Journal of Chemical Theory and Computation

Article

[pubs.acs.org/JCTC](https://pubs.acs.org/JCTC)

## A Collective Variable for the Rapid Exploration of Protein Druggability


Rémi Cuchillo, Kevin Pinto-Gil, and Julien Michel\*



**PLUMED**



**GROMACS**  
FAST. FLEXIBLE. FREE.






# Grid Update



If the atoms move, we need to update the grid according to that movement!



- 1) Calculate binding site  $COM_0$  at time  $t_0$
- 2) Move atoms
- 3) calculate binding site COM at time  $t=t_0+dt$
- 4) Center binding site at time  $t$  at the COM at time  $t_0$  (not shown)
- 5) Calculate the rotation matrix  $R$  that fits the translated binding site at time  $t$  to the binding site at time  $t_0$  with the lowest RMSD


# Behaviour of the JEDI scores in MD

## MD Setup details

- 1 ns MD at 300 K
- $dt=0.002$  ps
- Explicit solvent (TIP3P)
- Ligand cavities filled with water
- Saving *JEDI* every 1000 steps
- GROMACS 5.1<sup>4</sup> + PLUMED 2.2b<sup>5</sup>

<https://github.com/michellab/plumed2>


<https://github.com/michellab/jedi-utilities>




**Short simulations don't show big fluctuations of the JEDI score and the correlation does not decrease significantly**

# Sampling of hPNMT conformations


Big Pocket (liganded)




Big Pocket (unliganded)




Small Pocket (liganded)



Small Pocket (unliganded)



JEDI



## Part 3: Biased MD



**How to make hPNMT sample the conformation that binds the big ligand (F83) from the one that binds the small one (SKF)?**

# Biasing Potential and Force



## CONTINUOUS FUNCTION

$$JEDI = F(\vec{r}_{gridpoints}, \vec{r}_{atoms})$$



$$U_{JEDI} = F(JEDI) = F(\vec{r}_{gridpoints}, \vec{r}_{atoms})$$



$$U_{system} = U_{FF} + U_{JEDI}$$

## DIFFERENTIABLE FUNCTION

$$\frac{d(JEDI)}{dx_{atom_i}} = (...)$$



$$\vec{F}_{x_i} = -\frac{d(U_{system})}{dx_i} = -\left[ \frac{d(U_{FF})}{dx_i} + \frac{d(U_{JEDI})}{dx_i} \right]$$



"Classic"  
MD  
force



JEDI  
bias  
force



# Biasing Potential and Force



## HARMONIC POTENTIAL

$$U_{JEDI} = k(JEDI - JEDI_0)^2$$



$$\vec{F}_{JEDI_i} = -\frac{d[k(JEDI - JEDI_0)^2]}{dx_i} = -2k(JEDI - JEDI_0) \frac{d(JEDI)}{dx_i}$$

### IN COMPARISON TO MD FORCES WE WANT JEDI FORCES TO BE:

- Slowly varying absolute value (difference in force norm between steps)
- Slowly varying in direction (angle between force vectors of consecutive steps)




# Removal of net force and torque



Grid point i  
pushes atom j...

...but atom j does  
not push back



$$F_{ij} + F_{ji} \neq 0$$

$$\nabla JEDI_x = \sum_{j=1}^M \frac{\partial JEDI}{\partial x_{p_j}}$$

$$\nabla JEDI_y = \sum_{j=1}^M \frac{\partial JEDI}{\partial y_{p_j}}$$

$$\nabla JEDI_z = \sum_{j=1}^M \frac{\partial JEDI}{\partial z_{p_j}}$$

$$\nabla TorJEDI_x = \sum_{j=1}^M y_{p_j} \frac{\partial JEDI}{\partial z_{p_j}} - z_{p_j} \frac{\partial JEDI}{\partial y_{p_j}}$$

$$\nabla TorJEDI_y = \sum_{j=1}^M z_{p_j} \frac{\partial JEDI}{\partial x_{p_j}} - x_{p_j} \frac{\partial JEDI}{\partial z_{p_j}}$$

$$\nabla TorJEDI_z = \sum_{j=1}^M x_{p_j} \frac{\partial JEDI}{\partial y_{p_j}} - y_{p_j} \frac{\partial JEDI}{\partial x_{p_j}}$$

Type equation here


$$\neq \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

# Removal of net force and torque



Grid point i  
pushes atom j...

...but atom j does  
not push back



$$F_{ij} + F_{ji} \neq 0$$

$$\begin{pmatrix} \nabla JEDI_x^* = \sum_{j=1}^M \frac{\partial JEDI}{\partial x_{p_j}} + z_{3j} \\ \nabla JEDI_y^* = \sum_{j=1}^M \frac{\partial JEDI}{\partial y_{p_j}} + z_{3j+1} \\ \nabla JEDI_z^* = \sum_{j=1}^M \frac{\partial JEDI}{\partial z_{p_j}} + z_{3j+2} \\ \nabla TorJEDI_x^* = \sum_{j=1}^M y_{p_j} \left( \frac{\partial JEDI}{\partial z_{p_j}} + z_{3j+2} \right) - z_{p_j} \left( \frac{\partial JEDI}{\partial y_{p_j}} + z_{3j+1} \right) \\ \nabla TorJEDI_y^* = \sum_{j=1}^M z_{p_j} \left( \frac{\partial JEDI}{\partial x_{p_j}} + z_{3j} \right) - x_{p_j} \left( \frac{\partial JEDI}{\partial z_{p_j}} + z_{3j+2} \right) \\ \nabla TorJEDI_z^* = \sum_{j=1}^M x_{p_j} \left( \frac{\partial JEDI}{\partial y_{p_j}} + z_{3j+1} \right) - y_{p_j} \left( \frac{\partial JEDI}{\partial x_{p_j}} + z_{3j} \right) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

Undetermined Linear Equation system:  $Ax=y$

# Removal of net force and torque



1) Calculate The Moore-Penrose pseudo inverse matrix  $\mathbf{A}^+ = \mathbf{A}^t(\mathbf{A}\mathbf{A}^t)^{-1}$

2) Solving the linear equation system  $\mathbf{z} = \mathbf{A}^+ \mathbf{y}$

The solution of 2) minimizes the sum:

$$\|\mathbf{z}\| = \sum_{i=1}^{3M} z_i^2$$

$$\frac{\partial JEDI^*}{\partial x_{pj}} = \frac{\partial JEDI}{\partial x_{pj}} + z_{3j}$$


$$\frac{\partial JEDI^*}{\partial y_{pj}} = \frac{\partial JEDI}{\partial y_{pj}} + z_{3j+1}$$

$$\frac{\partial JEDI^*}{\partial z_{pj}} = \frac{\partial JEDI}{\partial z_{pj}} + z_{3j+2}$$

# Behaviour of the JEDI forces



Change in the absolute value of the force between consecutive steps




# Behaviour of the JEDI forces




Change in the direction of the force between consecutive steps

JEDI Forces



MD Forces



Difference in the force  
[kJ/(mol\*nm)]

## Part 4: Performance




**Can high druggability conformations  
be sampled faster using JEDI?**



# Serial performance




1 GPU / 4 CPU



```
for atom in binding_site:
    do_stuff()
    for point in grid:
        if point too_far_from atom:
            skip_point()
        endif
        do_more_stuff()
        for neighbour in neighbours[point]:
            do_even_more_stuff()
        endfor
    endfor
    still_more_stuff()
    for point in grid:
        just_a_bit_more()
    endfor
    last_things_to_do()
endfor
```

$$maxIter = M \left[ \sum_{i=0}^N (1 + neighbours[i]) \right]$$

# Parallel performance using openMP



Up to 3x-6x increase in speed using openMP

```
#pragma omp parallel for
```

```
for atom in binding_site:
```

```
do_stuff()
```

```
for point in grid:
```

```
if point too_far_from atom:
```

```
skip_point()
```

```
endif
```

```
do_more_stuff()
```

```
for neighbour in neighbours[point]:
```

```
do_even_more_stuff()
```

```
endfor
```

```
endfor
```

```
still_more_stuff()
```

```
for point in grid:
```

```
just_a_bit_more()
```

```
endfor
```

```
last_things_to_do()
```

```
endfor
```

$$maxIter = M \left[ \sum_{i=0}^N (1 + neighbours[i]) \right]$$

## Part 5: Conclusions and Outlook



**How to make JEDI still better? What to do with it then?**

# Conclusions



- JEDI shows a proper correlation with manually assigned druggability scores and allows to distinguish two conformations of the binding site of hPNMT
- The JEDI score is stable and no big fluctuations are seen in short MD simulations
- Different conformations of hPNMT can show similar JEDI scores in long MD simulations
- The biasing forces are stable in short biased simulations
- The performance of the code is still to be improved

# Further improvement



JCTC

Journal of Chemical Theory and Computation

Article


[pubs.acs.org/JCTC](https://pubs.acs.org/JCTC)

## Accurate Multiple Time Step in Biased Molecular Simulations

Marco Jacopo Ferrarotti, Sandro Bottaro, Andrea Pérez-Villa, and Giovanni Bussi\*

Scuola Internazionale Superiore di Studi Avanzati (SISSA), via Bonomea 265, 34136 Trieste, Italy

Evaluation of forces every  $n$  steps instead of 1 to boost performance



Overlapping a big grid onto the whole protein and clustering the grid points in order to detect the opening of transient pockets

# Application to hPNMT




- Perform JEDI-biased simulations of hPNMT to search for the desired conformational change
- Perform free energy calculations to determine the population of the conformations with higher JEDI scores
- Perform docking calculations of known ligands in order to verify the results



# Acknowledgements



## Michel research group



Dr. Julien Michel  
Royal Society University Research Fellow  
Room 263  
School of Chemistry  
University of Edinburgh  
Joseph Black Building  
King's Buildings  
David Brewster Road  
Edinburgh  
EH9 3FJ  
United Kingdom

## Funding



European Research Council



### Antonia Mey

**Degree:**  
*PhD. Physics, University of Nottingham, 2013*

**Keywords:**  
Free energy calculations, Molecular simulation workflows



### Jordi Juárez Jiménez

**Degree:**  
*PhD. Biomedicine, University of Barcelona, 2014*


**Keywords:**  
Cyclophilins inhibition, Docking and scoring, Protein dynamics




# Acknowledgements



# What is JEDI?



# Kernel Functions



$$S_v^{on}(k, v_{min}, \Delta) = \begin{cases} 0 & \text{if } m < 0 \\ k[1 - (1 - m^2)^2(1 + 2m^2)] & \text{if } 0 \leq m \leq 1 \\ k & \text{if } m > 1 \end{cases}$$

$$S_v^{off}(k, v_{min}, \Delta) = \begin{cases} k & \text{if } m < 0 \\ k[(1 - m^2)^2(1 + 2m^2)] & \text{if } 0 \leq m \leq 1 \\ 0 & \text{if } m > 1 \end{cases}$$

$$m = \frac{v - v_{min}}{\Delta}$$

Michel, J., JEDI Theory, Feb. 2016

**Penalize the descriptors with values too big or too low regarding those of the benchmark set**

# Activity calculation

$$a_i = S_{BS_i}^{off}(1.0, BS_{min}, \Delta BS) S_{mind_i}^{on}(1.0, CC_{mind}, \Delta CC) S_{exposure_i}^{on}(1.0, E_{min}, \Delta E)$$

Distance of grid point  $i$  to a known ligand  
(1 if not defined)

Distance of grid point  $i$  to binding site atoms

Exposure of grid point  $i$  to the solvent



# Hydrophobicity Descriptor

$$h_i = \frac{apolar_i}{contacts_i}$$

$$contacts_i = \sum_{j=1}^M S_{\|r_{ij,t}\|}^{off}(a_i, d_{hydro}, \Delta d_{hydro})$$

$$apolar_i = \sum_{j=1}^M I_{apolar}(j) S_{\|r_{ij,t}\|}^{off}(a_i, d_{hydro}, \Delta d_{hydro})$$

$$I_{apolar}(j) \begin{cases} 1 & \text{if } j \in \text{apolar group} \\ 0 & \text{if } j \in \text{polar group} \end{cases}$$

$$H_a = \sum_{i=1}^N \frac{h_i a_i}{\sum_{i=1}^N a_i}$$

**Apolar and polar atoms are defined at the same time as the grid, before the simulation**



# Active Volume Descriptor

$$V = \sum_{i=1}^N a_i V_g$$

$$V_g = \textit{spacing}^3 \longrightarrow \text{ONLY FOR EVENLY SPACED GRIDS!!!!}$$

$$V_a = \frac{V}{V_{max}} \longrightarrow \text{PARAMETER}$$

# Druglike Volume Descriptor

$$V_{druglike} = S_V^{off}(1.0, V_{max}, \Delta V_{max}) S_V^{on}(1.0, V_{min}, \Delta V_{min})$$



**Penalize  
pockets that  
are too big**



**Penalize  
pockets that  
are too small**

# JEDI score

$$JEDI = V_{druglike}(\alpha V_a + \beta H_a + \gamma)$$

Symbol	Definition	Value
$\alpha$	PLS derived volume coefficient	5.31
$\beta$	PLS derived hydrophobicity coefficient	24.29
$\gamma$	PLS derived intercept	-13.39
$\Delta_g$	grid spacing	0.15 nm
$BS_{min}$	Minimum distance to <i>ligand group</i> from which the $lig_i$ value starts to decrease	0.2 nm
$\Delta BS$	distance interval over which $lig_i$ decreases to 0	0.6 nm
$CC_{mind}$	distance below which a grid point is fully in close contact with the <i>binding site group</i>	0.15 nm
$\Delta CC$	distance interval over which a grid point is in partial contact with the <i>binding site group</i>	0.15 nm
$E_{min}$	minimum exposure value from which a grid point is considered to be partially exposed to the <i>binding site group</i>	10.0
$\Delta E$	interval over which a grid point becomes fully exposed to the <i>binding site group</i>	20.0
$CC2_{min}$	minimum distance below which a grid point is overlapping the binding site group	0.15 nm
$\Delta CC2$	distance interval over which a grid point is in partial contact with the binding site group	0.14 nm

Symbol	Definition	Value
$GP_{min}$	distance above which a grid point $k$ is considered neighbor of grid point $i$	0.25 nm
$GP_{max}$	distance below which a grid point $k$ is considered neighbor of grid point $i$	0.35 nm
$d_{hydro}$	distance below which a grid point $i$ is in contact with a binding site atom (for hydrophobicity calculation)	0.40 nm
$\Delta d_{hydro}$	distance interval over which a grid point $i$ is in partial contact with a binding site atom (for hydrophobicity calculation)	0.05 nm
$V_{max}$	volume below which $V_{druglike}$ is equal to 1	0.5 nm <sup>3</sup>
$\Delta V_{max}$	volume interval over which $V_{druglike}$ goes from 1 to 0	0.050 nm <sup>3</sup>
$V_{min}$	volume below which $V_{druglike}$ is equal to 0	0.0 nm <sup>3</sup>
$\Delta V_{min}$	volume interval over which $V_{druglike}$ goes from 0 to 1	0.050 nm <sup>3</sup>

# JEDI version 1

- Written in C
- Parameters hard-coded
- Apolar and polar atoms definition hard coded
- Works in implicit solvent
- Gromacs 4.5.5
- PLUMED 1.3

# JEDI version 2

- Written in C++
- Parameters supplied by *jedi.params*
- Grid, apolar and polar atoms file generated with a preprocessor *jedi\_setup.py*
- Handles explicit solvent and PBC
- Gromacs 5.1.0
- PLUMED 2.2