

```

data wide_lead;
input ID TRT $ PB1 - PB4;
datalines;
  1      P      30.8      26.9      25.8      23.8
  2      A      26.5      14.8      19.5      21.0
..... data omitted .....
;
run;

proc print data=wide_lead;
run;

```

Obs	ID	TRT	PB1	PB2	PB3	PB4
1	1	P	30.8	26.9	25.8	23.8
2	2	A	26.5	14.8	19.5	21.0
3	3	A	25.8	23.0	19.1	23.2
4	4	P	24.7	24.5	22.0	22.5
5	5	A	20.4	2.8	3.2	9.4

*Now we'll change the wide formatted data to long formatted;

```

data long_lead;
set wide_lead;
  week=0;
  PB=PB1;
  output;
  week=1;
  PB=PB2;
  output;
  week=4;
  PB=PB3;
  output;
  week=6;
  PB=PB4;
  output;
  drop PB1-PB4;
run;

proc print data=long_lead (obs=10);
run;

```

Obs	ID	TRT	week	PB
1	1	P	0	30.8
2	1	P	1	26.9
3	1	P	4	25.8
4	1	P	6	23.8
5	2	A	0	26.5

Obs	ID	TRT	week	PB
6	2	A	1	14.8
7	2	A	4	19.5
8	2	A	6	21.0
9	3	A	0	25.8
10	3	A	1	23.0

```

data long_lead2;
  set wide_lead;
  array APB(1:4) PB1-PB4;
  array Aweek(1:4) (0 1 4 6);
  do i=1 to 4;
    PB = APB[i];
    week = Aweek[i];
    output;
  end;
  drop PB1-PB4 Aweek1 - Aweek4 i;
run;
proc print data=long_lead2 (obs=10);
run;


```

Obs	ID	TRT	week	PB
1	1	P	0	30.8
2	1	P	1	26.9
3	1	P	4	25.8
4	1	P	6	23.8
5	2	A	0	26.5
6	2	A	1	14.8
7	2	A	4	19.5
8	2	A	6	21.0
9	3	A	0	25.8
10	3	A	1	23.0


```

*First we'll do a simple "spagetti plot" of the data;
Proc SGplot data = long_lead;
series x=week y=PB / group =ID LineAttrs= (pattern=1 color="black");
run;


```



```
Proc SGplot data = long_lead;
series x=week y=PB / group =ID LineAttrs= (pattern=1);
run;
```



```
Proc SGplot data = long_lead;
series x=week y=PB / group =ID;
run;
```



*Now we will include the mean line on the graph;

```
proc sort data=long_lead;
by week;
```

*Calculate the mean by week;

```
proc means mean data=long_lead;
by week;
var PB;
output out = MN_dat mean = mn_PB;
```


```
proc print data = MN_dat;
run;
```

*Stack the mean data onto the long dataset;

```
data stacked;
set long_lead MN_dat;
run;
```

Obs	week	_TYPE_	_FREQ_	mn_PB
1	0	0	100	26.406
2	1	0	100	19.091
3	4	0	100	19.792
4	6	0	100	22.204

```
Proc SGplot data = stacked;
series x=week y=PB / group =ID LineAttrs= (pattern=1 color="black");
series x=week y=mn_PB / LineAttrs= (pattern=1 color="blue" thickness=4);
run;
```





```

**We can also create separate plots by TRT group;
proc sort data=long_lead;
by TRT;

Proc SGplot data = long_lead;
by TRT;
series x=week y=PB / group =ID LineAttrs= (pattern=1);
run;


```



```

* Here we'll do separate plots (panels) for each TRT group;
Proc SGpanel data = long_lead;
PanelBy TRT / columns=2;
series x=week y=PB / group =ID LineAttrs= (pattern=1);
run;

```



*Now we will include the mean line on the graph by TRT;

```
proc sort data=long_lead;
by TRT week;
```


*Calculate the mean by week;

```
proc means mean data=long_lead;
by TRT week;
var PB;
output out = MN_TRT_dat mean = mn_TRT_PB;
```

```
proc print data = MN_TRT_dat;
run;
```

*First, let's look at the mean by TRT group;

```
Proc SGplot data = MN_TRT_dat;
series x=week y=mn_TRT_PB / group =TRT LineAttrs= (pattern=1 thickness=3);
run;
```



```


*Now we'll look at the means by TRT group with the rest of the data;
*Stack the mean data onto the long dataset;
data stacked_TRT;
set long_lead MN_TRT_dat;
run;

proc sort data=stacked_TRT;
by TRT week;

*First we'll do separate plots for each TRT group with the mean on each;
Proc SGplot data = stacked_TRT;
by TRT;
series x=week y=PB / group =ID LineAttrs= (pattern=1 color="black");
series x=week y=mn_TRT_PB / LineAttrs= (pattern=1 color="blue" thickness=4);
run;

* Now we'll combine them onto one plot with two panels;
Proc SGpanel data = stacked_TRT;
PanelBy TRT / columns=2;
series x=week y=PB / group =ID LineAttrs= (pattern=1 color="black");
series x=week y=mn_TRT_PB / LineAttrs= (pattern=1 color="blue" thickness=4);
run;

```




*Let's go back to the simple plot, but this time limit our sample.
First, we need to create a new variable to randomly thin our sample;

```
proc sort data=long_lead;
by ID week;
run;
```

```
data long_lead_rand;
set long_lead;
by ID week;
if (first.ID=1) then U=ranuni(37);
retain U;
run;
```

```
Proc SGplot data = long_lead_rand;
where U>0.90;
series x=week y=PB / group =ID LineAttrs= (pattern=1 color="black");
run;
```

```
*****
*****      Example Two (Unbalanced data)      *****
*****;
```

*The Six Cities Study of Air Pollution and Health was a longitudinal study designed to characterize lung growth as measured by changes in pulmonary function in children and adolescents, and the factors that influence lung function growth. A cohort of 13,379 children born on or after 1967 was enrolled in six communities across the U.S.: Watertown (Massachusetts), Kingston and Harriman (Tennessee), a section of St. Louis (Missouri), Steubenville (Ohio), Portage (Wisconsin), and Topeka (Kansas). Most children were enrolled in the first or second grade (between the ages of six and seven) and measurements of study participants were obtained annually until graduation from high school or loss to follow-up. At each annual examination, spirometry, the measurement of pulmonary function, was performed and a respiratory health questionnaire was completed by a parent or guardian.;


```
data air_pol;
input  ID Height Age INI_Height  INI_Age  Log_FEV1;
datalines;
```

1	1.20		9.3415	1.20	9.3415	0.21511
1	1.28		10.3929	1.20	9.3415	0.37156
1	1.33		11.4524	1.20	9.3415	0.48858
1	1.42		12.4600	1.20	9.3415	0.75142
1	1.48		13.4182	1.20	9.3415	0.83291
1	1.50		15.4743	1.20	9.3415	0.89200
1	1.52		16.3723	1.20	9.3415	0.87129
2	1.13		6.5873	1.13	6.5873	0.30748
2	1.19		7.6496	1.13	6.5873	0.35066


```
.....Data omitted .....
;
```

```
run;
*First, we'll take a couple of snapshots of the data;
```

```
Proc sgplot data = air_pol;
scatter x=Age y=Log_FEV1 / group =ID markerattrs=(color=black symbol=plus
size=5) ;
run;
```



```
Proc SGplot data = air_pol;
series x=Age y=Log_FEV1 / group =ID LineAttrs= (pattern=1 );
run;
```



*Plotting a mean line for unbalanced data is a bit more challenging and requires a loess line smoother;

```
proc loess data=air_pol plots=none;  
  ods output outputstatistics=out_pol;  
  model Log_FEV1=Age;  
run;
```

Independent Variable Scaling	
Scaling applied: None	
Statistic	Age
Minimum Value	6.43390
Maximum Value	18.69130

Optimal Smoothing Criterion	
AICC	Smoothing Parameter
-2.82204	0.30667

*Note: from here on we use the dataset "out_pol" and "DepVar" instead of "Log_FEV1";

```
proc sort data=out_pol;  
  by Age DepVar;  
run;
```

```
proc sgplot data=out_pol;  
  scatter x=Age y=DepVar/ markerattrs=(color=black size=5) ;  
  series x=Age y=pred/ lineattrs=(color=blue thickness=5) ;  
run;
```

