

Distributivity and the Commensurability of Value-Definite Realism and Empirical Adequacy

Patrick S. Mahon

Abstract

When does an empirically adequate theory admit a value-definite completion — an assignment of simultaneous definite values to all observables? The answer depends on the algebraic structure of the observations. For a directed system of Boolean algebras carrying compatible charges, the Stone construction produces an unconditional measure on the dual space, and every observable admits a simultaneous definite value via ultrafilters. The passage to a probability measure on an underlying state space requires only σ -additivity, and the choice of state space is unconstrained by the algebra [17].

For orthomodular lattices — the algebraic framework for incompatible observations — the situation is structurally different. The McDonald–Bimbó dual space carries principal filters as distinguished structure, and the Kochen–Specker theorem blocks global value-definiteness. Probabilistic realism (descent to a probability measure) remains available via Gleason-type results, but value-definite realism does not.

Distributivity guarantees value-definiteness; for the physically central case $L(H)$, $\dim H \geq 3$, non-distributivity blocks it. The state space admits a natural filtration $\mathcal{S} \supseteq \mathcal{S}_\sigma \supseteq \mathcal{S}_{\text{df}}$, whose three transitions — pasting, regularity, sharpness — are of different mathematical character. Distributivity collapses the filtration; non-distributivity can (but need not) open it.

2020 Mathematics Subject Classification. 03G12 (primary); 06C15, 81P10, 60A10 (secondary).

Keywords. orthomodular lattice, Stone duality, value-definiteness, Kochen–Specker theorem, Gleason’s theorem, empirical adequacy, constructive empiricism, Boolean algebra, directed system, σ -additivity.

1 Introduction

A central question in the philosophy of science asks when an empirically successful theory can be completed to a realist one — one that assigns definite properties to an underlying reality, not merely adequate predictions to observable phenomena. Van Fraassen [21] argues that science need only aim at *empirical adequacy*: agreement with all observable phenomena. The scientific realist demands more — an underlying state space whose properties generate the observations.

This debate is usually conducted in philosophical terms. We give it a mathematical formulation.

Three nested positions, each a strengthening of the previous:

- (i) **Empirical adequacy (EA).** A measure on the dual space of the observation algebra that agrees with all observed charges. No commitment to an underlying state space.
- (ii) **Probabilistic realism (PR).** A probability measure on an underlying state space $(\Omega, \sigma(\mathcal{C}))$ that generates the observations. Requires descent from the dual-space measure.
- (iii) **Value-definite realism (VDR).** Every observable simultaneously has a definite value. A global dispersion-free state — equivalently, a 2-valued homomorphism on the full observation algebra.

These are nested: VDR implies PR implies EA. Which implications reverse depends on the observation algebra:

Compatible observations. When all observations can be performed simultaneously, the observation algebra is a Boolean algebra — a complemented distributive lattice. Every Boolean algebra admits 2-valued homomorphisms (ultrafilters exist by Zorn’s lemma). In this setting, all three positions are available and mutually commensurable. The passage from EA to PR requires σ -additivity of the charges [17]; the further passage to VDR is free.

Incompatible observations. When some observations cannot be performed simultaneously — as in quantum mechanics, contextual experimental design, or distributed systems with interfering sensors — the observation algebra is an orthomodular lattice (OML), a complemented but non-distributive lattice. The Kochen–Specker theorem [16] shows that for the prototypical case $L(H)$ with $\dim H \geq 3$, no 2-valued homomorphism exists: VDR is blocked. Probabilistic realism remains available via Gleason’s theorem [8], and empirical adequacy is unconditional.

Distributivity guarantees value-definiteness; for $L(H)$ with $\dim H \geq 3$, its absence blocks it. The individual ingredients — Stone duality, the Kochen–Specker theorem, Gleason’s theorem, the Bub–Clifton uniqueness result, and the recent McDonald–Bimbó duality for orthomodular lattices [18] — are known. What has not been stated is the synthesis: that distributivity of the observation algebra is the condition controlling which of the three positions are available, and that the state space admits a filtration whose transitions are of different mathematical character. We begin with the Boolean case [17].

Let $(\iota, \leq)$ be a directed set indexing a family of Boolean algebras $\{B_i\}_{i \in \iota}$ with injective homomorphisms $\varphi_{ij} : B_i \rightarrow B_j$ for $i \leq j$. Write $\mathcal{C} = \bigcup_i B_i$ for the direct limit — the *cylinder algebra*.

A *compatible family of normalised charges* is a collection $\{\ell_i\}$ with each $\ell_i : B_i \rightarrow [0, 1]$ finitely additive, $\ell_i(1) = 1$, and

$$\ell_j \circ \varphi_{ij} = \ell_i \quad (i \leq j).$$

The Stone space $\text{St}(\mathcal{C})$ is the space of ultrafilters on $\mathcal{C}$, compact and totally disconnected. Each $\omega \in \Omega$ determines a principal ultrafilter

$$\text{pure}(\omega) = \{E \in \mathcal{C} : \omega \in E\}.$$

Any compatible family of normalised charges extends to a σ -additive Baire probability measure $\hat{\mu}$ on $\text{St}(\mathcal{C})$ — unconditionally, with no assumption of σ -additivity [3, Theorem 1], [17]. This is the *empirically adequate* object: it agrees with all charges on cylinder events.

The measure $\hat{\mu}$ lives on $\text{St}(\mathcal{C})$, not on Ω . Descent to a probability measure P on $(\Omega, \sigma(\mathcal{C}))$ requires

$$\hat{\mu}(\text{pure}(\Omega)) = 1,$$

i.e., that $\hat{\mu}$ concentrates on the principal ultrafilters. This holds if and only if each ℓ_i is σ -additive [17].

Every Boolean algebra admits lattice ultrafilters — maximal proper filters — by Zorn’s lemma. On a Boolean algebra, maximality forces a dichotomy: for every a , either $a \in F$ or $a^c \in F$. The proof extends F to include a^c whenever $a \notin F$; the extension remains proper because $a^c \wedge f \neq \perp$ for all $f \in F$, which follows from distributivity. This dichotomy makes every lattice ultrafilter a 2-valued homomorphism, assigning each $a \in \mathcal{C}$ a definite value in $\{0, 1\}$. On a non-distributive OML, the extension argument fails, and maximal proper filters need not be 2-valued.

Moreover, any subset $S \subseteq \text{St}(\mathcal{C})$ that projects surjectively onto each $\text{St}(B_i)$ can serve as a realisation space: set $\Omega = S$ with evaluation maps given by the restriction maps. The algebra imposes no constraint on the choice of Ω [17].

Proposition 1.1 (Boolean commensurability). *For a directed system of Boolean algebras with compatible charges:*

- (a) *EA is unconditional: $\hat{\mu}$ on $\text{St}(\mathcal{C})$ always exists.*
- (b) *PR holds iff each ℓ_i is σ -additive.*
- (c) *VDR holds unconditionally: $\text{St}(\mathcal{C})$ consists entirely of 2-valued homomorphisms.*

The only obstruction is σ -additivity at the EA–PR transition, and it is a condition on the charges, not the algebra.

Proof. Part (a) is the Stone construction [3, 17]. Part (b) is [17, Theorem 1]. Part (c): ultrafilters on a Boolean algebra are exactly the 2-valued homomorphisms; their existence follows from Zorn’s lemma applied to the filter generated by any non-zero element. $\square$

2 Orthomodular observation algebras

Proposition 1.1 depends on distributivity at exactly one point: the existence of 2-valued homomorphisms. When observations are incompatible, this fails. The non-distributive structure need not be postulated: Gunji et al. [9] show that gluing Boolean algebras along shared boundaries via categorical pushout produces an orthomodular lattice, generically non-distributive. The OML is *forced* by the context structure.

Definition 2.1. An *ortholattice* is a bounded lattice $A = (A, \wedge, \vee, {}^\perp, 0, 1)$ with an orthocomplementation ${}^\perp : A \rightarrow A$ satisfying $a \wedge a^\perp = 0$, $a \vee a^\perp = 1$, $a^{\perp\perp} = a$, and $a \leq b \Rightarrow b^\perp \leq a^\perp$.

Definition 2.2. An *orthomodular lattice* (OML) is an ortholattice satisfying the *orthomodular law*:

$$a \leq b \implies b = a \vee (a^\perp \wedge b).$$

Every Boolean algebra is an OML (the orthomodular law follows from distributivity), but the converse fails. An OML is Boolean if and only if it is distributive [18, Proposition 2.3].

Example 2.3. The lattice $L(H)$ of closed linear subspaces of a separable Hilbert space H with $\dim H \geq 3$, ordered by inclusion with $U^\perp = \{x \in H : \langle x, y \rangle = 0 \ \forall y \in U\}$, is an orthomodular lattice that is not distributive [15, Chapter 1].

Example 2.4 (Incompatible measurements). Three binary measurements A, B, C , where (A, B) and (A, C) can be performed jointly but (B, C) cannot. The jointly measurable pairs generate Boolean subalgebras; the full observation algebra is orthomodular, not Boolean.

Two elements a, b in an OML *commute* if they generate a Boolean subalgebra. A maximal set of pairwise commuting elements forms a *Boolean context* — a maximal Boolean subalgebra of A .

Definition 2.5. A *state* on an OML A is a function $s : A \rightarrow [0, 1]$ with $s(1) = 1$ and

$$a \perp b \implies s(a \vee b) = s(a) + s(b),$$

where $a \perp b$ means $a \leq b^\perp$.

A state is additive on *orthogonal* pairs only — not on arbitrary joins. This is the OML analogue of a finitely additive charge on a Boolean algebra.

Definition 2.6. A state s is *dispersion-free* if $s(a) \in \{0, 1\}$ for all $a \in A$.

A dispersion-free state is a 2-valued homomorphism: it assigns a definite truth value to every proposition in A . On a Boolean algebra, dispersion-free states are exactly the ultrafilters.

Theorem 2.7 (Kochen–Specker [16]). *If $\dim H \geq 3$, there is no dispersion-free state on $L(H)$.*

For $\dim H = 2$, dispersion-free states exist; the obstruction is specific to $\dim H \geq 3$.

Theorem 2.8 (Gleason [8]). *If $\dim H \geq 3$, every σ -additive state on $L(H)$ is of the form $s(P) = \text{tr}(\rho P)$ for a unique density operator ρ on H .*

The analogue of Stone duality for OMLs is the McDonald–Bimbó duality [18].

Let A be an OML. A *filter* on A is a non-empty subset $x \subseteq A$ that is upward closed ($a \in x$ and $a \leq b$ implies $b \in x$) and closed under finite meets ($a, b \in x$ implies $a \wedge b \in x$). A filter is *proper* if $0 \notin x$; the unique *improper* filter is $\omega = A$.

For each $a \in A$, the *principal filter* generated by a is $\uparrow a = \{b \in A : a \leq b\}$. Write $F(A)$ for the collection of all filters on A , and $P(A) \subseteq F(A)$ for the subcollection of principal filters.

Definition 2.9 (McDonald–Bimbó [18]). Let A be an OML. The *dual space* of A is the ordered relational topological space

$$S_0(A) = (F(A), \subseteq, \perp_A, P(A), \mathcal{T}),$$

where:

- (i) $F(A)$ is the set of all filters on A ;
- (ii) $\subseteq$ is set inclusion;
- (iii) $x \perp_A y$ iff there exists $a \in A$ with $a \in x$ and $a^\perp \in y$;
- (iv) $P(A)$ is the collection of principal filters;
- (v) $\mathcal{T}$ is the topology generated by the subbasis $\{h(a) : a \in A\} \cup \{F(A) \setminus h(a) : a \in A\}$, where $h(a) = \{x \in F(A) : a \in x\}$.

Theorem 2.10 (McDonald–Bimbó [18]). *$S_0(A)$ is an orthomodular space (compact, totally disconnected). The category of OMLs and homomorphisms is dually equivalent to the category of orthomodular spaces and continuous weak p -morphisms. Moreover, $A \cong \text{CO}(S_0(A))^\dagger$, the OML of clopen $\perp$ -stable subsets of $S_0(A)$.*

The critical differences from Stone duality:

Feature	Boolean (Stone)	OML (McDonald–Bimbó)
Dual points	Ultrafilters	All filters
Distinguished subset	None	$P(A)$ (principal filters)
Clopes	Boolean algebra	OML
Orthogonality	Trivial	Non-trivial ($\perp_A$)
2-valued homomorphisms	Plentiful (Zorn)	May not exist (KS)

In Stone duality, the principal filters (i.e., the principal ultrafilters $\text{pure}(\omega)$ for $\omega \in \Omega$) are *invisible*: they play no distinguished role in the dual space. Which ultrafilters are principal depends on the choice of Ω [17].

In the McDonald–Bimbó duality, $P(A)$ is *part of the structure* of the dual space: the orthomodular frame conditions (Definition 2.9) explicitly reference $P(A)$. This is the algebraic formalisation: for non-distributive algebras, the distinction between realised and unrealised

observation-patterns is constrained by the algebra. Distributivity is the condition under which this constraint vanishes.

$S_0(A)$ is compact, so the topological machinery applies. But the algebraic input is weaker. A state s on A defines

$$\mu(h(a)) = s(a)$$

on $\text{CO}(S_0(A))^\dagger$.

In the Boolean case, $\text{Clop}(\text{St}(\mathcal{C}))$ is a Boolean algebra of sets, and s is finitely additive on it. The Carathéodory–Choksi extension machinery applies: compactness of $\text{St}(\mathcal{C})$ gives σ -subadditivity, and the extension to a Baire measure is automatic.

In the OML case, $\text{CO}(S_0(A))^\dagger$ is an OML of sets — closed under $\cap$ and $^\perp$ but *not* under arbitrary $\cup$. The state s is additive on orthogonal pairs only. Carathéodory does not apply: there is no Boolean algebra of sets on which s is finitely additive.

For the prototypical case $A = L(H)$ with $\dim H \geq 3$, Gleason’s theorem (Theorem 2.8) supplies the extension for σ -additive states: every such state arises from a density operator, giving a σ -additive measure via the Born rule. For general OMLs, the extension problem is open.¹

Even when probabilistic descent succeeds, value-definite descent does not. The Bub–Clifton theorem [2] addresses what *can* be salvaged:

Theorem 2.11 (Bub–Clifton [2]). *Given a state ρ on $L(H)$ and a preferred observable R , there is a unique maximal sublattice $D(\rho, R) \subseteq L(H)$ such that:*

- (i) $D(\rho, R)$ is a Boolean algebra;
- (ii) $D(\rho, R)$ contains the spectral projections of R ;
- (iii) The restriction of ρ to $D(\rho, R)$ is a mixture of dispersion-free states.

In the present vocabulary, condition (iii) says that $D(\rho, R)$ is the maximal Boolean subalgebra on which VDR is available — restating Halvorson and Clifton [10] in the EA/PR/VDR framework. It depends on R : different preferred observables yield different maximal Boolean subalgebras.

3 Distributivity as the commensurability condition

The status of the three positions depends on the algebra:

$$\underbrace{\text{EA}}_{\text{always}} \xleftarrow{\sigma\text{-add.}} \underbrace{\text{PR}}_{\text{Gleason}} \xleftarrow{\text{VDR}} \underbrace{\text{dispersion-free}}_{\text{Zorn (Boolean) / KS-blocked (OML)}}$$

The rightmost arrow is free in the Boolean case and obstructed in the OML case. The middle (EA $\leftarrow$ PR) arrow is labelled by σ -additivity only in the Boolean case, where extension to the dual-space measure is free and σ -additivity governs descent alone. In the OML case this arrow also carries the extension step (state $\rightarrow$ measure on $S_0(A)$), which is governed by distributivity, not σ -additivity; for $L(H)$ Gleason supplies both at once.

Theorem 3.1 (Commensurability). *Let A be a complemented lattice.*

¹The extension splits into two axes that the Boolean case fuses. The *extension axis* — passing from the orthogonally additive state on the OML of $\perp$ -stable clopens to a charge on the full Boolean algebra of clopens of $S_0(A)$ — is the classical Boolean extension problem [13]: since the representation map preserves meets, meet-zero elements map to disjoint sets, so the condition is $\sum_i s(a_i) \leq 1$ over pairwise-meet-zero families, a Pitowsky polytope condition [19] strictly stronger than orthoadditivity when the OML is non-distributive. It is not governed by σ -additivity, and in the finite case is a decidable linear program. The *descent axis* — whether the resulting measure concentrates on the physical points — is what σ -additivity governs. For $L(H)$, Gleason resolves both at once.

(a) If A is distributive (Boolean), then EA, PR, and VDR are mutually commensurable:

- EA is unconditional (Stone construction).
- PR holds iff the charges are σ -additive [17].
- VDR holds unconditionally (ultrafilters exist).

Moreover, the choice of realisation space Ω is unconstrained by the algebra.

(b) If A is non-distributive (OML with $A \cong L(H)$, $\dim H \geq 3$), then:

- EA is unconditional.²
- PR holds (Gleason, Theorem 2.8).
- VDR fails globally (Kochen–Specker, Theorem 2.7).
- VDR holds locally within each Boolean context (Bub–Clifton, Theorem 2.11).

Proof. Part (a) is Proposition 1.1. For part (b): PR follows from Gleason’s theorem (Theorem 2.8), which supplies the σ -additive measure on $S_0(L(H))$ for any σ -additive state; EA follows from PR, since $PR \Rightarrow EA$; VDR-failure from Kochen–Specker; local VDR from Bub–Clifton. $\square$

Corollary 3.2. *Distributivity is sufficient for VDR. For the physically central case $A = L(H)$ with $\dim H \geq 3$, non-distributivity blocks VDR.*

Proof. Sufficiency: Theorem 3.1(a). Blockage: Theorem 3.1(b). $\square$

Remark 3.3. The dimensional restriction is necessary. For $\dim H = 2$, $L(\mathbb{C}^2)$ is non-distributive but admits dispersion-free states. More generally, horizontal sums of Boolean algebras and certain finite OMLs are non-distributive yet admit 2-valued homomorphisms [15, Chapter 4]. The sharp characterisation for general OMLs is: VDR is available iff A admits a dispersion-free state, which is a property of A , not of distributivity alone. Distributivity is sufficient (ultrafilters exist) but not necessary in full generality. For the physically and epistemologically central case $\dim H \geq 3$, it is the effective dividing line.

The containment $\mathcal{S}(A) \supseteq \mathcal{S}_\sigma(A) \supseteq \mathcal{S}_{\text{df}}(A)$ is standard in quantum logic [20, 15]. What we add is the identification of three transitions of different mathematical character, and the observation that distributivity controls which are trivial.

Definition 3.4 (Coherence filtration). Let A be a complemented lattice with a family of charges. The *state space* of A admits a descending filtration:

$$\mathcal{S}(A) \supseteq \mathcal{S}_\sigma(A) \supseteq \mathcal{S}_{\text{df}}(A),$$

where:

- $\mathcal{S}(A)$ is the set of states on A (contextual coherence);
- $\mathcal{S}_\sigma(A) \subseteq \mathcal{S}(A)$ is the subset of σ -additive states (probabilistic coherence);
- $\mathcal{S}_{\text{df}}(A) \subseteq \mathcal{S}_\sigma(A)$ is the subset of dispersion-free states (value-definite coherence).

Beneath the filtration sits a local condition: *consistency* requires only that each maximal Boolean subalgebra of A individually admits a compatible state.

²Under the reading of EA as the orthogonally additive set function on the $\perp$ -stable clopens. Promoting this to a genuine Baire measure on $S_0(A)$ is the extension step, which for $L(H)$ is supplied by Gleason and for general OMLs is open. The unconditional claim is thus about the existence of the clopen set function, not about the dual-space measure.

Each inclusion can be strict. The three transitions have different mathematical character:

Transition	Character	Key	Obstruction
$\text{Consistency} \rightarrow \mathcal{S}(A)$	Pasting (topological)	Do local sections glue?	Contextuality
$\mathcal{S}(A) \rightarrow \mathcal{S}_\sigma(A)$	Regularity (analytic)	Does the state have good limits?	σ -additivity
$\mathcal{S}_\sigma(A) \rightarrow \mathcal{S}_{\text{df}}(A)$	Sharpness (algebraic)	Can probabilities collapse to $\{0, 1\}$?	Kochen–Specker

Distributivity collapses the filtration. For a Boolean algebra:

- *Pasting* is trivial: every local assignment extends to a global state (distributivity guarantees compatibility).
- *Sharpness* is free: ultrafilters exist (Zorn’s lemma), so $\mathcal{S}_{\text{df}}(A) \neq \emptyset$ unconditionally.
- The only non-trivial transition is *regularity*: σ -additivity of the charges. This is the content of [17].

For an OML with $\dim \geq 3$, every transition is non-trivial:

- *Pasting* is non-trivial: contextuality obstructions are possible.
- *Regularity* requires Gleason-type results. (This filtration grades *internal* states of A ; the further passage from a σ -additive state to a measure on the dual space $S_0(A)$ — the extension step, free in the Boolean case but blocked by non-distributivity in general — is a separate question, supplied by Gleason for $L(H)$ and open otherwise.)
- *Sharpness* is blocked: $\mathcal{S}_{\text{df}}(A) = \emptyset$ (Kochen–Specker).

Distributivity is sufficient to collapse the filtration. For $L(H)$ with $\dim \geq 3$, non-distributivity opens it fully.

Remark 3.5 (Relation to contextuality hierarchies). Abramsky and Brandenburger [1] introduced a hierarchy of contextuality — probabilistic, possibilistic, and strong — grading the *strength of the obstruction* to a global section of a presheaf of probability distributions. Our filtration is complementary: it grades *what the algebra can support*, not how badly a global section fails. Their axis measures obstruction strength; ours measures coherence demands.

4 Discussion

The standard philosophical debate between scientific realism and constructive empiricism treats the question as uniform: either theories aim at truth or they aim at empirical adequacy. The algebraic analysis above shows that the question is *parametrised* by the observation algebra. For Boolean observations, the mathematics does not prefer one position over another. For non-Boolean observations ($\dim \geq 3$), VDR is algebraically impossible; the constructive empiricist position [21] is the generic one, and value-definite realism is the special case — the case in which the observation algebra happens to be distributive.

This reframing retroactively sharpens classical debates. The EPR argument for completeness [7] amounts to the demand that all observables simultaneously possess definite values — i.e., VDR. The Kochen–Specker theorem shows this is algebraically blocked for $\dim H \geq 3$. Bohr’s complementarity corresponds to the non-triviality of the pasting transition: incompatible contexts cannot be simultaneously realised. Bell’s beables [10] — the observables that *can*

have definite values — are the elements of the maximal Boolean subalgebra $D(\rho, R)$ (Theorem 2.11). De Finetti’s operationalism is EA in the Boolean case; Pitowsky [19] develops the quantum analogue via Dutch book arguments on $L(H)$.

The analysis is not specific to quantum mechanics. Any collection of individually Boolean observation contexts — jointly performable measurements in behavioural science [6], jointly queryable attributes in machine learning [4], simultaneously administrable treatments in experimental design — produces an observation algebra via categorical gluing [9]. If the contexts are mutually compatible, the colimit is Boolean and VDR is available. If not, the colimit is orthomodular and VDR may be blocked. The commensurability theorem thus provides a diagnostic across domains: when a deterministic latent-state model fails, the failure may be algebraic — the observation algebra does not support VDR — rather than a consequence of noise or insufficient data.

Several directions remain open:

1. *The OML extension problem.* Does every state on a general OML extend to a σ -additive measure on $S_0(A)$, as in the Boolean case? Gleason’s theorem answers this for $L(H)$; the general case is open (§2).
2. *Dynamical structure.* If the observation algebra carries a group action (stationarity, exchangeability), how does symmetry interact with the coherence hierarchy? The Boolean case is partially understood (de Finetti’s theorem for exchangeability); the OML case is unexplored.
3. *Causal inference.* Holland’s fundamental problem of causal inference [12] — that both potential outcomes $Y(0)$ and $Y(1)$ cannot be observed for the same unit — is structurally analogous to Bohr complementarity. Whether partial identification constraints generate non-Boolean algebraic structure, and whether the commensurability theorem applies to counterfactual reasoning, is unexplored.
4. *Relation to the topos programme.* The Bohrfication programme [14, 5, 11] encodes related structure more powerfully: the spectral presheaf over the poset of Boolean subalgebras captures VDR-failure as absence of global sections (Kochen–Specker), and local VDR within each Boolean context is built into the presheaf’s local sections. Our filtration operates at a coarser level — distinguishing EA, PR, and VDR as properties of the algebra’s state space rather than as sheaf-cohomological obstructions. The McDonald–Bimbó duality offers a complementary perspective: a single topological dual space (not a presheaf over contexts) in which principal filters are structurally distinguished. Whether the EA/PR/VDR vocabulary translates into the topos framework, and whether the filtration refines or merely simplifies the sheaf-theoretic picture, remains open.

References

- [1] Samson Abramsky and Adam Brandenburger. The sheaf-theoretic structure of non-locality and contextuality. *New Journal of Physics*, 13:113036, 2011.
- [2] Jeffrey Bub and Rob Clifton. A uniqueness theorem for ‘no collapse’ interpretations of quantum mechanics. *Studies in History and Philosophy of Modern Physics*, 27(2):181–219, 1996.
- [3] J. R. Choksi. Inverse limits of measure spaces. *Proceedings of the London Mathematical Society*, s3-8(3):321–342, 1958.
- [4] Rabanus Derr and Robert C. Williamson. Systems of precision: coherent probabilities on pre-Dynkin-systems and coherent previsions on linear subspaces. *arXiv preprint arXiv:2302.03522*, 2023.

- [5] Andreas Döring and Christopher J. Isham. A topos foundation for theories of physics: II. Daseinisation and the liberation of quantum theory. *Journal of Mathematical Physics*, 49(5):053516, 2008.
- [6] Ehtibar N. Dzhafarov and Janne V. Kujala. Contextuality-by-default 2.0: Systems, connections, and measures of contextuality. *Foundations of Physics*, 47:1009–1029, 2017.
- [7] Albert Einstein, Boris Podolsky, and Nathan Rosen. Can quantum-mechanical description of physical reality be considered complete? *Physical Review*, 47:777–780, 1935.
- [8] Andrew M. Gleason. Measures on the closed subspaces of a Hilbert space. *Journal of Mathematics and Mechanics*, 6:885–893, 1957.
- [9] Yukio-Pegio Gunji, Taichi Haruna, and Daisuke Uragami. Contextuality as a left adjoint: a categorical generation of orthomodular structure. *arXiv preprint arXiv:2603.22353*, 2026.
- [10] Hans Halvorson and Rob Clifton. Maximal beable subalgebras of quantum-mechanical observables. *International Journal of Theoretical Physics*, 38:2441–2484, 1999.
- [11] Chris Heunen, Nicolaas P. Landsman, and Bas Spitters. A topos for algebraic quantum theory. *Communications in Mathematical Physics*, 291:63–110, 2009.
- [12] Paul W. Holland. Statistics and causal inference. *Journal of the American Statistical Association*, 81(396):945–960, 1986.
- [13] Alfred Horn and Alfred Tarski. Measures in Boolean algebras. *Transactions of the American Mathematical Society*, 64(3):467–497, 1948.
- [14] Christopher J. Isham and Jeremy Butterfield. Topos perspective on the Kochen–Specker theorem: I. Quantum states as generalised valuations. *International Journal of Theoretical Physics*, 37:2669–2733, 1998.
- [15] Gudrun Kalmbach. *Orthomodular Lattices*. Academic Press, 1983.
- [16] Simon Kochen and Ernst P. Specker. The problem of hidden variables in quantum mechanics. *Journal of Mathematics and Mechanics*, 17:59–87, 1967.
- [17] Patrick S. Mahon. When does observational coherence determine probability? Preprint, 2026.
- [18] Joseph McDonald and Katalin Bimbó. Topological duality for orthomodular lattices. *Mathematical Logic Quarterly*, 69(4):514–541, 2023.
- [19] Itamar Pitowsky. *Quantum Probability — Quantum Logic*, volume 321 of *Lecture Notes in Physics*. Springer, 1989.
- [20] Pavel Pták and Sylvia Pulmannová. *Orthomodular Structures as Quantum Logics*. Kluwer Academic Publishers, 1991.
- [21] Bas C. van Fraassen. *The Scientific Image*. Oxford University Press, 1980.