

# Dynamics and Reconstruction in the Observable Measure

Patrick S. Mahon

## Abstract

Starting from a  $\sigma$ -additive probability measure on an observable algebra [6], this paper derives two structures that are standardly assumed rather than earned: the dynamics of observation under time, and the recoverability of the underlying state space. The Rokhlin disintegration theorem forces a conditional regularity kernel  $\kappa_Q$  into existence once  $\sigma$ -additivity holds; composing with  $Q$  gives the minimal sufficient factor  $Q_*$ , the coarsest reduction carrying full conditional information. Indexing over time, temporal coherence forces the Chapman–Kolmogorov equation — derived, not assumed — yielding a Markov semigroup  $\{\Pi_t\}$  and Koopman–Perron duality. In a measure-preserving system  $Q_*$  becomes the delay map  $\Phi_h$ , and reconstruction — that  $\mathcal{O}_h = \mathcal{B} \bmod \mu$ , density of the delay algebra in  $L^2$ , and injectivity of  $\Phi_h$  are equivalent — is the faithfulness question about this factor. The Stone space [6] built to derive  $\sigma$ -additivity is identified with  $X$  itself when reconstruction holds.

**2020 Mathematics Subject Classification.** 37A30 (primary); 47D06, 60J05, 37A05, 28A60, 06E15, 46E30 (secondary).

**Keywords.** Koopman operator, conditional regularity kernel, minimal sufficient factor, Markov semigroup, Koopman–Perron duality, observable dynamical system, reconstruction theorem, delay embedding, observable algebra, density bridge, measure-theoretic Takens, Stone duality, measure-preserving system.

## 1 Introduction

To study a dynamical system through its observables, Koopman [4] begins from the transformation  $T$  and defines an operator:  $f \mapsto f \circ T$ . To study prediction, the stochastic-process tradition begins from a filtration and builds conditional expectations within it [1]. In both cases the dynamical or temporal structure is written in first, and the probability measure is defined on top of it. This is the natural order.

But the probability measure, once in hand, already contains more than it was given credit for. Any two observations  $Q$  and  $F$  — no hypothesis on time or order — determine a conditional regularity of  $F$  given  $Q$ : the Markov kernel  $\kappa_Q(q, \cdot) = P(F \in \cdot \mid Q = q)$ , which the Rokhlin disintegration theorem [7] forces into existence once  $\sigma$ -additivity holds. It is not added; it is already there. Composing with  $Q$  gives the minimal sufficient factor  $Q_*$ , the coarsest reduction of the sample point that carries full conditional information about  $F$ . Temporal prediction is the special case in which  $F$  is a future observable; nothing in the construction requires it.

Index the conditional regularity over time and the kernels  $\{\Pi_t\}$  must compose consistently. That coherence forces Chapman–Kolmogorov — derived, not assumed — and from it a Markov semigroup and Koopman–Perron duality  $\int K_t g d\mu = \int g d(\mu \star \Pi_t)$ . In a measure-preserving system with  $Q_t = h \circ T^t$ , the factor  $Q_*$  becomes the delay map  $\Phi_h(x) = (h(T^n x))_{n \in \mathbb{Z}}$ , and the question becomes internal: does this factor see all of  $X$ , or only a proper quotient? The answer — that  $\mathcal{O}_h = \mathcal{B} \bmod \mu$ , density of the delay algebra in  $L^2$ , and injectivity of  $\Phi_h$  are all equivalent — is the reconstruction theorem. The Stone space [6] built to derive  $\sigma$ -additivity is identified with  $X$  itself when reconstruction holds.

## 2 Conditional regularity and the canonical factor

### 2.1 The conditional regularity kernel

Fix a probability space  $(\Omega, \mathcal{F}, P)$  and two measurable maps:

- $Q : \Omega \rightarrow \alpha$ , where  $(\alpha, \mathcal{A})$  is a measurable space;
- $F : \Omega \rightarrow \beta$ , where  $(\beta, \mathcal{B})$  is a standard Borel space.

The standard Borel hypothesis on  $\beta$  is used once: it is the hypothesis under which the Rokhlin disintegration theorem [7] guarantees the existence of a regular conditional distribution. All other results hold in greater generality.

We write  $\mathcal{P}(\beta)$  for the space of probability measures on  $\beta$ , equipped with its canonical (Giry)  $\sigma$ -algebra. The joint pushforward  $\rho := P \circ (Q, F)^{-1}$  on  $\alpha \times \beta$  has marginals  $\rho_\alpha = P \circ Q^{-1}$  and  $\rho_\beta = P \circ F^{-1}$ .

The regular conditional distribution of  $F$  given  $Q$  is the unique (up to  $P \circ Q^{-1}$ -null sets) Markov kernel  $kappa_Q : \alpha \rightarrow \mathcal{P}(\beta)$  satisfying

$$\int_A kappa_Q(q, B) d(P \circ Q^{-1})(q) = P(Q \in A, F \in B)$$

for all measurable  $A \subseteq \alpha$  and  $B \subseteq \beta$ .

**Definition 2.1** (Conditional regularity kernel). The *conditional regularity kernel* of  $F$  given  $Q$  is

$kappa_Q$ , the disintegration kernel of  $\rho = P \circ (Q, F)^{-1}$  over its first marginal:

$kappa_Q(q, A) = P(F \in A \mid Q = q)$ . When  $F$  is a future observable the kernel describes conditional prediction; the object itself is neutral on time.

The kernel is covariant under measurable maps: if  $\pi : \alpha \rightarrow \alpha'$  is measurable and  $Q' = \pi \circ Q$ , then

$kappa_Q(q, \cdot) = \kappa_{Q'}(\pi(q), \cdot)$  for  $(P \circ Q^{-1})$ -almost every  $q$ , by uniqueness of disintegration.

Composing

$kappa_Q$  with  $Q$  gives the map sending each sample point to its conditional regularity.

### 2.2 The minimal sufficient factor

**Definition 2.2** (Minimal sufficient factor). The *minimal sufficient factor* is

$$Q_* : \Omega \rightarrow \mathcal{P}(\beta), \quad Q_*(\omega) = kappa_Q(Q(\omega), \cdot).$$

Since

$kappa_Q$  is a Markov kernel it is measurable, and  $Q_*$  inherits measurability by composition.

**Theorem 2.3** (Factorization via the canonical factor). For every bounded measurable  $g : \beta \rightarrow \mathbb{R}$ :

$$E[g(F) \mid \sigma(Q)] = \int g dQ_*(\cdot) \quad P\text{-a.e.}$$

*Proof.* Let  $h(\omega) := \int g dQ_*(\omega)$ . We verify the two defining properties.

*Measurability.* The map  $q \mapsto \int g d$

$kappa_Q(q, \cdot)$  is strongly measurable; since  $Q_* =$

$kappa_Q(Q(\cdot), \cdot)$ , the map  $h$  is  $\sigma(Q)$ -measurable.

*Set-integral identity.* For  $S = Q^{-1}(A)$ :

$$\begin{aligned} \int_S h dP &= \int_A \int g d \\ \kappa_Q(q, \cdot) d(P \circ Q^{-1})(q) &\quad (\text{pushforward}) \\ &= \int_{A \times \beta} g(f) d\rho(q, f) \quad (\text{disintegration}) \\ &= \int_S g(F) dP. \end{aligned}$$

The conclusion follows from a.e. uniqueness of conditional expectation.  $\square$

Since  $\sigma(Q_*) \subseteq \sigma(Q)$  and the right-hand side is already  $\sigma(Q_*)$ -measurable,  $Q_*$  is a sufficient statistic for  $F$  given  $Q$ :  $E[g(F) \mid \sigma(Q)] = E[g(F) \mid \sigma(Q_*)]$   $P$ -a.e. It is *minimal* in the sense that every sufficient statistic factors through it:  $Q_*$  is the coarsest reduction of  $\omega$  that carries the full conditional regularity of  $F$  given  $Q$ . The operator  $(K_{Q_*}g)(q_*) = \int g dq_*$  on  $\mathcal{P}(\beta)$  recasts this as  $E[g(F) \mid \sigma(Q)] = (K_{Q_*}g) \circ Q_*$   $P$ -a.e.

### 3 Dynamics and duality

#### 3.1 Markov semigroup

When the conditional regularity kernel

$\kappa_Q$  is indexed over a time parameter, the kernels cannot be chosen independently: the regularity of  $F$  at time  $t + s$  given  $Q$  at time 0 must be consistent with first observing at time  $s$  and then at time  $t$ . This consistency requirement forces the Chapman–Kolmogorov equation.

Concretely, index over a discrete time horizon  $\mathbb{N}$  and let  $\Pi_t : \gamma \rightarrow \mathcal{P}(\gamma)$  be a Markov kernel on a state space  $\gamma$ , with  $\Pi_0(q, \cdot) = \delta_q$ . Temporal coherence of the conditional regularity demands

$$\Pi_{t+s}(q, \cdot) = \int_{\gamma} \Pi_t(q', \cdot) d\Pi_s(q, q') \quad (\text{i.e., } \Pi_{t+s} = \Pi_t \circ_{\kappa} \Pi_s)$$

— the Chapman–Kolmogorov equation, derived from the structure rather than assumed. A family  $\{\Pi_t\}$  satisfying it is a *Markov semigroup kernel*. The associated *Markov operators* are

$$(K_t g)(q) := \int g d\Pi_t(q, \cdot), \quad g : \gamma \rightarrow \mathbb{R}.$$

Each  $K_t$  is positive, unital, and contractive on bounded measurable functions, with  $K_0 = \text{id}$ ; positivity, unitality, and contractiveness are immediate from the Markov property.

**Proposition 3.1** (Semigroup property).  $K_{t+s}g = K_t(K_s g)$  for all bounded measurable  $g$ .

*Proof.*  $(K_{t+s}g)(q) = \int g d(\Pi_t \circ_{\kappa} \Pi_s)(q, \cdot) = \int \left( \int g d\Pi_t(q', \cdot) \right) d\Pi_s(q, q') = \int (K_t g)(q') d\Pi_s(q, q') = (K_t(K_s g))(q)$ , where the exchange of integrals is justified by Bochner–Fubini.  $\square$

The pushforward of a measure  $\mu$  on  $\gamma$  through the kernel is  $P_t^* \mu := \mu \star \Pi_t = \int \Pi_t(q, \cdot) d\mu(q)$ .

#### 3.2 Koopman–Perron duality

**Theorem 3.2** (Koopman–Perron duality). For every bounded measurable  $g$  and finite measure  $\mu$ :

$$\int (K_t g) d\mu = \int g d(P_t^* \mu).$$

*Proof.*  $\int g d(P_t^* \mu) = \int g d(\mu \star \Pi_t) = \int_{\gamma} \int g d\Pi_t(q, \cdot) d\mu(q) = \int (K_t g) d\mu$ , by Bochner–Fubini.  $\square$

When  $\Pi_t(q, \cdot) = \delta_{\varphi_t(q)}$  for a measurable map  $\varphi_t : \gamma \rightarrow \gamma$ , the stochastic picture collapses.

### 3.3 Specialisation to measure-preserving systems

**Proposition 3.3** (Deterministic limit). *If  $\Pi_t(q, \cdot) = \delta_{\varphi_t(q)}$ , then  $K_t g = g \circ \varphi_t$  and  $\varphi_{t+s} = \varphi_t \circ \varphi_s$ .*

*Proof.*  $(K_t g)(q) = \int g d\delta_{\varphi_t(q)} = g(\varphi_t(q))$ . The flow law follows from  $\delta_{\varphi_{t+s}(q)} = (\Pi_t \circ \kappa \Pi_s)(q, \cdot) = \delta_{\varphi_t(\varphi_s(q))}$  and injectivity of the Dirac embedding.  $\square$

In the deterministic case,  $K_t g = g \circ \varphi_t = U_T^t g$ : the Markov operator specialises to the classical Koopman operator. The Koopman–Perron duality  $\int K_t g d\mu = \int g d(P_t^* \mu)$  becomes the change-of-variables formula  $\int g \circ T^t d\mu = \int g d(T_*^t \mu)$ .

**Example 3.4** (Finite Markov chain). Let  $X = \{1, \dots, k\}$ ,  $\mathcal{B} = 2^X$ , and let  $P = (p_{ij})$  be the transition matrix of a finite irreducible Markov chain with unique stationary distribution  $\mu$  ( $\mu P = \mu$ ). Take full-state observation  $Q_t = X_t$ . The conditional regularity kernel of  $X_1$  given  $X_0$  is  $\kappa(i, \cdot) = P(i, \cdot)$ , the  $i$ -th row of  $P$ : this is the unique Markov kernel satisfying  $\mathbb{E}_\mu[f(X_1) \mid X_0 = i] = \sum_j p_{ij} f(j)$  for all bounded  $f$ . The  $t$ -step kernel  $\Pi_t(i, \cdot) = P^t(i, \cdot)$  is the corresponding semigroup kernel, and temporal coherence gives  $\Pi_{t+s} = \Pi_t \circ \kappa \Pi_s$ , i.e.  $P^{t+s} = P^t P^s$ . The semigroup acts as

$$K_t f(i) = \sum_j p_{ij}^t f(j).$$

Koopman–Perron duality reduces to the stationary-measure identity:

$$\int K_t g d\mu = \mu^\top P^t g = (\mu P^t)^\top g = \mu^\top g = \int g d\mu,$$

the fixed-point property  $\mu P = \mu$  of the stationary distribution. In this finite setting every charge is automatically  $\sigma$ -additive, so the admissibility condition of [6] is trivially satisfied. Reconstruction holds at lag 0:  $\mathcal{O}_h = \sigma(h) = \mathcal{B}(X)$  whenever  $h$  separates points of  $X$ .

The full structure produced by the construction is the triple  $(\mathcal{P}(\beta), \nu_{Q_*}, \{K_t\}_{t \in \mathbb{N}})$ : the canonical factor space, stationary distribution  $\nu_{Q_*} = P \circ Q_*^{-1}$ , and Markov semigroup. This is the *observable dynamical system* derived from  $(Q, F, P)$ : the conditional regularity of  $F$  relative to  $Q$ , promoted to a dynamical object by temporal coherence.

## 4 From regularity to reconstruction

Setting  $Q_t = h \circ T^t$  in a measure-preserving system  $(X, \mathcal{B}, \mu, T)$  with  $h \in L^\infty$  specialises the construction. Under this choice,  $Q_*(\omega) = (h(T^n \omega))_{n \in \mathbb{Z}}$  is the delay map  $\Phi_h(x) = (h(T^n x))_{n \in \mathbb{Z}}$ , and the observable dynamical system is the triple  $(\Phi_h(X), \mu \circ \Phi_h^{-1}, \{K_t\})$ .

The canonical factor space  $\Phi_h(X)$  is a subset of  $\mathbb{R}^\mathbb{Z}$  encoding the dynamical information that  $h$  can see. Whether it is a faithful image of  $X$  depends on how much the delay orbit  $\{h \circ T^n : n \in \mathbb{Z}\}$  collectively distinguishes points of  $X$ . Injectivity of  $\Phi_h$  modulo  $\mu$ -null sets is equivalent to the observable  $\sigma$ -algebra

$$\mathcal{O}_h = \sigma(\{h \circ T^n : n \in \mathbb{Z}\})$$

equalling  $\mathcal{B}(X)$  modulo  $\mu$ .

The Koopman–Perron duality (Theorem 3.2) connects this to a function-space question: the observable  $\sigma$ -algebra  $\mathcal{O}_h$  determines the Markov semigroup through the algebra it generates, and the reconstruction question is precisely whether that algebra is large enough to recover all of  $\mathcal{B}$ .

## 4.1 Separation defect and the role of CE

For a finite joined query algebra  $\mathcal{G} = \mathcal{E}_{n_1} \vee \dots \vee \mathcal{E}_{n_k}$  from a directed system  $(\mathcal{E}_n)_n$ , the *separation defect* is

$$\delta(\mathcal{G}) := \sup_{A \in \sigma(\mathcal{C})} \inf_{E \in \mathcal{G}} \mu(A \triangle E),$$

where  $\sigma(\mathcal{C})$  is the observable  $\sigma$ -algebra of the full system. It vanishes iff  $\sigma(\mathcal{G}) = \sigma(\mathcal{C}) \bmod \mu$ . A sequence  $(\mathcal{G}_k)$  is *honest* if it is increasing and  $\sigma(\bigcup_k \mathcal{G}_k) = \sigma(\mathcal{C}) \bmod \mu$ ; under an honest scheme  $\delta(\mathcal{G}_k)$  is non-increasing.

**Theorem 4.1** (CE drives separation defect to zero). *Let  $(\mathcal{E}_n)_n$  be a query system with realization measure  $\mu$  satisfying collective exhaustion [6]. Under any honest refinement scheme  $(\mathcal{G}_k)$ ,*

$$\delta(\mathcal{G}_k) \longrightarrow 0 \quad \text{as } k \rightarrow \infty.$$

*Proof.* Fix  $A \in \sigma(\mathcal{C})$ . Since  $(\mathcal{G}_k)$  is honest,  $\sigma(\bigcup_k \mathcal{G}_k) = \sigma(\mathcal{C}) \bmod \mu$ , so  $A$  is  $\sigma(\bigcup_k \mathcal{G}_k)$ -measurable. By the  $L^1(\mu)$  martingale convergence theorem,  $\mathbf{1}_{E_k} \rightarrow \mathbf{1}_A$  in  $L^1(\mu)$  where  $E_k = \mathbb{E}_\mu[\mathbf{1}_A \mid \mathcal{G}_k]$  rounded to the nearest atom; in particular  $\mu(A \triangle E_k) \rightarrow 0$ . CE [6] ensures no mass persists on the vanishing sets  $A \triangle E_k$ , confirming  $\inf_{E \in \mathcal{G}_k} \mu(A \triangle E) \rightarrow 0$  uniformly over  $A$ . Hence  $\delta(\mathcal{G}_k) \rightarrow 0$ .  $\square$

Sections 5–6 carry out the reconstruction in the delay-map setting.

**Remark 4.2** (The temporal specialisation). With  $Q_t = h \circ T^t$ , the algebra  $\mathcal{O}_h^{(L)}$  is an honest refinement scheme once reconstruction holds (§§5–6). The separation defect specialises to  $\delta(L) := \delta(\mathcal{O}_h^{(L)})$ , the central quantity of [5]: exponential decay of  $\delta(L)$  under mixing yields minimax-optimal certification rates. In the general query setting, exhaustion and mixing are separate conditions; the delay-map setting collapses the first by construction.

## 5 Density bridge and reconstruction

### 5.1 Density bridge lemma

Throughout this section and the next,  $(X, \mathcal{B}, \mu)$  is a standard probability space,  $T : X \rightarrow X$  is a measurable,  $\mu$ -preserving invertible bimeasurable bijection with  $T_*\mu = \mu$ , and  $h \in L^\infty(X, \mu)$ .

**Definition 5.1** (Observable algebra). The *observable algebra* generated by  $h$  is

$$\mathcal{O}_h := \sigma(\{h \circ T^n : n \in \mathbb{Z}\}) \subseteq \mathcal{B}(X).$$

The *delay algebra*  $\mathcal{A}_h$  is the smallest subalgebra of  $L^\infty(X, \mu)$  containing 1 and every  $h \circ T^n$ ; concretely, finite sums  $\sum_\alpha c_\alpha \prod_k (h \circ T^{n_k})^{e_k}$ .

**Definition 5.2** (Delay map). The *delay map* is

$$\Phi_h : X \rightarrow \mathbb{R}^{\mathbb{Z}}, \quad \Phi_h(x) = (h(T^n x))_{n \in \mathbb{Z}}.$$

Since  $\pi_n \circ \Phi_h = h \circ T^n$  for each coordinate projection  $\pi_n$ , the map  $\Phi_h$  is measurable and  $\mathcal{O}_h = \Phi_h^{-1}(\mathcal{B}(\mathbb{R}^{\mathbb{Z}}))$ . Thus  $\mathcal{O}_h = \mathcal{B} \bmod \mu$  is precisely the condition that  $\Phi_h$  is a measure-theoretic embedding.

The central lemma connects the Boolean algebra level to the  $L^2$  level. It holds for any subalgebra of  $L^\infty$ .

**Lemma 5.3** (Density bridge). *Let  $(X, \mathcal{B}, \mu)$  be a probability space and let  $\mathcal{A} \subseteq L^\infty(X, \mu)$  be a subalgebra containing the constant function 1. Then:*

$$\overline{\mathcal{A}}^{L^2} = L^2(X, \sigma(\mathcal{A}), \mu).$$

*Consequently,  $\overline{\mathcal{A}}^{L^2} = L^2(X, \mu)$  if and only if  $\sigma(\mathcal{A}) = \mathcal{B}$  modulo  $\mu$ .*

*Proof.* Write  $\mathcal{M} := \sigma(\mathcal{A})$  and  $V := \overline{\mathcal{A}}^{L^2}$ .

**Step 1:**  $V \subseteq L^2(X, \mathcal{M}, \mu)$ . Every  $f \in \mathcal{A}$  is  $\mathcal{A}$ -measurable, hence  $\mathcal{M}$ -measurable. So  $\mathcal{A} \subseteq L^2(X, \mathcal{M}, \mu)$ . Since  $L^2(X, \mathcal{M}, \mu)$  is a closed subspace of  $L^2(X, \mathcal{B}, \mu)$ , taking the closure gives  $V \subseteq L^2(X, \mathcal{M}, \mu)$ .

**Step 2:**  $L^2(X, \mathcal{M}, \mu) \subseteq V$ . It suffices to show that every indicator  $\mathbf{1}_E$  with  $E \in \mathcal{M}$  lies in  $V$ . By the functional monotone class theorem [2], the class

$$\mathcal{H} := \{f \in L^\infty(X, \mathcal{M}, \mu) : f \in V\}$$

is a vector space containing  $\mathcal{A}$  and closed under bounded monotone pointwise limits (since if  $f_n \in V$  are uniformly bounded and  $f_n \rightarrow f$  pointwise a.e. with  $f \in L^\infty$ , then  $f_n \rightarrow f$  in  $L^2$  by dominated convergence, so  $f \in V$ ). Since  $\mathcal{A}$  is an algebra containing 1 and  $\sigma(\mathcal{A}) = \mathcal{M}$ , the monotone class theorem gives  $\mathcal{H} = L^\infty(X, \mathcal{M}, \mu)$ . In particular every  $\mathcal{M}$ -indicator  $\mathbf{1}_E \in V$ .

Since the  $\mathcal{M}$ -simple functions are dense in  $L^2(X, \mathcal{M}, \mu)$  and  $V$  is closed,  $L^2(X, \mathcal{M}, \mu) \subseteq V$ .

**Conclusion.** Steps 1 and 2 give  $V = L^2(X, \mathcal{M}, \mu)$ . The final equivalence is immediate:  $V = L^2(X, \mathcal{B}, \mu)$  iff  $L^2(X, \mathcal{M}, \mu) = L^2(X, \mathcal{B}, \mu)$  iff  $\mathcal{M} = \mathcal{B}$  mod  $\mu$ .  $\square$

**Remark 5.4** (Role of the algebra structure). The algebra structure of  $\mathcal{A}$  is essential. The corresponding statement for the closed linear span  $\mathcal{H}_{\mathcal{A}} = \overline{\text{span}}(\mathcal{A})$  is false in general:  $\mathcal{H}_{\mathcal{A}} = L^2$  does not imply  $\sigma(\mathcal{A}) = \mathcal{B}$ .

For example, take  $X = [0, 1]$ ,  $\mu = \text{Lebesgue}$ ,  $T(x) = 2x \bmod 1$  (the doubling map), and  $h(x) = x$ . The orbit  $\{h \circ T^n\}(x) = \{2^n x \bmod 1\}$  generates  $\mathcal{B}([0, 1])$  as a  $\sigma$ -algebra (the dyadic structure separates points). But  $\text{span}\{2^n x \bmod 1 : n \geq 0\}$  is a proper subspace of  $L^2[0, 1]$ : it does not contain  $x^2$ . So the linear span can be a proper dense subspace even when  $\sigma(\mathcal{A}) = \mathcal{B}$ .

In the other direction,  $h$  cyclic for  $U_T$  (linear span dense) always implies  $\sigma(\mathcal{A}_h) = \mathcal{B}$  mod  $\mu$  — see Corollary 6.2.

## 5.2 Reconstruction theorem

**Theorem 5.5** (Reconstruction theorem). *Let  $(X, \mathcal{B}, \mu, T)$  be an invertible measure-preserving system and  $h \in L^\infty(X, \mu)$ . The following are equivalent:*

- (i)  $\mathcal{O}_h = \mathcal{B}$  modulo  $\mu$ ;
- (ii)  $\mathcal{A}_h$  is dense in  $L^2(X, \mu)$ ;
- (iii)  $\Phi_h : X \rightarrow \mathbb{R}^\mathbb{Z}$  is a measure-theoretic embedding:  $\Phi_h^{-1}(\mathcal{B}(\mathbb{R}^\mathbb{Z})) = \mathcal{B}$  modulo  $\mu$ .

*Proof.* (i)  $\Leftrightarrow$  (ii). Apply the density bridge (Lemma 5.3) with  $\mathcal{A} = \mathcal{A}_h$  and  $\sigma(\mathcal{A}_h) = \mathcal{O}_h$ :  $\overline{\mathcal{A}_h}^{L^2} = L^2(X, \mu)$  iff  $\mathcal{O}_h = \mathcal{B}$  mod  $\mu$ .

(i)  $\Leftrightarrow$  (iii). By Definition 5.2,  $\mathcal{O}_h = \Phi_h^{-1}(\mathcal{B}(\mathbb{R}^\mathbb{Z}))$ . So  $\mathcal{O}_h = \mathcal{B}$  mod  $\mu$  iff  $\Phi_h^{-1}(\mathcal{B}(\mathbb{R}^\mathbb{Z})) = \mathcal{B}$  mod  $\mu$ , which is exactly condition (iii).  $\square$

Condition (iii) means  $\Phi_h$  induces an isomorphism of measure algebras  $(X/\mu, \mathcal{B}/\mu) \cong (\mathbb{R}^\mathbb{Z}/(\Phi_h)_*\mu, \mathcal{B}(\mathbb{R}^\mathbb{Z})/(\Phi_h)_*\mu)$ . Since  $(\Phi_h(Tx))_n = h(T^{n+1}x) = (\sigma(\Phi_h(x)))_n$ , the measure  $(\Phi_h)_*\mu$  is shift-invariant, and  $(X, T, \mu)$  is isomorphic to  $(\mathbb{R}^\mathbb{Z}, \sigma, (\Phi_h)_*\mu)$  restricted to  $\Phi_h(X)$ .

## 6 Cyclic vectors and Stone space identification

### 6.1 Cyclic vectors

**Definition 6.1** (Cyclic vector). The function  $h \in L^2(X, \mu)$  is a *cyclic vector* for  $U_T$  if

$$\mathcal{H}_h := \overline{\text{span}}\{U_T^n h : n \in \mathbb{Z}\} = L^2(X, \mu).$$

**Corollary 6.2** (Cyclic implies reconstruction). *If  $h \in L^\infty(X, \mu)$  is a cyclic vector for  $U_T$ , then  $\mathcal{O}_h = \mathcal{B}$  modulo  $\mu$ .*

*Proof.* Since  $\mathcal{A}_h$  contains  $\text{span}\{h \circ T^n\} = \text{span}\{U_T^n h\}$  as a subset,  $\overline{\mathcal{A}_h}^{L^2} \supseteq \mathcal{H}_h = L^2$ . By the density bridge,  $\mathcal{O}_h = \mathcal{B} \bmod \mu$ .  $\square$

**Remark 6.3** (The implication is strict). The converse of Corollary 6.2 fails in general:  $\mathcal{O}_h = \mathcal{B}$  does not imply  $h$  is cyclic. The doubling map example of Remark 5.4 witnesses this.

The cyclic vector condition asks for density of the *linear span* of the orbit; reconstruction asks only for density of the *algebra*. For a bounded  $h$ , the linear span is contained in the algebra, so cyclicity is the stronger condition.

The two conditions coincide precisely when  $U_T$  has *simple  $L^2$ -spectrum*:  $L^2(X, \mu)$  is a cyclic  $U_T$ -module, meaning there exists some cyclic vector. In this case, every generating observation  $h$  (satisfying  $\mathcal{O}_h = \mathcal{B}$ ) is automatically cyclic. Simple spectrum is a generic property of ergodic transformations in the Baire category sense [3] but is not universal.

### 6.2 Stone space identification

The Stone space  $\text{St}(\mathcal{O}_h)$  of the observable algebra is the compact Hausdorff space whose points are ultrafilters on  $\mathcal{O}_h$ , with the natural Stone embedding  $\iota : X \rightarrow \text{St}(\mathcal{O}_h)$  sending  $x$  to the principal ultrafilter  $\{E \in \mathcal{O}_h : x \in E\}$ .

**Theorem 6.4** (Stone space identification). *Suppose  $\mathcal{O}_h = \mathcal{B}$  modulo  $\mu$ . Then  $\iota$  is a measure-theoretic isomorphism:  $\iota$  identifies the measure algebra  $(X/\mu, \mathcal{B}/\mu)$  with  $(\text{St}(\mathcal{O}_h)/\iota_*\mu, \mathcal{B}(\text{St}(\mathcal{O}_h))/\iota_*\mu)$ .*

*Proof.* *Injectivity mod  $\mu$ .* Suppose  $\iota(x) = \iota(x')$ . For any  $E \in \mathcal{B}$  with  $x \in E \not\equiv x'$ , find  $F \in \mathcal{O}_h$  with  $\mu(E \triangle F) = 0$ . Since  $x \in F$  and  $\iota(x) = \iota(x')$ , also  $x' \in F$ , so  $x' \in (E \triangle F) \cup F^c$ , placing  $x'$  in a null set. Hence  $\iota$  is injective  $\mu$ -a.e.

*$\sigma$ -algebra identification.* The pullback  $\iota^{-1}$  maps the Baire  $\sigma$ -algebra of  $\text{St}(\mathcal{O}_h)$  (generated by the clopen sets  $[E] = \{u : E \in u\}$ ) to  $\mathcal{O}_h = \mathcal{B} \bmod \mu$ .

*Measure recovery.* Since  $\mu$  is  $\sigma$ -additive on  $\mathcal{O}_h = \mathcal{B}$ , the collective exhaustion condition of [6] forces  $\iota_*\mu$  to concentrate on the principal ultrafilters  $\iota(X)$ , so  $\iota_*\mu$  is supported on the image and the measure algebras are identified.  $\square$

The Stone space [6] — built as a compact ambient space for deriving  $\sigma$ -additivity — is, under the reconstruction condition, the state space  $X$  itself. The construction that began by seeking a compact completion of the sample space ends by recognising it as the object the observer was trying to recover.

**Example 6.5** (Cantor set: reconstruction without smoothness). Let  $\mathcal{C} \subset [0, 1]$  be the middle-thirds Cantor set, and let  $\mu_{\mathcal{C}}$  be the Cantor measure — the pushforward of the fair-coin product measure  $\nu = (\frac{1}{2}\delta_0 + \frac{1}{2}\delta_2)^{\mathbb{N}}$  under the ternary coding map  $\pi(d_1, d_2, \dots) = \sum_{k \geq 1} d_k/3^k$ . Let  $S : \mathcal{C} \rightarrow \mathcal{C}$  be the shift map

$$S(x) = \begin{cases} 3x & x \in \mathcal{C} \cap [0, \frac{1}{3}], \\ 3x - 2 & x \in \mathcal{C} \cap [\frac{2}{3}, 1], \end{cases}$$

and let  $h(x) = x$ . The system  $(\mathcal{C}, \mu_{\mathcal{C}}, S)$  is measurably conjugate to the Bernoulli shift  $(\{0, 2\}^{\mathbb{N}}, \nu, \sigma)$  via  $\pi$ ; in particular  $S$  is  $\mu_{\mathcal{C}}$ -invariant and ergodic.

For  $x = \pi(d_1, d_2, \dots)$  we have  $S^n x = \pi(d_{n+1}, d_{n+2}, \dots)$ . Since every point of  $\mathcal{C}$  lies in  $[0, \frac{1}{3}] \cup [\frac{2}{3}, 1]$ ,

$$d_{n+1} = 0 \iff S^n x \in [0, \frac{1}{3}], \quad d_{n+1} = 2 \iff S^n x \in [\frac{2}{3}, 1],$$

so  $h \circ S^n$  recovers the  $(n+1)$ -st ternary digit of  $x$ . The delay vector  $\Phi_h^{(L)}(x) = (x, Sx, \dots, S^L x)$  reads off digits  $d_1, \dots, d_{L+1}$ ; its fibres are the generation- $(L+1)$  cylinder sets

$$F_z = \{x \in \mathcal{C} : d_k(x) = z_k, 1 \leq k \leq L+1\}, \quad z \in \{0, 2\}^{L+1},$$

each carrying  $\mu_{\mathcal{C}}(F_z) = 2^{-(L+1)}$ , so  $\mathcal{O}_h^{(L)} = \sigma(\Phi_h^{(L)}) = \sigma\{F_z : z \in \{0, 2\}^{L+1}\}$ . Since  $\mu_{\mathcal{C}}$ -a.e. pair of distinct points differs at some finite digit, these algebras separate points as  $L \rightarrow \infty$ :

$$\mathcal{O}_h = \sigma\left(\bigcup_{L \geq 0} \mathcal{O}_h^{(L)}\right) = \mathcal{B}(\mathcal{C}) \mod \mu_{\mathcal{C}}.$$

By Theorem 5.5,  $\Phi_h$  is a measure-theoretic embedding: the finite-lag observable algebras refine to the full Borel structure as  $L \rightarrow \infty$ .

$\mathcal{C}$  is not a smooth manifold: it is compact and zero-dimensional, with Hausdorff dimension  $\log 2 / \log 3 \approx 0.63$ . The differential-geometric hypotheses of Takens's theorem —  $C^2$  ambient dynamics, integer manifold dimension  $d$ , embedding bound  $2d+1$  — are unavailable before one even asks about regularity of  $S$ . Takens has no domain of application here. Theorem 5.5 applies because it requires only that the observable algebra be measurably generated; smoothness plays no role.

The classical Takens embedding theorem [8] reaches a related conclusion by a different route: for a  $C^2$  diffeomorphism on a compact  $d$ -manifold and a generic smooth  $h$ , the finite-delay map  $\Phi_h^{(k)}(x) = (h(x), h(Tx), \dots, h(T^{k-1}x))$  is a diffeomorphic embedding into  $\mathbb{R}^{2d+1}$  for  $k \geq 2d+1$ . The reconstruction theorem here requires only measurability, uses all delays simultaneously, and replaces the dimension count with the algebraic density condition; in exchange, the embedding is measure-theoretic rather than topological. On a compact manifold with generic smooth  $h$ , the orbit  $\{h \circ T^n\}$  separates points, so Stone–Weierstrass gives  $\mathcal{A}_h$  uniformly dense in  $C(M)$  and hence  $L^2$ -dense: the density condition holds and the two results are compatible.

Reconstruction is an exact asymptotic condition: it asks whether the delay algebra generates all of  $\mathcal{B}$  in the limit of all lags, and says nothing about rates, finite approximations, or what a finite time series can certify. Those questions are taken up in [5].

## References

- [1] Joseph L. Doob. *Stochastic Processes*. Wiley, New York, 1953.
- [2] Eugene B. Dynkin. *Markov Processes*. Springer, 1965.
- [3] Paul R. Halmos. In general a measure preserving transformation is mixing. *Annals of Mathematics*, 45(4):786–792, 1944.
- [4] Bernard O. Koopman. Hamiltonian systems and transformation in Hilbert space. *Proceedings of the National Academy of Sciences*, 17(5):315–318, 1931.
- [5] Patrick S. Mahon. Certifying reconstruction from finite data. Preprint, 2026.
- [6] Patrick S. Mahon. Probability from observation: Carathéodory and Stone routes to the observable extension theorem. Preprint, 2026.

- [7] Vladimir Abramovich Rokhlin. On the fundamental ideas of measure theory. *American Mathematical Society Translations*, 10:1–54, 1952.
- [8] Floris Takens. Detecting strange attractors in turbulence. In *Dynamical Systems and Turbulence, Warwick 1980*, volume 898 of *Lecture Notes in Mathematics*, pages 366–381. Springer, 1981.