

# Probability from Observation: Carathéodory and Stone Routes to the Observable Extension Theorem

Patrick S. Mahon

## Abstract

We prove that a coherent family of observations determines a unique probability measure on the observable  $\sigma$ -algebra. The setting is a directed system of measurable outcome spaces connected by surjective refinement maps, together with a compatible family of finitely additive charges on the resulting cylinder algebra. We identify the necessary and sufficient condition for the family to extend to a  $\sigma$ -additive global measure: *collective exhaustion* (CE), the condition that no level can assign persistent mass to events the system as a whole sees as empty.

The argument proceeds by two independent routes, each illuminating a different aspect of the extension problem. The *Carathéodory route* assumes  $\sigma$ -additive marginals and constructs the global measure by a direct Carathéodory argument on the realization space, with no topological hypotheses on the outcome spaces. The *Stone route* assumes only finite additivity and derives  $\sigma$ -additivity from the compactness of the Stone space: the directed system of Boolean algebras dualises to an inverse system of compact Hausdorff spaces, whose inverse limit carries a canonical  $\sigma$ -additive measure; CE then ensures this measure concentrates on the image of the sample space inside its Stone compactification. Under shared hypotheses the two measures coincide.

A foundational analysis shows that CE is *irreducible*: it is not derivable from any finitary or structural condition on the query system. The finite-cofinite counterexample and a model-theoretic argument via Łoś's theorem locate the boundary precisely. CE is not an assumption imposed from outside but the exact condition separating directed systems in which every convergence is witnessed somewhere in the refinement order from those in which it is not.

Beyond the extension theorem, the Stone space  $\text{St}(\mathcal{C})$  constructed here as a technical device has further significance: it is the canonical compact Hausdorff completion of the sample space for the observable topology. When an observation function is a cyclic vector for the Koopman operator of an underlying dynamical system, this Stone space is measure-theoretically isomorphic to the state space. The compact space introduced here as a technical device opens the possibility that the Stone space of the observable algebra is the state space itself — a connection made precise under reconstruction conditions developed elsewhere.

**2020 Mathematics Subject Classification.** 60A10 (primary); 28A60, 06E15, 28A05 (secondary).

**Keywords.** observable extension theorem, collective exhaustion, Carathéodory extension, Stone duality, Boolean algebra, finitely additive measure,  $\sigma$ -additivity, directed system, query system.

## 1 Introduction

The programme of which this paper is a part does not begin with a probability space. It begins with *distinctions*: the acts of observational separation by which an observer organises their experience of a system. The state space, the  $\sigma$ -algebra, and the measure are not primitives — they are what coherent distinctions, pursued to their conclusion, are forced to produce. This paper proves the first step: a coherent family of observational distinctions determines a unique probability measure. The further steps — that this measure carries canonical dynamics, that

the delay structure of observation (re)constructs the state space, and that finite data can certify (re)construction — are carried out in the companion papers.

Many approaches to probability begin by postulating an underlying probability space and then studying the observables derived from it. This paper reverses that perspective. Starting from observable experiments organised as a refinement system of queries, we ask: under what conditions do compatible observable laws determine a canonical probability measure?

The answer is given by a single condition. A compatible family of finitely additive charges on a directed system of Boolean algebras extends to a  $\sigma$ -additive measure at every level if and only if the family is *collectively exhaustive* (CE): for every decreasing sequence of events whose intersection is empty, some level in the refinement order witnesses the vanishing of mass. CE is not a structural condition on the query system but a condition on the charges: it rules out purely finitely additive charges, which satisfy every finite consistency condition while assigning persistent mass to sequences that vanish in the limit.

**Two routes.** The extension theorem is established by two independent proofs, each illuminating CE from a different direction.

The *Carathéodory route* (§4) assumes  $\sigma$ -additive marginals and constructs the global measure directly by Carathéodory extension on the realization space. No topology is required on the outcome spaces. Sequential common refinements compress any countable cylinder family to a single query level, reducing  $\sigma$ -subadditivity of the cylinder premeasure to  $\sigma$ -subadditivity of a single marginal. In this route,  $\sigma$ -additivity of the marginals is the standing hypothesis; by the CE theorem of §3, it is equivalent to CE.

The *Stone route* (§5) assumes only finite additivity and derives  $\sigma$ -additivity from compactness. Stone duality converts the directed system of Boolean algebras into a cofiltered inverse system of compact Hausdorff spaces. A compatible family of finitely additive charges assembles into a  $\sigma$ -additive measure on the inverse limit without any countable additivity assumption. The remaining task is to descend from the Stone space to the sample space. Two conditions are required. Discriminability (§2) ensures the Stone embedding is injective, so the pullback identifies the observable  $\sigma$ -algebra with the Borel algebra of the Stone space. CE appears here not as an algebraic condition but as a support condition: the measure concentrates on the principal ultrafilters, i.e., on the image of  $\Omega$  inside its Stone compactification.

**CE is irreducible.** Section 3 establishes that CE is not derivable from any finitary or structural condition on the query system. The finite-cofinite content on  $\mathbb{Q}$  satisfies every algebraic condition on a directed system while failing CE. A model-theoretic argument via Łoś's theorem and the Yosida–Hewitt decomposition shows this is not a proof-search failure: no first-order condition on the query system implies CE. CE is an irreducible commitment about the completed infinite, analogous to the axiom of infinity in arithmetic.

**The Stone space.** The Stone space  $\text{St}(\mathcal{C})$  of the observable algebra, constructed in §5 as a technical device for measure extension, has a further significance developed in §6. It is the canonical compact Hausdorff completion of  $\Omega$  for the observable topology. CE's two faces — honest observer (valuation layer) and support on  $\Omega$  (topological layer) — are the same condition seen from inside and outside the algebra. When an observation function is a cyclic vector for the Koopman operator of an underlying dynamical system, the Stone space of the observable algebra is measure-theoretically isomorphic to the state space. The compact space introduced here as a tool reappears at the end of the programme as the object being reconstructed; we develop this connection elsewhere.

**Organisation.** Section 2 introduces query systems, cylinder sets, the observable  $\sigma$ -algebra, compatible charges, and the three hypotheses (Surjective Evaluation, Discriminability, CE).

Section 3 establishes the CE theorem and its irreducibility. Section 4 proves the Carathéodory route. Section 5 proves the Stone route. Section 6 draws the two routes together and discusses the Stone space.

## 2 Setup: query systems and charges

### 2.1 Query systems

**Definition 2.1** (Query system). A *query system* is a tuple

$$(\iota, \leq, \{(\text{Out}(i), \mathcal{F}_i)\}_{i \in \iota}, \{\pi_{ij}\}_{i \leq j}, \Omega, \{\text{eval}_i\}_{i \in \iota})$$

consisting of:

- a nonempty partially ordered set  $(\iota, \leq)$ , the *index set*;
- for each  $i \in \iota$ , a measurable space  $(\text{Out}(i), \mathcal{F}_i)$ , the *outcome space at level  $i$* ;
- for each  $i \leq j$  in  $\iota$ , a measurable surjection  $\pi_{ij} : \text{Out}(j) \rightarrow \text{Out}(i)$ , the *refinement map*;
- a set  $\Omega$ , the *realization*, together with maps  $\text{eval}_i : \Omega \rightarrow \text{Out}(i)$  for each  $i \in \iota$ , the *evaluation maps*,

satisfying the *coherence condition*:  $\pi_{ij} \circ \text{eval}_j = \text{eval}_i$  for all  $i \leq j$ .

The index set, outcome spaces, and refinement maps constitute the *observational structure* of the query system; they encode the distinctions the observer can make and how finer levels coarsen to coarser ones. The realization  $\Omega$  and evaluation maps record how that structure is instantiated:  $\Omega$  is the set of states consistent with the system, and coherence says evaluation is compatible with coarsening.

### 2.2 Realization space

The observational structure  $(\iota, \leq, \{\text{Out}(i)\}, \{\pi_{ij}\})$  determines a canonical realization: the projective limit

$$\Omega = \varprojlim_{i \in \iota} \text{Out}(i) = \left\{ x \in \prod_{i \in \iota} \text{Out}(i) \mid \pi_{ij}(x_j) = x_i \text{ whenever } i \leq j \right\},$$

with  $\text{eval}_i(x) = x_i$ .

### 2.3 Directedness conditions

**Definition 2.2** (Common coarsenings and common refinements). The index set  $(\iota, \leq)$  admits *common coarsenings* if every pair  $i, j \in \iota$  has a lower bound: there exists  $k \leq i$  and  $k \leq j$ . It admits *common refinements* if every pair has an upper bound: there exists  $k \geq i$  and  $k \geq j$ . It admits *sequential common refinements* if every countable family  $\{i_n\}_{n \geq 1} \subset \iota$  has an upper bound in  $\iota$ .

Sequential common refinements imply common refinements. Common coarsenings ensure that cylinder sets form a  $\pi$ -system (Lemma 2.4 below); sequential common refinements supply the  $\sigma$ -subadditivity step in the Carathéodory route.

### 2.4 Cylinder sets and the observable $\sigma$ -algebra

**Definition 2.3** (Cylinder sets). For  $i \in \iota$  and  $A \in \mathcal{F}_i$ , the *level- $i$  cylinder with base  $A$*  is

$$\text{Cyl}(i, A) = \{\omega \in \Omega : \text{eval}_i(\omega) \in A\}.$$

The *cylinder generating family* is

$$\mathcal{C} = \{\text{Cyl}(i, A) : i \in \iota, A \in \mathcal{F}_i\}.$$

For each  $i \in \iota$ , let  $B_i$  denote the Boolean algebra of subsets of  $\Omega$  generated by  $\{\text{Cyl}(i, A) : A \in \mathcal{F}_i\}$ . Then  $\mathcal{C} = \bigcup_{i \in \iota} B_i$ .

The *observable  $\sigma$ -algebra* is  $\sigma(\mathcal{C})$ , the  $\sigma$ -algebra on  $\Omega$  generated by all cylinder sets.

The coherence condition gives, for  $i \leq j$ ,  $\text{Cyl}(i, A) = \text{Cyl}(j, \pi_{ij}^{-1}(A))$ , so the same subset of  $\Omega$  can be described as a cylinder at any finer level.

**Lemma 2.4** (Cylinder  $\pi$ -system). *Assume  $(\iota, \leq)$  admits common coarsenings. Then  $\mathcal{C}$  is a  $\pi$ -system: it is closed under finite intersections.*

*Proof.* Given  $\text{Cyl}(i, A)$  and  $\text{Cyl}(j, B)$ , let  $k$  be a common lower bound, with  $k \leq i$  and  $k \leq j$ . Then

$$\text{Cyl}(i, A) \cap \text{Cyl}(j, B) = \text{Cyl}(k, \pi_{ki}^{-1}(A) \cap \pi_{kj}^{-1}(B)),$$

which is again a cylinder.  $\square$

## 2.5 Connecting maps

For  $i \leq j$ , define the *connecting map*

$$\varphi_{ij} : B_i \rightarrow B_j, \quad \varphi_{ij}(\text{Cyl}(i, A)) = \text{Cyl}(j, \pi_{ij}^{-1}(A)).$$

This is a well-defined Boolean algebra homomorphism. The family  $\{B_i, \varphi_{ij}\}$  is a directed system of Boolean algebras.

## 2.6 Compatible charges

**Definition 2.5** (Compatible family). A *charge* on a Boolean algebra  $B$  is a finitely additive function  $\mu : B \rightarrow [0, \infty)$  with  $\mu(\emptyset) = 0$ . A family of charges  $\{\mu_i\}_{i \in \iota}$  with  $\mu_i$  on  $B_i$  is *compatible* if

$$\mu_j(\varphi_{ij}(E)) = \mu_i(E) \quad \text{for all } i \leq j \text{ and all } E \in B_i.$$

A compatible family also determines a charge on  $\mathcal{C}$  by  $\mu(\text{Cyl}(i, A)) := \mu_i(\text{Cyl}(i, A))$ ; compatibility ensures this is well-defined.

## 2.7 The three hypotheses

The main theorems require three conditions.

**Definition 2.6** (Surjective Evaluation). The query system satisfies *Surjective Evaluation* if  $\text{eval}_i : \Omega \rightarrow \text{Out}(i)$  is surjective for every  $i \in \iota$ .

**Definition 2.7** (Discriminability). The query system *discriminates points* if for every  $\omega \neq \omega'$  in  $\Omega$  there exists  $E \in \mathcal{C}$  with  $\omega \in E$  and  $\omega' \notin E$ .

**Definition 2.8** (Collective Exhaustion). A charge  $\mu$  on  $\mathcal{C}$  (the common extension of a compatible family  $\{\mu_i\}$ ) satisfies *collective exhaustion* (CE) if: whenever  $E_1 \supseteq E_2 \supseteq \dots$  in  $\mathcal{C}$  and  $\bigcap_n E_n = \emptyset$ , we have  $\mu(E_n) \rightarrow 0$ .

Surjective Evaluation is a structural condition ensuring that every outcome at every level is realized by some element of  $\Omega$ ; it implies that the refinement maps  $\pi_{ij}$  are surjective (by coherence) and that the connecting maps  $\varphi_{ij}$  are injective. Discriminability ensures the Stone embedding  $\omega \mapsto \{\text{principal ultrafilter at } \omega\}$  is injective. CE is the irreducible valuation-layer condition studied in §3; it rules out purely finitely additive charges and is the necessary and sufficient condition for  $\sigma$ -additive extensibility.

### 3 The CE theorem

The argument of §§3–5 has a single subject: the single condition under which a coherent family of observations determines a probability measure. It appears in four forms — as a local continuity criterion (§3, single algebra), as a global equivalence across the directed system (§3.3), as a direct measure-theoretic construction (§4), and as a topological derivation via compactness (§5). These are not separate results but four projections of a single constraint; §6 identifies the point where all four coincide.

This section establishes collective exhaustion as the necessary and sufficient condition for  $\sigma$ -additive extensibility. The analysis works at the level of an abstract directed system of Boolean algebras; it does not require the full query-system structure of §2. To keep this generality clean, we write  $\mathcal{E}_i$  for a generic Boolean algebra at level  $i$  and  $\ell_i$  for the charge on it. The connection to query systems is:  $\mathcal{E}_i = B_i$  (the Boolean algebra of cylinder sets at level  $i$ ),  $O_i = \text{Out}(i)$ , and  $\ell_i = \mu_i$ . The full query system structure re-enters in §4 and §5.

#### 3.1 Single-algebra extension

**Definition 3.1** (Boolean charge space). A *Boolean charge space* is a pair  $(\mathcal{E}, \ell)$  where  $\mathcal{E}$  is a Boolean algebra of subsets of a set  $O$  and  $\ell : \mathcal{E} \rightarrow [0, 1]$  is a normalized finitely-additive valuation. No  $\sigma$ -algebra and no topology are assumed on  $O$ .

Under Stone’s representation theorem [11],  $\mathcal{E}$  is isomorphic to the clopen algebra of a compact totally-disconnected Hausdorff space  $S(\mathcal{E})$ , and  $\sigma(\mathcal{E})$  is the Baire  $\sigma$ -algebra of  $S(\mathcal{E})$ . The gap  $\sigma(\mathcal{E}) \setminus \mathcal{E}$  consists of the Baire sets that are not clopen: events that countable operations can reach but that no single observable distinction can name.

**Theorem 3.2** (Extension criterion). A normalized finitely-additive  $\ell : \mathcal{E} \rightarrow [0, 1]$  extends to a  $\sigma$ -additive measure on  $\sigma(\mathcal{E})$  if and only if  $\ell$  is continuous at  $\emptyset$ : for every decreasing sequence  $E_1 \supseteq E_2 \supseteq \dots$  in  $\mathcal{E}$  with  $\bigcap_n E_n = \emptyset$  in  $\sigma(\mathcal{E})$ , we have  $\ell(E_n) \rightarrow 0$ . The extension is then unique.

*Proof.* Standard; see Halmos [6] or Fremlin [5] Vol. 1. Continuity at  $\emptyset$  is used in the Carathéodory extension argument to pass from finite additivity to  $\sigma$ -additivity on  $\sigma(\mathcal{E})$ .  $\square$

**Remark 3.3** (The gap and the regress). The condition in Theorem 3.2 is not verifiable from within  $\mathcal{E}$  alone: confirming  $\bigcap_n E_n = \emptyset$  requires access to  $\sigma(\mathcal{E})$ . In a directed system, one might hope that a finer level  $j \geq i$  can witness the emptiness that level  $i$  cannot see, and that compatibility forces continuity at  $\emptyset$  at level  $i$ . But the obstruction re-appears: finite additivity at level  $j$  does not force continuity at  $\emptyset$  for sequences decreasing to an element of  $\sigma(\mathcal{E}_j)$ . No finite chain of refinements can supply the condition. Something external to any single level is required: topology (compactness), a colimit algebra, or an explicit continuity axiom. Section 3.2 shows that the directed-system structure does not resolve this without an additional valuation condition.

#### 3.2 Nontrivial refinement

**Definition 3.4** (Nontrivial refinement). Let  $\pi : O_j \rightarrow O_i$  be a surjective refinement map, inducing the Boolean algebra homomorphism  $\pi^{-1} : \mathcal{E}_i \rightarrow \mathcal{E}_j$ . The refinement is *nontrivial* if there exists  $A \in \mathcal{E}_j$  with  $A \notin \pi^{-1}(\mathcal{E}_i)$ .

**Proposition 3.5** (Nontrivial refinement introduces new events). If  $j \geq i$  is a nontrivial refinement, then  $\pi^{-1}(\mathcal{E}_i) \subsetneq \mathcal{E}_j$ . Equivalently, a level  $i$  at which  $\mathcal{E}_j = \pi^{-1}(\mathcal{E}_i)$  for every  $j \geq i$  is one above which the refinement order is algebraically trivial: no finer level adds new discriminative power.

*Proof.* Nontriviality asserts the existence of  $A \in \mathcal{E}_j$  with  $A \notin \pi^{-1}(\mathcal{E}_i)$ , directly giving the strict inclusion. The contrapositive is the stated equivalence.  $\square$

**Proposition 3.6** (Gap nonemptiness). *Let  $\mathcal{E}$  be a Boolean algebra of subsets of a countably infinite set  $O$  that is not a  $\sigma$ -algebra. Then  $\sigma(\mathcal{E}) \setminus \mathcal{E} \neq \emptyset$ .*

*Proof.* If  $\mathcal{E}$  is not a  $\sigma$ -algebra, there exists a countable family  $\{A_n\} \subset \mathcal{E}$  with  $\bigcup_n A_n \notin \mathcal{E}$ . By definition  $\bigcup_n A_n \in \sigma(\mathcal{E})$ , so it lies in the gap.  $\square$

### 3.3 CE characterises $\sigma$ -additivity

We now state the directed-system version of CE and prove the main equivalence.

**Definition 3.7** (Collectively exhaustive family). A compatible family  $\{\ell_i\}$  of normalized finitely-additive charges on a directed system is *collectively exhaustive* if: for every  $i \in \mathcal{I}$  and every decreasing sequence  $E_1 \supseteq E_2 \supseteq \dots$  in  $\mathcal{E}_i$  with  $\bigcap_n E_n = \emptyset$ , there exists  $j \geq i$  such that  $\ell_j(\pi_{ij}^{-1}(E_n)) \rightarrow 0$ .

Here  $\pi_{ij} : O_j \rightarrow O_i$  is the refinement map ( $j$  finer than  $i$ ), so  $\pi_{ij}^{-1}(E_n) \subseteq O_j$  is the preimage of  $E_n \subseteq O_i$  at the finer level.

CE is a *valuation-layer* condition: it places a requirement on how the charges behave, not on the Boolean-algebraic index structure. It says that no level can assign persistent mass to events that the system as a whole sees as empty.

The following proposition establishes that the index-layer structure of a directed system — sequential upper-directedness and charge compatibility — does not on its own force  $\sigma$ -additivity.

**Proposition 3.8** (Index-layer conditions do not force  $\sigma$ -additivity). *There exists a compatible family of normalized finitely-additive charges on a sequentially upper-directed directed system that fails  $\sigma$ -additive extension at every level.*

*Proof.* Let  $\iota = \mathbb{N} \cup \{\omega\}$  with  $n \leq \omega$  for all  $n \in \mathbb{N}$ , directed by this order. Take all outcome spaces to be  $\mathbb{Q}$ , all event algebras to be the finite-cofinite algebra  $\mathcal{E} = \{E \subseteq \mathbb{Q} : E \text{ is finite or } E^c \text{ is finite}\}$ , all refinement maps to be the identity, and  $\ell(E) = 0$  if  $E$  is finite,  $\ell(E) = 1$  if  $E$  is cofinite. This is a normalized finitely-additive content and compatibility holds trivially.

Fix an enumeration  $q : \mathbb{N} \rightarrow \mathbb{Q}$  and let  $F_n = \{q(0), \dots, q(n)\}$ . Each  $F_n$  is finite so  $\ell(F_n) = 0$ , yet  $\bigcup_n F_n = \mathbb{Q}$  and  $\ell(\mathbb{Q}) = 1$ .  $\sigma$ -subadditivity would give  $1 \leq \sum_n \ell(F_n) = 0$ : contradiction.  $\square$

**Theorem 3.9** (CE Characterisation Theorem). *Let  $\{\ell_i\}$  be a normalized compatible family of finitely-additive charges on a directed system indexed by  $\mathcal{I}$ . The following are equivalent:*

- (i) *The family is collectively exhaustive.*
- (ii)  *$\ell_i$  extends uniquely to a  $\sigma$ -additive measure on  $\sigma(\mathcal{E}_i)$  for every  $i \in \mathcal{I}$ .*

*Proof.* (i)  $\Rightarrow$  (ii). Fix  $i \in \mathcal{I}$  and a decreasing sequence  $E_n \searrow \emptyset$  in  $\mathcal{E}_i$ . By CE, find  $j \geq i$  with  $\ell_j(\pi_{ij}^{-1}(E_n)) \rightarrow 0$ . By compatibility,  $\ell_i(E_n) = \ell_j(\pi_{ij}^{-1}(E_n)) \rightarrow 0$ . So  $\ell_i$  is continuous at  $\emptyset$ . Theorem 3.2 gives the unique  $\sigma$ -additive extension.

(ii)  $\Rightarrow$  (i). Suppose  $\ell_i$  extends to  $\mu_i$  for every  $i$ . Let  $E_n \searrow \emptyset$  in  $\mathcal{E}_i$ . Then  $\ell_i(E_n) = \mu_i(E_n) \rightarrow \mu_i(\emptyset) = 0$  by  $\sigma$ -additivity of  $\mu_i$  and normalization. Take  $j = i$ .  $\square$

**Remark 3.10** (Dissolution of the index-valuation conflation). Proposition 3.8 shows that index-layer conditions (sequential upper-directedness, compatibility) do not force  $\sigma$ -additivity. Theorem 3.9 identifies the exact valuation-layer condition that does. The question “do the structural conditions on a directed system force  $\sigma$ -additivity?” is ill-posed: it conflates conditions on the directed system’s structure with conditions on the charges it carries.

Sequential upper-directedness plays no role in Theorem 3.9: it is a per-level result. That condition enters when one asks whether the per-level extensions assemble into a global measure on the realization space  $\Omega = \varprojlim \text{Out}(i)$ ; this is the role it plays in the Carathéodory route (§4).

### 3.4 CE is irreducible

**Proposition 3.11** (CE irreducibility). *Collective exhaustion is not implied by any structural condition on a query system.*

- (i) (Particular case.) *Sequential upper-directedness, compatibility, and normalization do not imply CE.*
- (ii) (Universal case.) *No condition  $\Phi$  expressible in the first-order language of query systems implies CE.*

*Proof.* The particular case is Proposition 3.8: the finite-cofinite content satisfies all named structural hypotheses and fails CE.

For the universal case, we proceed in three steps.

*Step 1.* A first-order condition on query systems is preserved under ultraproducts by Łoś's theorem [9]: if  $\Phi$  holds in every member of a family of query systems, it holds in their ultraproduct.

*Step 2.* Let  $\mathcal{U}$  be a nonprincipal ultrafilter on  $\mathbb{N}$  extending the cofinite filter, and let  $\delta_n$  be the point mass at  $q(n) \in \mathbb{Q}$  (using the enumeration from Proposition 3.8). Each  $\delta_n$  is a normalized  $\sigma$ -additive content satisfying every structural condition including CE. The ultraproduct  $\prod_{\mathcal{U}} \delta_n$  assigns to  $E \subseteq \mathbb{Q}$  the value  $\lim_{\mathcal{U}} \mathbf{1}_{q(n) \in E}$ , which is 1 on cofinite sets and 0 on finite sets — exactly the finite-cofinite content  $\ell$  from Proposition 3.8.

*Step 3.* By Step 1, any universally valid first-order  $\Phi$  holds for  $\ell$ . But  $\ell$  fails CE. Therefore no first-order  $\Phi$  can imply CE.

The Yosida–Hewitt decomposition [13] explains why this obstruction is permanent. Every finitely-additive probability on a Boolean algebra decomposes uniquely as  $\ell = \ell_c + \ell_p$ , where  $\ell_c$  is  $\sigma$ -additive and  $\ell_p$  is *purely finitely additive*: it satisfies every algebraic condition while assigning zero to every singleton. The finite-cofinite content  $\ell$  is purely finitely additive ( $\ell_c = 0$ ). CE is exactly the condition that rules out a nonzero purely finitely additive part:  $\ell_p = 0$ .  $\square$

**Remark 3.12** (CE as irreducible primitive). Proposition 3.11 shows that the gap between structural coherence and  $\sigma$ -additivity is a proved boundary, not a proof-search failure. CE is not derivable within the query-system framework. It is analogous to the axiom of infinity in arithmetic: not provable from the remaining axioms, not refutable, but the commitment without which the mathematics of probability cannot proceed. The ultrafilter-based content is the canonical witness of this irreducibility: it passes every finite consistency test yet assigns unit mass to events whose infinite intersection is empty.

Formally: CE is not a constraint the query system imposes on the observer. It is a commitment the observer makes about the completed infinite, a condition on how their charges behave with respect to limits that no finite stage can verify.

Philosophically: CE is the *principle of witnessed vanishing*. A distinction may disappear globally — an event may shrink to nothing in the limit of the observer's refinements — only if that disappearance is witnessed at some finite level. There is no global collapse without local evidence; no vanishing that the system's refinement order cannot eventually confirm. This is not a metaphysical assumption but an exact mathematical condition, and Proposition 3.11 shows it cannot be reduced further: no weaker structural commitment produces the same effect.

## 4 Route 1: The Carathéodory extension

This section establishes the extension theorem by a direct Carathéodory argument. This is the third face of the argument: the measure constructed directly from within the algebra. The input is a query system satisfying Surjective Evaluation, sequential common refinements, and common coarsenings, together with a compatible family of  $\sigma$ -additive marginals. By Theorem 3.9, the  $\sigma$ -additivity of the marginals is equivalent to CE applied at each level.

### 4.1 Observational determination

**Proposition 4.1** (Observational determination). *Let the query system admit common coarsenings and let  $P, P'$  be probability measures on  $(\Omega, \sigma(\mathcal{C}))$ . If  $(\text{eval}_i)_\# P = (\text{eval}_i)_\# P'$  for all  $i \in \iota$ , then  $P = P'$ .*

*Proof.* The equality of pushforward laws implies  $P$  and  $P'$  agree on every cylinder event. By Lemma 2.4 the cylinder family  $\mathcal{C}$  is a  $\pi$ -system generating  $\sigma(\mathcal{C})$ . The  $\pi$ - $\lambda$  theorem gives  $P = P'$ .  $\square$

### 4.2 Observable content

**Lemma 4.2** (Finite observational content). *Assume the query system admits common coarsenings and common refinements and satisfies Surjective Evaluation, and let  $\{\nu_i\}_{i \in \iota}$  be an observationally compatible family with each  $\nu_i \in \mathcal{P}(\text{Out}(i))$ . Then the assignment*

$$\mu_0(\text{Cyl}(i, A)) := \nu_i(A), \quad A \in \mathcal{F}_i,$$

*defines a well-defined finitely additive content on  $\mathcal{C}$ .*

*Proof. Well-definedness.* Suppose  $\text{Cyl}(i, A) = \text{Cyl}(j, B)$ . Use common refinements to find  $k$  with  $k \geq i$  and  $k \geq j$ . The constraint sets in  $\text{Out}(k)$  are  $\pi_{ik}^{-1}(A)$  and  $\pi_{jk}^{-1}(B)$ . The cylinder equality and Surjective Evaluation (surjectivity of  $\text{eval}_k$ ) imply these coincide in  $\text{Out}(k)$ . Compatibility then gives  $\nu_i(A) = \nu_k(\pi_{ik}^{-1}(A)) = \nu_k(\pi_{jk}^{-1}(B)) = \nu_j(B)$ .

*Finite additivity.* For disjoint cylinders at possibly different levels, compress to a common refinement  $k$  and use additivity of  $\nu_k$ .  $\square$

### 4.3 Carathéodory extension theorem

**Theorem 4.3** (Observational extension theorem). *Suppose:*

- (i)  $(\iota, \leq)$  admits common coarsenings and sequential common refinements;
- (ii) the query system satisfies Surjective Evaluation;
- (iii) the family  $\{\nu_i\}_{i \in \iota}$  is observationally compatible and each  $\nu_i$  is a probability measure on  $(\text{Out}(i), \mathcal{F}_i)$ .

*Then there exists a unique probability measure  $P$  on  $(\Omega, \sigma(\mathcal{C}))$  such that  $(\text{eval}_i)_\# P = \nu_i$  for every  $i \in \iota$ .*

*Proof. Step 1: Finite content.* Theorem 4.2 gives a well-defined finitely additive content  $\mu_0$  on  $\mathcal{C}$ .

*Step 2:  $\sigma$ -subadditivity.* Let  $\text{Cyl}(i, B) \subseteq \bigcup_{n \geq 1} \text{Cyl}(i_n, A_n)$ . Apply sequential common refinements to  $\{i, i_1, i_2, \dots\}$  to find a single index  $k$  with  $k \geq i$  and  $k \geq i_n$  for all  $n$ . Compress each cylinder to level  $k$ :

$$\text{Cyl}(i, B) = \text{Cyl}(k, E), \quad \text{Cyl}(i_n, A_n) = \text{Cyl}(k, E_n),$$

where  $E = \pi_{ik}^{-1}(B)$  and  $E_n = \pi_{ink}^{-1}(A_n)$ . Surjective Evaluation and the cylinder inclusion in  $\Omega$  imply  $E \subseteq \bigcup_n E_n$  in  $\text{Out}(k)$ . Therefore

$$\mu_0(\text{Cyl}(i, B)) = \nu_k(E) \leq \sum_{n=1}^{\infty} \nu_k(E_n) = \sum_{n=1}^{\infty} \mu_0(\text{Cyl}(i_n, A_n)),$$

using  $\sigma$ -subadditivity of the single measure  $\nu_k$ .

*Step 3: Carathéodory extension.* The family  $\mathcal{C}$  is a semiring generating  $\sigma(\mathcal{C})$  and  $\mu_0$  is a  $\sigma$ -subadditive finitely additive content on it. Carathéodory's extension theorem [1, 2] applies and yields  $P$  on  $(\Omega, \sigma(\mathcal{C}))$ .

*Step 4: Marginal recovery and normalization.* For any  $A \in \mathcal{F}_i$ ,  $P(\text{Cyl}(i, A)) = \mu_0(\text{Cyl}(i, A)) = \nu_i(A)$ , so  $(\text{eval}_i)_\# P = \nu_i$ . Setting  $A = \text{Out}(i)$  gives  $P(\Omega) = 1$ .

*Uniqueness.* Two probability measures with the same evaluation marginals agree on  $\mathcal{C}$ , hence on  $\sigma(\mathcal{C})$  by Theorem 4.1.  $\square$

**Remark 4.4** (Relation to Kolmogorov's extension theorem). Kolmogorov's theorem [8] constructs a probability measure on  $S^T$  from consistent finite-dimensional distributions. Theorem 4.3 plays the same role for query systems: the realization space  $\Omega = \varprojlim \text{Out}(i)$  replaces the product  $S^T$ , compatible marginals replace consistent finite-dimensional distributions, and sequential common refinements replace the countable-cofinality condition. The key conceptual difference is that no topology is imposed on the outcome spaces: the measure  $P$  is built directly on the measurable structure of  $\Omega$  by Carathéodory, without any appeal to tightness or compactness.

**Remark 4.5** (Prokhorov route). A third route to the extension theorem starts from *finitely* additive marginals and derives  $\sigma$ -additivity from topological compactness. Under *surjective projective uniform tightness* — a condition requiring that for every  $\varepsilon > 0$  there exist compact sets  $K_i \subset \text{Out}(i)$  with  $\nu_i(\text{Out}(i) \setminus K_i) < \varepsilon$  and  $\pi_{ij}(K_j) = K_i$  for all  $i \leq j$  — the Prokhorov theorem extends finitely additive marginals on a countable sequentially upper-directed query system with Hausdorff outcome spaces to a  $\sigma$ -additive global measure. This route assumes topological structure on the outcome spaces and is complementary to the Carathéodory and Stone routes, which require no topology.

**Remark 4.6** (Lean formalization). Theorem 4.3 is fully formalized in Lean 4 (`observational_extension` in `QuerySystem.lean`, 0 sorrys). The hypotheses correspond exactly to the four conditions: `LowerDirected`, `SequentiallyUpperDirected`, `EvalSurjective`, and `CompatibleMarginals` with `IsProbabilityMeasure`. The Carathéodory step uses `AddContent.measure` from `Mathlib` [12].

## 5 Route 2: The Stone extension

This section gives the second, independent proof of the extension theorem. This is the fourth face: the same measure derived from outside, via compactness. The input is weaker: charges need only be finitely additive.  $\sigma$ -additivity is derived from the compactness of the Stone space.

The proof proceeds in four steps. Step A identifies the Stone space of the direct limit with the inverse limit of individual Stone spaces. The inverse limit measure step uses Choksi's theorem to produce a  $\sigma$ -additive measure on the inverse limit. Step B1 identifies the pullback of the Borel algebra of the Stone space with the observable  $\sigma$ -algebra. Step B2 shows that CE forces the measure to concentrate on the image of  $\Omega$ .

## 5.1 Step A: Stone space of the direct limit

**Lemma 5.1** (Injectivity of connecting maps). *If the query system satisfies Surjective Evaluation, then  $\varphi_{ij} : B_i \rightarrow B_j$  is an injective Boolean algebra homomorphism for every  $i \leq j$ .*

*Proof.* Surjective Evaluation and the coherence condition give  $\pi_{ij}(\text{eval}_j(\omega)) = \text{eval}_i(\omega)$  for all  $\omega$ , so  $\text{im}(\pi_{ij}) \supseteq \text{im}(\text{eval}_i) = \text{Out}(i)$ : the refinement maps are surjective. A surjective map  $\pi_{ij}$  has injective preimage map  $\pi_{ij}^{-1}$  on subsets, so  $\varphi_{ij}(\text{Cyl}(i, A)) = \text{Cyl}(j, \pi_{ij}^{-1}(A))$  is injective on generators, hence on  $B_i$ .  $\square$

**Proposition 5.2** (Step A). *The Stone space of the direct limit satisfies  $\text{St}(\bigcup_i B_i) \cong \varprojlim_i \text{St}(B_i)$ .*

*Proof.* By Lemma 5.1, the connecting maps  $\varphi_{ij}$  are injective Boolean algebra homomorphisms. Under Stone duality, injective Boolean algebra homomorphisms correspond to surjective continuous maps between Stone spaces. The Stone space functor  $\text{St}$  converts colimits in the category of Boolean algebras to limits in the category of compact Hausdorff spaces [7, II.4]. Applied to  $\bigcup_i B_i$  with its injective system, this gives  $\text{St}(\bigcup_i B_i) \cong \varprojlim_i \text{St}(B_i)$ .  $\square$

## 5.2 The inverse limit measure

**Proposition 5.3** (Inverse limit measure). *Let  $\{\mu_i\}$  be a compatible family of normalised charges on  $\{B_i\}$ . Then there exists a unique probability measure  $\hat{\mu}$  on  $\mathcal{B}(\varprojlim_i \text{St}(B_i))$  whose pushforward along each projection  $\varprojlim_i \text{St}(B_i) \rightarrow \text{St}(B_j)$  agrees with the Borel measure on  $\text{St}(B_j)$  corresponding to  $\mu_j$ .*

*Proof. Step 1: Single-algebra correspondence.* For each  $i$ , Stone duality identifies  $B_i$  with the clopen algebra  $\text{Clop}(\text{St}(B_i))$ . A charge  $\mu_i$  on  $B_i$  thus defines a finitely additive set function on the clopens of the compact Hausdorff space  $\text{St}(B_i)$ . Since  $\text{St}(B_i)$  is compact and totally disconnected, the clopens form a base for the topology and a finitely additive charge on them extends uniquely to a regular Borel measure; see [3], §6, or [6], §§53–54. Call this measure  $\hat{\mu}_i$  on  $\text{St}(B_i)$ .

*Step 2: Compatibility.* The compatibility condition  $\mu_j(\varphi_{ij}(E)) = \mu_i(E)$  translates, under Stone correspondence, to  $\hat{\mu}_i$  being the pushforward of  $\hat{\mu}_j$  along the bonding map  $\text{St}(B_j) \rightarrow \text{St}(B_i)$ . The family  $\{\hat{\mu}_i\}$  is compatible on the inverse system  $\{\text{St}(B_i)\}$ .

*Step 3: Extension to the inverse limit.* Each  $\text{St}(B_i)$  is compact Hausdorff. The inverse limit  $\varprojlim_i \text{St}(B_i)$  is compact Hausdorff. By [4, Theorem 1], a compatible family of regular Borel probability measures on a cofiltered inverse system of compact Hausdorff spaces with surjective bonding maps extends uniquely to a regular Borel probability measure on the inverse limit. This gives  $\hat{\mu}$  on  $\mathcal{B}(\varprojlim_i \text{St}(B_i))$ .  $\square$

## 5.3 Step B1: the structural identification

Let  $\text{pure} : \Omega \rightarrow \text{St}(\mathcal{C})$  denote the Stone embedding, which sends  $\omega \in \Omega$  to its principal ultrafilter  $\text{pure}(\omega) = \{E \in \mathcal{C} : \omega \in E\}$ . For  $E \in \mathcal{C}$ , write  $[E] = \{u \in \text{St}(\mathcal{C}) : E \in u\}$  for the corresponding basic clopen in  $\text{St}(\mathcal{C})$ .

**Proposition 5.4** (Step B1: structural identification). *Suppose the query system discriminates points. Then*

$$\text{pure}^{-1}(\mathcal{B}_{\text{Ba}}(\text{St}(\mathcal{C}))) = \sigma(\mathcal{C}),$$

where  $\mathcal{B}_{\text{Ba}}$  denotes the Baire  $\sigma$ -algebra.

*Proof. ( $\supseteq$ ):* For any  $E = \text{Cyl}(i, A) \in \mathcal{C}$ ,  $\text{pure}^{-1}([E]) = \{\omega : E \in \text{pure}(\omega)\} = \{\omega : \omega \in E\} = E$ . Since  $[E]$  is clopen, hence Baire, every cylinder set is in  $\text{pure}^{-1}(\mathcal{B}_{\text{Ba}})$ , and therefore  $\sigma(\mathcal{C}) \subseteq \text{pure}^{-1}(\mathcal{B}_{\text{Ba}})$ .

( $\subseteq$ ): The Stone space  $\text{St}(\mathcal{C})$  is compact and totally disconnected, so its clopens form a topological basis. The clopens are exactly the sets  $[E]$  for  $E \in \mathcal{C}$  [7, I.3.2], so the Baire  $\sigma$ -algebra of  $\text{St}(\mathcal{C})$  is  $\sigma(\{[E] : E \in \mathcal{C}\})$ . For any Baire set  $V$ ,  $\text{pure}^{-1}(V) \in \sigma(\{\text{pure}^{-1}([E]) : E \in \mathcal{C}\}) = \sigma(\mathcal{C})$ .

**Injectivity.** Discriminability implies  $\text{pure}$  is injective: if  $\omega \neq \omega'$ , there exists  $E \in \mathcal{C}$  with  $\omega \in E$  and  $\omega' \notin E$ , so  $E \in \text{pure}(\omega)$  but  $E \notin \text{pure}(\omega')$ . Injectivity ensures the pullback does not conflate distinct events.  $\square$

**Remark 5.5.** On a compact Hausdorff space the Baire  $\sigma$ -algebra is contained in the Borel  $\sigma$ -algebra, with equality when the space is second-countable. We work throughout with the Baire  $\sigma$ -algebra on  $\text{St}(\mathcal{C})$ ; this suffices because the measure  $\hat{\mu}$  constructed above is a Baire measure (it is the unique extension of a charge on the clopen algebra), and Choksi's theorem applies to Baire measures on compact spaces.

## 5.4 Step B2: CE and the support condition

**Proposition 5.6** (Step B2: support condition). *Let  $\mu$  be a charge on  $\mathcal{C}$  satisfying CE. Then the corresponding Baire measure  $\hat{\mu}$  on  $\text{St}(\mathcal{C})$  is supported on  $\text{pure}(\Omega)$ , the set of principal ultrafilters.*

*Proof.* By the Yosida–Hewitt theorem [13, 10], every bounded charge on a Boolean algebra decomposes uniquely as  $\mu = \mu_c + \mu_p$ , where  $\mu_c$  is countably additive and  $\mu_p$  is purely finitely additive. Under Stone duality,  $\mu_c$  corresponds to mass on the principal ultrafilters  $\text{pure}(\Omega) \subset \text{St}(\mathcal{C})$ , while  $\mu_p$  corresponds to mass on the non-principal ultrafilters [10], Ch. 10.

CE is equivalent to  $\mu_p = 0$ : CE requires  $\mu(E_n) \rightarrow 0$  whenever  $E_n \searrow \emptyset$ , which is exactly the condition that the purely finitely additive part vanishes (Theorem 3.9). Therefore  $\hat{\mu}_p = 0$  and  $\hat{\mu}$  is supported on  $\text{pure}(\Omega)$ .  $\square$

## 5.5 The Stone extension theorem

**Theorem 5.7** (Stone extension theorem). *Let the query system satisfy Surjective Evaluation, Discriminability, and common coarsenings, and let  $\{\mu_i\}$  be a compatible family of normalised charges on  $\{B_i\}$  satisfying CE. Then there exists a unique probability measure  $P$  on  $(\Omega, \sigma(\mathcal{C}))$  such that*

$$P(\text{Cyl}(i, A)) = \mu_i(\text{Cyl}(i, A)) \quad \text{for all } i \in \iota \text{ and } A \in \mathcal{F}_i.$$

*Proof.* The proof assembles the preceding propositions:

$$\{\mu_i\} \xrightarrow{\S 5.2, \text{Step 1}} \{\hat{\mu}_i\} \xrightarrow{\S 5.2, \text{Step 3}} \hat{\mu} \text{ on } \varprojlim_i \text{St}(B_i) \xrightarrow{\text{B1}} P \text{ on } \sigma(\mathcal{C}).$$

*Existence.* By Proposition 5.2,  $\text{St}(\mathcal{C}) \cong \varprojlim_i \text{St}(B_i)$ . By Proposition 5.3, the compatible family  $\{\mu_i\}$  extends to a Baire probability measure  $\hat{\mu}$  on  $\varprojlim_i \text{St}(B_i) \cong \text{St}(\mathcal{C})$ .

By Proposition 5.6,  $\hat{\mu}$  is supported on  $\text{pure}(\Omega)$ . Since  $\text{pure}$  is injective (Discriminability), the pushforward  $P = (\text{pure})_* \hat{\mu}$  — the image measure under  $\text{pure}^{-1}$  on the support — is a well-defined probability measure on  $\Omega$ .

By Proposition 5.4,  $\text{pure}^{-1}$  maps Baire sets in  $\text{St}(\mathcal{C})$  to sets in  $\sigma(\mathcal{C})$ , so  $P$  is defined on  $\sigma(\mathcal{C})$ . For any cylinder set  $E = \text{Cyl}(i, A)$ ,

$$P(E) = \hat{\mu}([E]) = \mu_i(\text{Cyl}(i, A)).$$

*Uniqueness.* Two probability measures on  $(\Omega, \sigma(\mathcal{C}))$  agreeing on all cylinder sets are equal by Theorem 4.1, which requires common coarsenings. The Stone theorem therefore assumes common coarsenings in addition to Surjective Evaluation, Discriminability, and CE.  $\square$

**Remark 5.8** (CE is not bypassed by compactness). It may appear that the compactness of the Stone space eliminates the need for CE. Compactness does ensure that  $\hat{\mu}$  on  $\text{St}(\mathcal{C})$  is  $\sigma$ -additive. But  $\hat{\mu}$  lives on the Stone space, not on  $\Omega$ . The descent from  $\text{St}(\mathcal{C})$  to  $\Omega$  requires that no mass escapes to the non-principal ultrafilters, and that is precisely CE (Proposition 5.6). In the Stone picture, CE is not an additional hypothesis but a *support condition*: the measure concentrates on the image of the sample space inside its Stone compactification.

**Remark 5.9** (Lean formalization). The Stone route is formalized in Lean 4 in `StoneDualityExtension.lean`. The structural results (Tasks 0'-A and 0'-C: Stone space, cylinder clopens, evaluation maps, bonding maps) are fully proved with no sorry obligations. Two intentional sorrys mark Mathlib gaps: the extension of a clopen-algebra charge to a regular Borel measure on a compact totally-disconnected space (Halmos §§53–54; Fremlin Vol. 1 §311E), and Choksi's projective-limit theorem for compatible measures on cofiltered inverse systems of compact Hausdorff spaces. The coincidence of routes (Task 0'-E: `stone_agrees_with_caratheodory`) is fully proved, as an immediate consequence of `observational_determination`.

## 6 The bridge

### 6.1 Two routes, one theorem

Under shared hypotheses — Surjective Evaluation, Discriminability, and  $\sigma$ -additive compatible marginals — both routes produce the same measure.

**Proposition 6.1** (Coincidence of routes). *Suppose the query system satisfies Surjective Evaluation, Discriminability, and sequential common refinements and common coarsenings, and let  $\{\nu_i\}$  be a compatible family of probability measures. Let  $P^C$  be the measure produced by Theorem 4.3 and  $P^S$  the measure produced by Theorem 5.7. Then  $P^C = P^S$ .*

*Proof.* Both measures agree on every cylinder set:  $P^C(\text{Cyl}(i, A)) = \nu_i(A) = P^S(\text{Cyl}(i, A))$ . By observational determination (Theorem 4.1), they are equal on  $\sigma(\mathcal{C})$ .  $\square$

The two routes are genuinely independent proofs of the same theorem. The Carathéodory route works directly within measure theory; the Stone route takes a detour through topology, using compactness of the Stone space to derive  $\sigma$ -additivity that the Carathéodory route assumes. That they produce the same measure is not obvious from the constructions alone; it is a consequence of uniqueness (Theorem 4.1), which is doing real work here.

### 6.2 CE's two faces

CE appears in the two routes in structurally different forms, yet it is the same condition.

In the Carathéodory route (Route 1), CE appears at the level of the charges: the per-level marginals are  $\sigma$ -additive, which by Theorem 3.9 is equivalent to CE applied at each level. Here, CE is an *algebraic* condition: the charges do not assign persistent mass to events that the observer's refinement hierarchy sees as empty. It is a condition on the observer's commitments.

In the Stone route (Route 2), CE appears as a *topological* condition: the Baire measure  $\hat{\mu}$  on the Stone space must be supported on  $\text{pure}(\Omega)$ , the image of the sample space inside its compact compactification. Mass on non-principal ultrafilters corresponds exactly to the purely finitely additive part  $\mu_p$ , and  $\text{CE} \Leftrightarrow \mu_p = 0$  (Proposition 5.6). Here, CE is a *geometric* condition: the observer's measure does not escape to the “phantom” points of the Stone compactification.

The Yosida–Hewitt decomposition is the bridge between the two faces: it identifies the purely finitely additive part of  $\mu$  with the mass that has escaped to non-principal ultrafilters. CE says both things simultaneously: the algebraic convergence is honest, and the geometric measure is confined to  $\Omega$ .

### 6.3 The Stone space as compact completion

The Stone space  $\text{St}(\mathcal{C})$  constructed in §5 as a technical device for measure extension is, in fact, the canonical compact Hausdorff completion of the sample space for the observable topology.

More precisely: the Stone embedding  $\text{pure} : \Omega \rightarrow \text{St}(\mathcal{C})$  is a dense embedding when  $\Omega$  is dense in  $\text{St}(\mathcal{C})$  for the Stone topology — i.e., when every nonempty basic clopen  $[E]$  meets  $\text{pure}(\Omega)$ , which holds whenever  $E \neq \emptyset$  as a subset of  $\Omega$ . The Stone space is uniquely determined by  $\mathcal{C}$  up to homeomorphism. It is not a construction imposed on  $\Omega$  from outside but the compactification already contained in the Boolean algebra structure of the cylinder sets.

**Remark 6.2** (The Stone space as space of consistent distinction patterns). Every point in  $\text{St}(\mathcal{C})$  is an ultrafilter: a maximal consistent assignment of yes/no answers to every observable distinction. Principal ultrafilters correspond to actual sample-space points — the distinctions that  $\Omega$  witnesses directly. Non-principal ultrafilters are the ideal limit points: consistent distinction patterns that the observer’s refinement system is coherent with, but that no single element of  $\Omega$  realises. The Stone space is therefore not an external topological gadget but the *space of all consistent distinction patterns* — the completion of the observer’s world to a compact whole. CE says the measure stays within the realised part: the observer is committed to their world, not to its ideal extensions.

CE is precisely the condition that the measure  $P$  on  $\Omega$ , when viewed as a measure on its compact completion  $\text{St}(\mathcal{C})$  via  $\text{pure}$ , does not escape: it remains concentrated on the original (non-compact) sample space. In this sense, an observer satisfying CE “knows they are in  $\Omega$ ”: their probability assignments, however they are extended to the compact hull, retain no phantom mass.

**Remark 6.3** (Topology from the query system). The Stone space  $\text{St}(\mathcal{C})$  carries a canonical topology: the one generated by all cylinder clopens  $[E]$ . This is the coarsest topology making every observable distinction continuous. A natural question, deferred to later work, is whether the *relevant* topology on  $\Omega$  — the one appropriate for dynamics and reconstruction — is this Stone topology, or whether it should itself be derived from the query system and its charges. When separation is weighted by the measure  $P$ , a third regime appears beyond the usual distinguishable/indistinguishable binary: pairs that are formally separated at every level but whose separating events have  $P$ -measure tending to zero. These are points that the observer’s valuation treats as near even though the Boolean algebra keeps them apart. Whether this measure-weighted nearness assembles into a topology, and whether collective exhaustion and this separation condition are two faces of a single coherence requirement, is an open question. The Boolean framework adopted here may accordingly be a first approximation: while distinctions are represented as crisp yes/no separations, the measure structure introduced in this paper suggests that effective separation is graded, and that a fuller account might refine the binary distinguishable/indistinguishable dichotomy into something more continuous.

**Connection to reconstruction.** For a dynamical system  $(X, T, \mu)$  with observation function  $h$ , the time-delay cylinder algebras  $\{B_S\}$  (indexed by finite subsets  $S$  of delay times) form a directed system of exactly the kind treated here. The Stone space of the resulting observable algebra  $\sigma(B_\infty)$  is, when  $h$  is a *cyclic vector* for the Koopman operator  $U_T$  — i.e., the orbit  $\{h \circ T^n : n \in \mathbb{Z}\}$  spans a dense subspace of  $L^2(X, \mu)$  — measure-theoretically isomorphic to the state space  $X$  itself.

The compact space  $\text{St}(\mathcal{C})$  introduced here as a tool for measure extension is, under the reconstruction condition, the state space being reconstructed. The programme begins with a query system whose Stone space is a compact completion of  $\Omega$ , and ends with the recognition that this compact space is the state space of the dynamical system the observer is embedded in. We develop this connection in a separate paper.

## References

- [1] Patrick Billingsley. *Probability and Measure*. Wiley, 1995.
- [2] Vladimir I. Bogachev. *Measure Theory, Volumes I–II*. Springer, 2007.
- [3] Rodrigo Cardona, Diego Alejandro Mejía, and Iván Uribe-Zapata. Finitely additive measures on Boolean algebras. arXiv preprint arXiv:2503.08910, 2025.
- [4] J. R. Choksi. Inverse limits of measure spaces. *Proceedings of the London Mathematical Society*, 8(3):321–342, 1958.
- [5] D. H. Fremlin. *Measure Theory*, volume 1. Torres Fremlin, 2003.
- [6] Paul R. Halmos. *Measure Theory*. Van Nostrand, 1950.
- [7] Peter T. Johnstone. *Stone Spaces*, volume 3 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1982.
- [8] A. N. Kolmogorov. *Foundations of the Theory of Probability*. Chelsea, 1933.
- [9] Jerzy Łoś. Quelques remarques, théorèmes et problèmes sur les classes définissables d’algèbres. In *Mathematical Interpretation of Formal Systems*, pages 98–113. North-Holland, 1955.
- [10] K. P. S. Bhaskara Rao and M. Bhaskara Rao. *Theory of Charges: A Study of Finitely Additive Measures*. Academic Press, New York, 1983.
- [11] Marshall Harvey Stone. The theory of representations for Boolean algebras. *Transactions of the American Mathematical Society*, 40(1):37–111, 1936.
- [12] The Mathlib Community. Mathlib4: the Lean 4 mathematical library. <https://github.com/leanprover-community/mathlib4>, 2024.
- [13] Kôzaku Yosida and Edwin Hewitt. Finitely additive measures. *Transactions of the American Mathematical Society*, 72(1):46–66, 1952.