

A σ -Essential Contextual State in ZFC: Localization and the Product Ulam Witness

Patrick S. Mahon*

Abstract

A two-valued state on a concrete quantum logic of subsets of a set Ω assigns a definite truth value to each event, coherently across overlapping measurement contexts. On a Boolean σ -algebra every finitely coherent local pattern of such assignments extends to a global σ -additive two-valued state. We ask whether the non-distributive case can differ: does there exist a concrete σ -complete non-Boolean orthomodular poset carrying a finitely coherent local pattern that extends to *no* global σ -additive two-valued state — a *σ -essential contextual state*? We first localize the question: a witness exists iff the pattern is finitely coherent while no point evaluation and no non-principal σ -additive state extends it, and we map its boundary — empty in the Polish-representable case (Derr–Williamson), irreducible to the Hilbert-space dichotomy of Blecher–Weaver, irreducible to a measurable cardinal. We then answer it affirmatively in ZFC. The witness is a *product Ulam carrier* on $\Omega = \omega_1 \times \{1, 2, 3, 4\}$: an Ulam matrix imported as generators makes every σ -additive two-valued state a point evaluation, a Specker triple of fibre events has empty kernel, and a trace-invariant normal form shows the two mechanisms cannot interact, so that a finitely additive extension survives, built from an arbitrary ultrafilter on $\mathcal{P}(\omega_1)/\text{ctble}$ through a parity code. Consequently the target sentence carries no large-cardinal strength. The quotient of the carrier by the countable ideal is a σ -complete non-Boolean orthomodular poset with finitely additive two-valued states but no σ -additive ones at all. The lattice (OML) form of the question remains open, and we conjecture it fails: latticehood appears to force σ -liftability.

1 Introduction

Contextuality is typified by the absence of a single underlying state realizing a consistent assignment of truth values to a finite family of events across overlapping measurement contexts. In finite scenarios one need not distinguish whether it is σ -additive or finitely additive states that are precluded: a finite carrier has no countable orthogonal families beyond the finite ones, so every finitely additive state is already σ -additive. Demanding countable additivity sharpens the question: can a *finitely* coherent pattern extend to no global σ -additive state, the failure visible only at the σ -level?

We isolate this phenomenon for concrete quantum logics, reduce it to a set-theoretic core, locate that core against the measure-theoretic and operator-algebraic literature, and resolve it affirmatively in ZFC by an explicit construction, the *product Ulam carrier*. Throughout we work with two-valued states, the non-distributive analogue of a $\{0, 1\}$ -valued measure.

A *concrete σ -complete orthomodular poset* (concrete σ -orthoposet, σ -class) is a family $\mathcal{L} \subseteq \mathcal{P}(\Omega)$ with

$$\emptyset, \Omega \in \mathcal{L}; \quad A \in \mathcal{L} \implies A^\perp := \Omega \setminus A \in \mathcal{L}; \quad \{A_n\} \subseteq \mathcal{L} \text{ pairwise disjoint} \implies \bigcup_n A_n \in \mathcal{L},$$

ordered by inclusion. This is the σ -class of Navara–Pták [19], a Gudder concrete quantum logic [11, 20]: the maximal Boolean subalgebras (“blocks”) overlap but their union need not

*Research Computing Group, Simon Fraser University, Burnaby, BC, Canada. pmahon@sfu.ca

be Boolean. (The σ -class axioms above, which are what every argument uses, do not provide lattice meets, and the witness constructed in §5 is provably *not* a lattice; “poset,” not “lattice,” is the operative term throughout. The lattice form of the problem remains open; see §6.) A *two-valued state* is $s: \mathcal{L} \rightarrow \{0, 1\}$ with $s(\Omega) = 1$ that is additive over orthogonal joins lying in its domain: whenever $\{A_n\} \subseteq \mathcal{L}$ is a *finite* pairwise disjoint family with $\bigcup_n A_n \in \mathcal{L}$,

$$s(\bigcup_n A_n) = \sum_n s(A_n)$$

(in particular $s(A) + s(A^\perp) = 1$ on complement pairs); it is σ -*additive* when this holds for countable such families. The same additivity is required of states on a sub-orthoposet $B \subseteq \mathcal{L}$, over orthogonal joins that exist in B . The *point evaluations* $\delta_\omega(A) := \mathbf{1}[\omega \in A]$ are σ -additive two-valued states; call these *Dirac* and all others *non-Dirac*. For a finite $\perp$ -closed sub-orthoposet $B \subseteq \mathcal{L}$ and a two-valued state s_0 on B , s_0 *extends* to s on $\mathcal{L}$ if $s \upharpoonright B = s_0$.

Definition 1.1 (finite coherence). Let $B \subseteq \mathcal{L}$ be a finite $\perp$ -closed sub-orthoposet and s_0 a two-valued state on B . Say s_0 is *finitely coherent on $\mathcal{L}$* if s_0 extends to a (finitely additive) two-valued state on all of $\mathcal{L}$.

Definition 1.2 (σ -essential contextual state). A σ -*essential contextual state* on $\mathcal{L}$ is a two-valued state s_0 on a finite $\perp$ -closed sub-orthoposet $B \subseteq \mathcal{L}$ that is finitely coherent on $\mathcal{L}$ but extends to *no* global σ -additive two-valued state on $\mathcal{L}$. A concrete σ -orthoposet $\mathcal{L}$ *carries a σ -essential contextual state* (is a *witness*) if such an s_0 exists. We require $\mathcal{L}$ to be non-Boolean and *essentially irreducible*: the quotient of $\mathcal{L}$ by the σ -ideal of countable sets has centre $\{0, 1\}$; see Remark 1.5.

Remark 1.3 (Why the coherence clause is necessary). Without Definition 1.1 the definition of a witness is satisfiable on a Boolean carrier, trivially and against its intent. Let $\Omega_7 = \{\pm\}^3 \setminus \{(+, +, +)\}$, let $\mathcal{L} = \mathcal{P}(\Omega_7)$, and let B be the $\perp$ -closure of the three coordinate events $A_i = \{x : x_i = +\}$. All twelve pairwise intersections among $\{A_i, A_i^\perp\}$ ($i \neq j$) are nonempty, so the only additivity constraints inside B are the complement pairs, and $s_0(A_1) = s_0(A_2) = s_0(A_3) = 1$ is a two-valued state on B . But $A_1 \cap A_2 \cap A_3 = \emptyset$, and on a finite power set every two-valued state is Dirac, so s_0 extends to no σ -additive state — on a Boolean σ -algebra. The pattern is Specker’s parable (pairwise but not jointly) embedded as a sub-orthoposet, and locality of coherence cannot distinguish distributive from non-distributive carriers. The phenomenon of interest is a failure *at σ* : finite additivity achievable globally, countable additivity not. Definition 1.1 makes this the definition, and Proposition 2.1 shows the Boolean baseline is then restored.

The phenomenon is a failure of σ -level compactness. Wright’s theorem [23], that every two-valued state on a *finite* OML is finitely realised, forbids any *finite* witness, so the question is whether a coherent finite pattern can fail to globalise once σ -additivity is imposed. On a Boolean carrier it cannot (§2). The main result of this paper is that on a non-distributive carrier it can, without hypotheses:

Theorem 1.4 (Main; proved as Theorem 5.18). *In ZFC there exist a set Ω with $|\Omega| = \aleph_1$, a concrete σ -complete orthomodular poset $\mathcal{L} \subseteq \mathcal{P}(\Omega)$, a finite $\perp$ -closed $B \subseteq \mathcal{L}$, and a two-valued state s_0 on B such that s_0 extends to a finitely additive two-valued state on $\mathcal{L}$ but to no σ -additive one. Moreover every σ -additive two-valued state on $\mathcal{L}$ is Dirac, and $\mathcal{L}$ is non-Boolean, essentially irreducible, non-segregated, and not Polish-representable.*

Remark 1.5 (Irreducibility: not vacuous, and modulo the countable ideal). The carrier class is inhabited and irreducibility is non-trivial: the horizontal sum MO_κ is concrete, non-Boolean, and irreducible (centre $\{\emptyset, \Omega\}$) for $\kappa \geq 2$ [14], σ -complete for $\kappa = \omega$, and concreteness — exactly order-determination by two-valued states [11, 12] — does not force a non-trivial centre.

Literal irreducibility is nonetheless unavailable to any carrier with all singletons, since countable sets are then central (Lemma 5.3); and the witness’s rigidity mechanism requires singletons (Theorem 5.13), so the amendment to *essential* irreducibility — trivial centre in the quotient by the countable ideal, computed exactly in Corollary 5.12 — is forced by the method. (MO_κ itself is excluded as a witness by Lemma 4.3.)

Verification status. The construction of §5 was developed jointly with a large language model (see Acknowledgements), hand-checked by two independent adversarial passes, and subsequently *formalized in Lean 4 in full*: the main theorem is machine-checked end-to-end in the author’s repository (`formalization/QuerySystem/`, modules `SigmaEssentialAmended` and `UlamWitness*`; final statement `psiAmended_ZFC`), with zero `sorry`s and axioms exactly `propext`, `Classical.choice`, `Quot.sound` — i.e. ZFC, no additional axioms and no large cardinals. The formalization covers the Ulam matrix (Lemma 5.1), the normal form and the full disjointness table (Theorem 5.8, Lemma 5.6 including case (IV)), rigidity (Theorem 5.13), the vote state and its additivity (Theorem 5.16), and the assembly at ω_1 (Theorem 5.18); the Ω_7 example of Remark 1.3 is also machine-checked, so the necessity of the coherence clause is itself certified. The remaining trust surface is definitional fidelity (that the formal definitions transcribe Definitions 1.1–1.2), a finite reading exercise.

2 The Boolean baseline

On a Boolean σ -algebra, Definition 1.2 is never satisfied. The proof isolates the obstructions a witness must overcome at once; this is the baseline the non-distributive case must escape.

Proposition 2.1 (No witness in the Boolean case). *Let $\mathcal{L} \subseteq \mathcal{P}(\Omega)$ be a Boolean σ -algebra and $B \subseteq \mathcal{L}$ a finite $\perp$ -closed sub-orthoposet with a two-valued state s_0 that is finitely coherent on $\mathcal{L}$. Then s_0 extends to a Dirac, hence to a global σ -additive two-valued state on $\mathcal{L}$.*

Proof. Let $\mu \supseteq s_0$ be a global finitely additive two-valued state on $\mathcal{L}$ (Definition 1.1) and $B_0 \subseteq \mathcal{L}$ the finite Boolean subalgebra generated by B . The atoms of B_0 partition Ω ; finite additivity of μ over this finite partition yields a unique atom $a \in B_0$ with $\mu(a) = 1$, and a , being a nonzero element of the concrete algebra B_0 , is a nonempty subset of Ω . Every element of B_0 either contains a or is disjoint from it, so for any $\omega \in a$, $\delta_\omega \upharpoonright B_0 = \mu \upharpoonright B_0$, whence

$$K(s_0) := \bigcap \{A \in B : s_0(A) = 1\} \supseteq a \neq \emptyset$$

(we call $K(s_0)$ the *kernel* of s_0), $\delta_\omega \upharpoonright B = s_0$ by Lemma 3.1, and δ_ω is σ -additive. $\square$

The division of labour is that coherence supplies the global finitely additive μ and distributivity converts it into a point. A witness must defeat three obstructions at once:

- (a) (*coherence*) the pattern must extend finitely at all (Remark 1.3: empty-kernel patterns exist even on Boolean carriers, but there they are incoherent);
- (b) on a Boolean carrier any finitely additive global extension concentrates on a nonempty atom, a common point. A witness breaks this: the events lie in incompatible blocks with empty meet, so no Dirac extends s_0 and the finitely additive extension is genuinely non-Dirac;
- (c) away from a measurable cardinal, every σ -additive two-valued state on a Boolean σ -algebra is Dirac [21, 13], so even where (b) fails no non-Dirac σ -additive state repairs the gap.

These pull against one another: breaking (b) requires non-distributivity, which is what makes (c) escapable by admitting non-Dirac global states. A witness must break (b) while leaving no such repair available and, by (a), without breaking finite coherence. The construction of §5 is engineered around this triple bind.

3 The Localization Theorem

When a Dirac extends s_0 is elementary.

Lemma 3.1. *Let $B \subseteq \mathcal{L}$ be a finite $\perp$ -closed sub-orthoposet with two-valued state s_0 . A point evaluation δ_ω satisfies $\delta_\omega \upharpoonright B = s_0$ iff $\omega \in K(s_0) = \bigcap \{A \in B : s_0(A) = 1\}$. Consequently no Dirac extends s_0 iff $K(s_0) = \emptyset$.*

Proof. By definition,

$$\delta_\omega \upharpoonright B = s_0 \iff (\omega \in A \iff s_0(A) = 1) \text{ for all } A \in B.$$

Split B by the value of s_0 . The true constraints give $\omega \in A$ for every A with $s_0(A) = 1$, i.e. $\omega \in K(s_0)$. A false A ($s_0(A) = 0$) is handled by its complement: $\perp$ -closure gives $A^\perp \in B$, and additivity gives $s_0(A^\perp) = 1 - s_0(A) = 1$, so $A^\perp$ is true and the false constraint reduces to a true one, already subsumed in $K(s_0)$. Hence the false constraints add nothing and $\delta_\omega \upharpoonright B = s_0 \iff \omega \in K(s_0)$. $\square$

Theorem 3.2 (Localization). *Let $\mathcal{L}$ be a concrete σ -orthoposet, $B \subseteq \mathcal{L}$ a finite $\perp$ -closed sub-orthoposet, and s_0 a two-valued state on B . Then s_0 is a σ -essential contextual state iff*

- (0) s_0 is finitely coherent on $\mathcal{L}$;
- (i) $K(s_0) = \emptyset$ (no point evaluation extends s_0); and
- (ii) no non-Dirac σ -additive two-valued state on $\mathcal{L}$ extends s_0 .

Moreover clauses (0) and (ii) are jointly freely arrangeable: there is a concrete σ -orthoposet and a finitely coherent s_0 with $K(s_0) = \emptyset$.

Proof. The Dirac/non-Dirac dichotomy on global σ -additive two-valued states is exhaustive, so s_0 has no σ -additive extension iff it has no Dirac extension and no non-Dirac extension. By Lemma 3.1 the first conjunct is (i); the second is (ii); clause (0) is Definition 1.2's coherence requirement. For joint arrangeability of (0) and (ii), the Navara–Pták construction (Example 3.4) realises $K(s_0) = \emptyset$ with a global extension that is even σ -additive. $\square$

Remark 3.3 (Scope). The Dirac/non-Dirac split being exhaustive, the equivalence unfolds the definition rather than reducing it to a separate problem. Its content is that (0)+(ii) is freely arrangeable, so all weight sits in (0) $\wedge$ (ii): coherence must survive while every non-Dirac σ -additive repair is destroyed. The witness of §5 decouples the two provably (Theorem 5.8).

The Navara–Pták construction supplies clauses (0)+(ii) and shows (i) is binding.

Example 3.4 (Navara–Pták [19]). Let $\Omega = Q_0 \times Q_0$ with $Q_0 = \mathbb{Q} \cap (0, 1)$, and let $f, g: \Omega \rightarrow \mathbb{R}$ be the coordinate projections. Let $\mathcal{L} = \mathcal{L}_{f,g}$ be the least σ -class containing the inverse images of Borel sets under f and g . Set $C_f = f^{-1}(\{1/3\})$, $C_g = g^{-1}(\{1/2\})$, $C_{f+g} = (f+g)^{-1}(\{1/2\})$. Define $m(A) = 1$ iff A contains one of C_f, C_g, C_{f+g} . Navara and Pták show m is a (two-valued, σ -additive) state on $\mathcal{L}$, and that since $C_f \cap C_g \cap C_{f+g} = \emptyset$ the state m is not concentrated at any point.

Remark 3.5 (The example is not itself a witness). Let B be the $\perp$ -closure of $\{C_f, C_g, C_{f+g}\}$ and s_0 the pattern giving each value 1. Then $C_f \cap C_g \cap C_{f+g} = \emptyset$ gives $K(s_0) = \emptyset$, so clause (i) holds; but the Navara–Pták state has $m \upharpoonright B = s_0$ and is non-Dirac σ -additive, so clause (ii) fails and s_0 is not σ -essential. Clauses (0)+(i) do not suffice: the repair exists because $\mathcal{L}_{f,g}$ is generated by two compatible observables, Boolean enough to support a non-Dirac global state through the point-level gap. A witness requires a carrier on which no non-Dirac state repairs the gap while, per clause (0), a finitely additive one still must.

Stripped of lattice scaffolding, the core is a selection problem. Each block is a maximal compatible context, equivalently a countable partition Π_α of Ω into elements of $\mathcal{L}$; a two-valued state selects one cell $c_\alpha \in \Pi_\alpha$ per block, subject to coherence, σ -additivity, and non-principality.

Question 3.6 (σ -point-selection, bare form). Given a set Ω and a family $\{\Pi_\alpha\}$ of countable partitions of Ω (the blocks of a concrete σ -orthoposet, pairwise overlapping in their common refinements), does there exist a selection assigning to each Π_α one cell $c_\alpha \in \Pi_\alpha$ such that

- (a) (coherence) the selections agree on events common to several blocks;
- (b) (σ -additivity) the induced $\{0, 1\}$ -valuation commutes with countable disjoint unions;
- (c) (non-principality) there is no $\omega_0 \in \Omega$ with $c_\alpha =$ the cell of Π_α containing ω_0 for all α ;
- (d) (prescription) the selection realises a prescribed finite pattern that no single point realises?

A witness exists iff, on some admissible carrier, some prescribed pattern that is realisable by a merely finitely additive selection admits *no* selection satisfying (a)–(d) and no principal one either: the witness question is the localised *negative* instance of the selection problem, with finite-additive realisability retained. Two features of the bare form, obscured in lattice language, fix its position: its distance from a measurable cardinal, and its place within contextuality.

Remark 3.7 (The Boolean shadow is a measurable cardinal; the question is not). For a compatible block family (a single Boolean base $\mathcal{B}$ on Ω), a coherent non-principal σ -additive selection $s: \mathcal{B} \rightarrow \{0, 1\}$ has $U_s := s^{-1}(1)$ a non-principal σ -complete ultrafilter on $\mathcal{B}$, so $\exists s \implies |\Omega|$ measurable [13]. Question 3.6 imposes coherence (a) across incompatible, mutually non-distributive blocks: strictly stronger than an ultrafilter on any single block. The Boolean shadow reduces to a measurable cardinal; the non-distributive coherence is exactly what the reduction cannot cross (§4). The witness of §5 shows how far the shadow misleads: there the carrier’s rigidity is supplied by an Ulam matrix on ω_1 — the classical device for proving ω_1 *non*-measurable — and the coherence is carried by a mere ultrafilter on $\mathcal{P}(\omega_1)/\text{ctble}$, never asked to be σ -complete.

Remark 3.8 (Relation to sheaf-theoretic contextuality). Question 3.6 concerns locally coherent $\{0, 1\}$ -valuations with no global section, contextuality in the sense of Abramsky–Brandenburger [2], σ -additive form [1]. The framing is standard; what was open is the conjunction the literature does not treat together: countable additivity + a non-Hilbert (abstractly concrete) carrier + a $\{0, 1\}$ -valued selection. Theorem 5.18 settles that conjunction affirmatively.

4 The boundary map

We locate the core between an upper boundary where it is empty and lower routes that fail to settle it. Each boundary is a constraint the witness of §5 is forced to obey.

The upper boundary is the Polish-representable case. Derr and Williamson [6] prove a σ -extension criterion for coherent probabilities on pre-Dynkin systems: when the carrier is Polish-representable (that is, Ω Polish, the generated σ -class equal to the Borel σ -algebra, with inner-regular blocks), finite coherence implies σ -extendability (their Theorem D.6, via Maharam [16]). Specialised to two-valued states, this says that on a Polish-representable carrier every finitely coherent s_0 globalises: no σ -essential contextual state exists. A witness must therefore live on a carrier that is not Polish-representable — as the one below, with $|\Omega| = \aleph_1$ and rigidity incompatible with inner regularity, does (Proposition 5.19).

A Hilbert-space analogue exists, but does not transfer down to the concrete case. On the projection lattice of $B(\ell^2(\kappa))$, Blecher–Weaver [5] show a singular countably additive pure state exists iff κ is Ulam measurable, the nearest positive precedent that countable additivity on a non-distributive σ -complete OML can carry set-theoretic strength. It does not settle the question here; three obstructions point the same way.

- (1) *Wrong carrier.* For $\dim H \geq 3$ the projection lattice of $B(H)$ admits no two-valued state (Kochen–Specker [15]), so it is not concrete; the Blecher–Weaver dichotomy concerns pure states, not two-valued ones, which are absent there by Gleason [10].
- (2) *Masa transfer fails.* Carrying a state-existence result to an abelian subalgebra means factoring through a masa; for the pure two-valued case this fails: Akemann–Weaver [3] give a pure state on $B(H)$ multiplicative on no masa, and Blecher–Weaver use Marcus–Spielman–Srivastava paving for exactly this reason.
- (3) *The consumed object is structurally absent.* Blecher–Weaver extract their cardinal from a genuine ultrafilter on the diagonal masa $\ell^\infty(\kappa)$ (projections = subsets, intersection-closed). On a concrete non-Boolean σ -orthoposet, $\mathcal{F}_s := \{A \in \mathcal{L} : s(A) = 1\}$ is an ultrafilter for *no* two-valued state s , not even a filter: meet-closure fails already for δ_ω , since in any concrete MO_2 the atoms a, b have $a \cap b \neq \emptyset$ (else orthogonal) yet $a \wedge b = 0$, so $\delta_\omega(a) = \delta_\omega(b) = 1$, $\delta_\omega(a \wedge b) = 0$. The countably-complete ultrafilter an Ulam-style argument would consume is absent from the carrier: the same non-distributivity (lattice meet $\neq$ set intersection), one level down.

Remark 4.1 (Strength of the claim). Remark 3.7 and (1)–(3) assemble cited facts; they are not a non-reducibility theorem in a fixed reduction calculus. The claim is only that the two obvious bridges — “measurable cardinal” and “masa transfer from Blecher–Weaver” — do not close the question, the obstruction in each being the same non-distributivity, and the witness consumes neither object. Item (3) is load-bearing there: the absence of meet-closure is what lets three pairwise-incompatible assignments cohere finitely (Remark 6.2).

The remaining route from below, faithful representation, is barred by the same obstruction. Routing the core through a faithful σ -tribe representation (embedding $\mathcal{L}$ into a σ -algebra of sets, so two-valued states become point evaluations) is unavailable: a faithful tribe-of-sets representation forces distributivity, hence Booleanness (the “Floor” [20]). σ -Loomis–Sikorski holds under the Riesz Decomposition Property (σ -complete MV-algebras, RDP effect algebras [8]), but OMLs lack RDP (MO_2 witnesses), and the available image is only σ -epimorphic, not point-separating. The trivialising route is blocked, again non-distributively.

“Segregated” carriers admit no witness; the notion making this true is gluing-theoretic, not centre-theoretic (a centre-theoretic definition would exclude nothing, since the exemplar MO_ω has trivial centre by Remark 1.5). The statement and its proof follow.

Definition 4.2 (horizontal σ -sum). A concrete σ -orthoposet $\mathcal{L}$ is a *horizontal σ -sum* if its blocks pairwise intersect in $\{\emptyset, \Omega\}$ and no two nonzero proper elements of distinct blocks are disjoint.

Lemma 4.3 (Horizontal σ -sums do not witness). *If $\mathcal{L}$ is a horizontal σ -sum then $\Phi(\mathcal{L})$ holds (notation of §6): every finitely coherent two-valued state on a finite $\perp$ -closed $B \subseteq \mathcal{L}$ extends to a σ -additive two-valued state on $\mathcal{L}$. In particular MO_κ , in any concrete realisation of this shape, admits no σ -essential contextual state.*

Proof. By hypothesis every pairwise-disjoint family in $\mathcal{L}$, in particular every countable orthogonal family, lies in a single block, so a function on $\mathcal{L}$ is a σ -additive state iff its restriction to each block is one: any tuple of blockwise σ -additive two-valued states glues to a global one, no cross-block constraint existing. Let s_0 on B be finitely coherent with global finitely additive extension μ . Each block $\mathcal{B}_\gamma$ is a Boolean σ -algebra of sets; $\mu \upharpoonright (\mathcal{B}_\gamma \cap \langle B \rangle)$ concentrates, as in Proposition 2.1, on a nonempty atom of the finite Boolean algebra generated by $B \cap \mathcal{B}_\gamma$ inside $\mathcal{B}_\gamma$, and any Dirac of that block at a point of that atom is a blockwise σ -additive state agreeing with s_0 on $B \cap \mathcal{B}_\gamma$. Gluing the blockwise choices yields $\mu' \in \text{St}_\sigma(\mathcal{L})$ with $\mu' \upharpoonright B = s_0$ (elements of B lie in blocks, and on shared elements — only $\emptyset, \Omega$ — all choices agree). $\square$

Remark 4.4 (Segregated carriers). Call a carrier *segregated* when every countable orthogonal family lies in a single block whose σ -additive two-valued states rescue every finite trace. Lemma 4.3 shows segregated carriers satisfy Φ , so a witness must be non-segregated — but the moral is sharper than “glue the blocks tightly.” In the witness below the blocks *are* loosely glued at the level of the pattern (no cross-core orthogonality exists, Lemma 5.6(IV)); what fails is the gluing’s *input*: one block is a Boolean σ -algebra on ω_1 containing an Ulam matrix, admitting no non-Dirac σ -additive state to contribute (Theorem 5.13). Segregation kills a witness only where blockwise rescuers exist; destroying them blockwise is the remaining freedom.

5 The witness: the product Ulam carrier

Lineage. The carrier shape — ω_1 times a countable fibre, coupled through the countable ideal, with Ulam–Solovay rigidity making every σ -additive state countably carried — goes back to Binder and Navara [4] (Thm. 4; restated as [18, Thm. 17.4]), who drew the *opposite* moral: such enlargements are “almost Boolean” (lattice state space, Jauch–Piron). Their framework is σ -additive throughout; the finitely additive two-valued layer of §5.4 — the coherent Specker pattern built from an ultrafilter through the parity code — is absent there, and it is exactly that layer which turns the carrier shape into a witness. The empty-kernel triple descends from Navara–Pták [19] (after Dravecký–Šipoš [7]), used there in the opposite direction: a σ -additive non-concentrated rescuer exists, whereas here rigidity destroys every one.

Throughout, M denotes ω_1 (a set of cardinality $\aleph_1$ with a fixed well-order of type ω_1), $F = \{1, 2, 3, 4\}$ is a four-element *fibre* set, and $\Omega = M \times F$. For $E \subseteq \Omega$ and $f \in F$ the *fibre trace* is $E_f = \{x \in M : (x, f) \in E\}$. For $X, Y \subseteq M$ write $X \approx Y$ when $X \Delta Y$ is countable, and $X^{(0)} = X$, $X^{(1)} = M \setminus X$. Let

$$E_4 = \{0000, 1111\} \cup \{\text{the six weight-2 vectors}\} \subseteq \mathbb{F}_2^4$$

be the even-weight code, and $N = \{\kappa \in E_4 : \kappa_4 = 0\} = \{0000, 1100, 1010, 0110\}$ the *normalised* patterns, one from each complement pair $\{\kappa, \kappa \Delta 1111\}$. The *cores* are

$$A_1 = M \times \{1, 2\}, \quad A_2 = M \times \{1, 3\}, \quad A_3 = M \times \{2, 3\},$$

with trace patterns 1100, 1010, 0110 in the coordinate order (1, 2, 3, 4). Then $A_1 \cap A_2 = M \times \{1\}$, $A_1 \cap A_3 = M \times \{2\}$, $A_2 \cap A_3 = M \times \{3\}$, $A_1 \cap A_2 \cap A_3 = \emptyset$, and $\Omega \setminus (A_1 \cup A_2 \cup A_3) = M \times \{4\}$: the four fibres are exactly the four inhabited Boolean regions of the triple, the three “solo” regions being empty. These are set identities; none of the four fibres is an element of the carrier (Corollary 5.9). This is the incidence geometry of Specker’s parable and of Example 3.4, realised with every inhabited region uncountable.

5.1 The carrier

Lemma 5.1 (Ulam matrix). *There is a family $\{C_{\alpha,n} : \alpha < \omega_1, n < \omega\}$ of subsets of M such that:*

- (1) (*row disjointness*) $C_{\alpha,n} \cap C_{\alpha,m} = \emptyset$ for $n \neq m$;
- (2) (*row covering*) $M \setminus \bigcup_{n < \omega} C_{\alpha,n} = \{\beta : \beta \leq \alpha\}$, a countable set;
- (3) (*column disjointness, exact*) $C_{\alpha,n} \cap C_{\beta,n} = \emptyset$ for $\alpha \neq \beta$.

Proof. For each $\beta < \omega_1$ fix an injection $g_\beta : \{\gamma : \gamma < \beta\} \rightarrow \omega$ (initial segments are countable; simultaneous choice by AC), and put $C_{\alpha,n} = \{\beta : \alpha < \beta < \omega_1, g_\beta(\alpha) = n\}$ [22]. (1): $g_\beta(\alpha)$ is a single value. (2): every $\beta > \alpha$ lies in $C_{\alpha,g_\beta(\alpha)}$. (3): $\beta \in C_{\alpha,n} \cap C_{\alpha',n}$ with $\alpha \neq \alpha'$ would give $g_\beta(\alpha) = n = g_\beta(\alpha')$, contradicting injectivity. $\square$

Definition 5.2 (the carrier). Let the *cells* be $D_{\alpha,n} := C_{\alpha,n} \times F \subseteq \Omega$, and let $\mathcal{L}$ be the least σ -class on Ω containing every singleton $\{\omega\}$, every cell $D_{\alpha,n}$, and the cores A_1, A_2, A_3 .

Rows ignore the fibre coordinate entirely; cores ignore the M coordinate entirely. This separation is the design principle: the Ulam machinery and the contextual geometry are prevented from interacting except through countable sets, and Theorem 5.8 is the proof that the σ -class operations cannot manufacture an interaction.

Lemma 5.3 (σ -class basics). *In any σ -class $\mathcal{L}$ containing all singletons:*

- (1) (relative complement) $X \subseteq Y, X, Y \in \mathcal{L} \implies Y \setminus X \in \mathcal{L}$;
- (2) every countable $N \subseteq \Omega$ is in $\mathcal{L}$;
- (3) (countable perturbation) $X \in \mathcal{L}, N$ countable $\implies X \cup N, X \setminus N \in \mathcal{L}$;
- (4) every two-valued state on $\mathcal{L}$ is monotone;
- (5) every countable set is compatible with every element of $\mathcal{L}$.

Proof. (1) X and $Y^\perp$ are disjoint, so $X \cup Y^\perp \in \mathcal{L}$; its complement is $Y \setminus X$. (2) A countable disjoint union of singletons. (3) $X \cup N = X \sqcup (N \setminus X)$; and $X \setminus N = (X^\perp \cup N)^\perp$ with $X^\perp \cup N \in \mathcal{L}$ by the previous sentence. (4) By (1), $s(Y) = s(X) + s(Y \setminus X) \geq s(X)$ for $X \subseteq Y$. (5) For countable N and $X \in \mathcal{L}$: $X = (X \setminus N) \sqcup (X \cap N)$ and $N = (N \setminus X) \sqcup (X \cap N)$ with all parts in $\mathcal{L}$ by (2)–(3), a joint orthogonal decomposition. $\square$

5.2 The trace invariant

Definition 5.4 (the invariant). Say $E \subseteq \Omega$ satisfies the invariant, written $E \in \tilde{P}$, if there exist $\xi \subseteq M$ and $\kappa \in E_4$ with

$$E_f \approx \xi^{(\kappa_f)} \quad (f = 1, 2, 3, 4);$$

call (ξ, κ) a *representation* of E . Intrinsically: $E \in \tilde{P}$ iff all pairwise symmetric differences $E_f \Delta E_g$ are countable or co-countable and the induced parity vector is even.

Lemma 5.5 (representation rigidity). *If (ξ, κ) and (η, λ) both represent E with $\kappa, \lambda \in E_4$, then either $\lambda = \kappa$ and $\eta \approx \xi$, or $\lambda = \kappa \Delta 1111$ and $\eta \approx \xi^\perp$. Consequently:*

- (1) each $E \in \tilde{P}$ has a unique representation with $\kappa \in N$, up to $\approx$ -change of ξ ;
- (2) E is countable iff its normalised representation is $(\xi, 0000)$ with $\xi \approx \emptyset$;
- (3) if $\kappa \in N \setminus \{0000\}$ (a weight-2 element) then E is uncountable: some trace is $\approx \xi$ and some $\approx \xi^\perp$, and these cannot both be countable.

Proof. If $\kappa_f = \lambda_f$ for some f , then $\xi^{(\kappa_f)} \approx \eta^{(\kappa_f)}$ gives $\xi \approx \eta$; substituting into the remaining coordinates forces $\kappa_g = \lambda_g$ for all g , since $\xi \approx \xi^\perp$ is impossible, M being uncountable. Otherwise $\kappa_f \neq \lambda_f$ for all f , i.e. $\lambda = \kappa \Delta 1111$, and any coordinate gives $\eta \approx \xi^\perp$. Exactly one of $\kappa, \kappa \Delta 1111$ has fourth coordinate 0, giving (1); (2), (3) are immediate. $\square$

Lemma 5.6 (disjointness table). *Let $E, G \in \tilde{P}$ have normalised representations $(\xi, \kappa), (\eta, \lambda)$, $\kappa, \lambda \in N$, and suppose $E \cap G = \emptyset$ (only $E_f \cap G_f$ countable, for each f , is used). Coordinatewise, disjointness mod countable of $\xi^{(\kappa_f)}$ and $\eta^{(\lambda_f)}$ reads*

$$(\kappa_f, \lambda_f) = (0, 0): \xi \cap \eta \approx \emptyset; \quad (0, 1): \xi \subseteq \eta \text{ mod ctble}; \quad (1, 0): \eta \subseteq \xi \text{ mod ctble}; \quad (1, 1): \xi \cup \eta \approx M.$$

The ten unordered pairs (κ, λ) resolve as:

Case	(κ, λ)	Forced relations	Conclusion
(I)	(0000, 0000)	$\xi \cap \eta \approx \emptyset$	$E \sqcup G \in \tilde{P}$, rep. $(\xi \cup \eta, 0000)$
(II)	(κ, κ) , κ weight-2	$\xi \cap \eta \approx \emptyset$ and $\xi \cup \eta \approx M$	$\eta \approx \xi^\perp$; $E \sqcup G$ co-ctble, rep. $(M, 0000)$
(III)	$(0000, \lambda)$, λ weight-2	$\xi \subseteq \eta$, $\xi \cap \eta \approx \emptyset$ mod ctble	$\xi \approx \emptyset$: E countable; rep. of $E \sqcup G$ is (η, λ)
(IV)	$\kappa \neq \lambda$, both weight-2	$\xi \approx \eta$, $\xi \cap \eta \approx \emptyset$, $\xi \cup \eta \approx M$	impossible ($\emptyset \approx M$)

Proof. The coordinate translations are immediate; (I)–(III) are read off. For (IV): any two distinct elements of $N \setminus \{0000\}$ agree in exactly one of the first three coordinates and both vanish in the fourth, so the multiset of coordinate pairs (κ_f, λ_f) is exactly $\{(1, 1), (1, 0), (0, 1), (0, 0)\}$. Then $(1, 0)$ and $(0, 1)$ give $\xi \approx \eta$; $(0, 0)$ gives $\xi \cap \eta \approx \emptyset$, hence $\xi \approx \emptyset$; $(1, 1)$ gives $\xi \cup \eta \approx M$, hence $\emptyset \approx M$ — false, M being uncountable. $\square$

Row (IV) is the theorem in miniature: no element of $\mathcal{L}$ carrying one core pattern is ever exactly disjoint from an element carrying a different core pattern, beyond countable degeneracy. This is what makes the three simultaneous assignments $s_0(A_i) = 1$ immune to additivity constraints; the parity-triangle certificate that dooms tighter designs (three exact events with $a + b = a + c = b + c = 1$ over $\{0, 1\}$, summing to the odd total 3) cannot form.

Lemma 5.7 (family trichotomy). *Let $\{E_i\}_{i < \omega} \subseteq \tilde{P}$ be pairwise exactly disjoint. Then exactly one of:*

- (1) *every E_i of uncountable class is diagonal ($\kappa_i = 0000$, $\xi_i \not\approx \emptyset$), pairwise $\xi_i \cap \xi_j \approx \emptyset$;*
- (2) *exactly one E_{i_0} is weight-2; all other E_i are countable;*
- (3) *exactly two E_{i_0}, E_{i_1} are weight-2, in the same coset with $\xi_{i_1} \approx \xi_{i_0}^\perp$; all other E_i are countable.*

Proof. By Lemma 5.6(IV) all weight-2 members share one coset. By (II) two same-coset weight-2 members are complementary mod countable; a third in that coset would be $\approx$ -complementary to both, giving $\xi \approx \xi^\perp$ — impossible. So there are at most two weight-2 members. If at least one exists, any diagonal member is disjoint from it, hence countable by (III); if two (complementary) weight-2 members exist, any further member has each trace class below the complement of their join, which is $\approx \emptyset$, hence is countable. $\square$

Theorem 5.8 (normal form). $\mathcal{L} \subseteq \tilde{P}$: *every element of the carrier satisfies the invariant.*

Proof. $\tilde{P}$ contains the generators: singletons and $\emptyset$ are $(\emptyset, 0000)$; Ω is $(M, 0000)$; cells are $(C_{\alpha, n}, 0000)$; the cores are $(\emptyset, 1100)$, $(\emptyset, 1010)$, $(\emptyset, 0110)$. It is complement-closed: $(E^\perp)_f = (E_f)^\perp$, so $(\xi, \kappa \Delta 1111)$ represents $E^\perp$. It is closed under countable pairwise exactly-disjoint unions, by Lemma 5.7: in case (1) the union is represented by $(\bigcup_i \xi_i^*, 0000)$, where $\xi_i^* := \xi_i \setminus \bigcup_{j < i} (\xi_i \cap \xi_j)$ trims the pairwise-countable overlaps — each $\bigcup_{j < i} (\xi_i \cap \xi_j)$ is countable, so $\xi_i^* \approx \xi_i$, the ξ_i^* are exactly disjoint, and coordinatewise the union's trace differs from $\bigcup_i \xi_i^*$ by a countable union of countable sets; in case (2) by $(\xi_{i_0}, \kappa_{i_0})$, the countable members perturbing each trace countably; in case (3) by $(M, 0000)$. Hence $\{E : E \in \tilde{P}\}$ is a σ -class containing the generators, so it contains $\mathcal{L}$. $\square$

Corollary 5.9 (the missing meet region). *No element of $\mathcal{L}$ is $\approx$ -equal (fibrewise) to a fibre $M \times \{f\}$, $f \in F$. In particular the pairwise core meets $A_1 \cap A_2 = M \times \{1\}$, $A_1 \cap A_3 = M \times \{2\}$, $A_2 \cap A_3 = M \times \{3\}$, and $M \times \{4\} = A_i^\perp \cap A_j^\perp$ ($i \neq j$), are not in $\mathcal{L}$.*

Proof. The trace classes of $M \times \{f\}$ are M at coordinate f and $\emptyset$ elsewhere, mod countable. In any representation, the coordinates take at most the two values $\xi, \xi^\perp$ mod countable; here they take the classes of M and $\emptyset$, so $\xi \approx \emptyset$ or $\xi \approx M$, and the pattern is κ or $\kappa \Delta 1111$ for some $\kappa \in E_4$. But a unit vector and its complement have odd weight. $\square$

Corollary 5.10 (stripping bar). *If $E \in \mathcal{L}$ has E_3, E_4 countable (i.e. $E \subseteq A_1$ mod countable), then E is countable or $E \approx A_1$ fibrewise. Likewise for A_2, A_3 and the core complements. If $E \subseteq A_1 \cap A_2$ exactly, E is countable.*

Proof. Take the normalised (ξ, κ) ; throughout, coordinate f carries $\xi^{(\kappa_f)}$ (Definition 5.4). If $\kappa = 0000$: coordinates 3, 4 carry ξ , so $\xi \approx \emptyset$ and E is countable. If $\kappa = 1100$: coordinates 3, 4 carry ξ ($\kappa_3 = \kappa_4 = 0$), so $\xi \approx \emptyset$; coordinates 1, 2 then carry $\xi^\perp \approx M$, so $E \approx A_1$. If $\kappa = 1010$ or 0110 : one of coordinates 3, 4 carries ξ and the other $\xi^\perp$, forcing $\xi \approx \emptyset$ and $\xi \approx M$ simultaneously — impossible. For the last claim, $E \subseteq M \times \{1\}$ makes coordinates 2, 3, 4 countable, and the same analysis kills all three weight-2 cosets, leaving $\xi \approx \emptyset$. $\square$

Corollary 5.11 (incompatibility; not a lattice). *A_1 and A_2 are incompatible in $\mathcal{L}$; hence $\mathcal{L}$ is non-Boolean. Moreover $A_1 \wedge A_2$ does not exist: the lower bounds of $\{A_1, A_2\}$ in $\mathcal{L}$ are exactly the countable subsets of $M \times \{1\}$, among which there is no maximum. Thus $\mathcal{L}$ is a σ -complete orthomodular poset and provably not a lattice.*

Proof. Compatibility in a concrete orthomodular poset means $A_1 = A' \sqcup D$, $A_2 = A'' \sqcup D$ with $A', A'', D \in \mathcal{L}$ pairwise disjoint; then $D = A_1 \cap A_2$ (expand $A_1 \cap A_2$ using $A' \cap A'' = \emptyset$), contradicting Corollary 5.9. The lower-bound computation is the last clause of Corollary 5.10. $\square$

Corollary 5.12 (centre). *$Z(\mathcal{L}) = \{E \in \mathcal{L} : E \text{ countable or co-countable}\}$: the quotient of $\mathcal{L}$ by the countable ideal has trivial centre, so $\mathcal{L}$ is essentially irreducible.*

Proof. ($\supseteq$) Lemma 5.3(5) and complement-closure of the centre. ($\subseteq$) Let E be central. Compatibility with A_1 puts $E \cap A_1 \in \mathcal{L}$ (as in Corollary 5.11, the common part of a compatible decomposition is the intersection), whose trace quadruple is $(E_1, E_2, \emptyset, \emptyset)$ with the last two coordinates exactly empty. Membership in $\tilde{P}$ (Theorem 5.8) then forces, by the analysis of Corollary 5.10, $[E_1] = [E_2] \in \{0, 1\}$ as classes mod countable. Compatibility with A_2 gives $[E_1] = [E_3]$, with A_3 gives $[E_2] = [E_3]$, all in $\{0, 1\}$; so the first three trace classes are a common constant t . Membership of E itself in $\tilde{P}$ forces the fourth class to equal t : with three coordinates constant t , the pairwise-difference and parity conditions of Definition 5.4 require the fourth difference to vanish. Thus E is countable ($t = 0$) or co-countable ($t = 1$).

For the quotient claim, let $[E]$ be central in $\mathcal{L}/\text{ctble}$. Compatibility of $[E]$ with $[A_1]$ in the quotient supplies $D \in \mathcal{L}$ with $D \approx E \cap A_1$ (the quotient decomposition, as in Corollary 5.11, computed mod countable). Then $D \in \tilde{P}$ has coordinates 3, 4 countable, and the analysis of Corollary 5.10 — which only ever uses trace *classes* — runs unchanged mod countable, giving $[E_1] = [E_2] \in \{0, 1\}$. Compatibility with $[A_2], [A_3]$ give $[E_1] = [E_3]$ and $[E_2] = [E_3]$, and membership of E in $\tilde{P}$ forces the fourth class exactly as above. Hence E is countable or co-countable, i.e. $[E] \in \{0, 1\}$: the quotient centre is trivial. $\square$

5.3 Rigidity

Theorem 5.13 (rigidity). *Every σ -additive two-valued state on $\mathcal{L}$ is Dirac.*

Proof. Let s be σ -additive and two-valued. If $s(\{\omega\}) = 1$ for some ω , monotonicity (Lemma 5.3(4)) gives $s(E) = 1$ whenever $\omega \in E$, and $\omega \notin E$ gives $s(E^\perp) = 1$, so $s(E) = 0$: $s = \delta_\omega$. Otherwise s vanishes on all singletons, hence, by σ -additivity, on all countable sets. Fix a row α : the cells $\{D_{\alpha, n}\}_{n < \omega}$ are pairwise exactly disjoint (Lemma 5.1(1)), their union U_α lies in $\mathcal{L}$ (σ -class closure), and $\Omega \setminus U_\alpha$ is countable (Lemma 5.1(2)), so $s(U_\alpha) = 1$ and σ -additivity over the row yields exactly one $n(\alpha) < \omega$ with $s(D_{\alpha, n(\alpha)}) = 1$. The map $\alpha \mapsto n(\alpha)$ sends ω_1 into ω , so some value n^* has uncountable fibre; fix $\alpha \neq \beta$ with $n(\alpha) = n(\beta) = n^*$. By column disjointness (Lemma 5.1(3)), $D_{\alpha, n^*} \cap D_{\beta, n^*} = \emptyset$ exactly; their union is in $\mathcal{L}$, and additivity gives s the value 2 there — contradiction. So the second case is void. $\square$

The proof consumes only singletons, cells, rows, columns, and σ -class closure — nothing about cores or the invariant. Rigidity is therefore robust to any enlargement of the generator set preserving these items. Note the inversion of Remark 3.7: the Ulam matrix, classically the device by which ω_1 fails to be measurable, here works *for* the witness — clause (ii) of Theorem 3.2 is achieved outright, in ZFC, by importing the matrix into the carrier as generators.

5.4 The state

Definition 5.14. Fix an ultrafilter $\mathcal{U}$ on M extending the co-countable filter (the filter is proper, M being uncountable), and write $u(\xi) \in \{0, 1\}$ for its characteristic function. Then u vanishes on countable sets, is constant on $\approx$ -classes, satisfies $u(\xi^\perp) = 1 - u(\xi)$, and is additive on pairs with $\xi \cap \eta \approx \emptyset$. For $E \in \mathcal{L}$ with normalised representation (ξ, κ) define the *vote*

$$v(E) := \kappa \Delta (u(\xi) \cdot 1111) \in E_4,$$

and, with $\hat{S} := \{1111, 1100, 1010, 0110\}$ (one member from each complement pair of E_4),

$$m(E) := \mathbf{1}[v(E) \in \hat{S}].$$

Lemma 5.15 (well-definedness). *v , hence m , is independent of the representation. Moreover $v(E^\perp) = v(E) \Delta 1111$, so $m(E^\perp) = 1 - m(E)$; $m(\Omega) = 1$; m vanishes on countable sets; and $m(A_1) = m(A_2) = m(A_3) = 1$.*

Proof. By Lemma 5.5 the normalised representation is unique up to $\approx$ -change of ξ , on which u is constant; computing from the non-normalised partner $(\xi^\perp, \kappa \Delta 1111)$ gives $\kappa \Delta 1111 \Delta (u(\xi^\perp) \cdot 1111) = \kappa \Delta ((1 + u(\xi^\perp)) \cdot 1111) = \kappa \Delta (u(\xi) \cdot 1111)$. Complement: representation $(\xi, \kappa \Delta 1111)$, vote $v(E) \Delta 1111$; since $\hat{S}$ selects one member per complement pair, m flips. Ω : $(M, 0000)$, $u(M) = 1$, vote $1111 \in \hat{S}$. Countable E : $(\emptyset, 0000)$, vote $0000 \notin \hat{S}$. Cores: $\xi = \emptyset$, $u = 0$, votes $1100, 1010, 0110 \in \hat{S}$. $\square$

Theorem 5.16 (m is a two-valued state). *For every finite pairwise-disjoint family $\{E_i\} \subseteq \mathcal{L}$ (whose union is automatically in $\mathcal{L}$), $m(\bigsqcup_i E_i) = \sum_i m(E_i)$. Hence m is a finitely additive two-valued state on $\mathcal{L}$ extending the pattern $s_0(A_i) = 1$, which is therefore finitely coherent on $\mathcal{L}$.*

Proof. Pairs suffice (finite families follow by induction, partial unions lying in $\mathcal{L}$). Let $E \cap G = \emptyset$, $H = E \sqcup G$, and run Lemma 5.6 on normalised representations. (I): $m(E) = u(\xi)$ and $m(G) = u(\eta)$ (votes $u \cdot 1111$; $0000 \notin \hat{S}$, $1111 \in \hat{S}$), and H is represented by $(\xi \cup \eta, 0000)$ up to countable trimming, so $m(H) = u(\xi \cup \eta) = u(\xi) + u(\eta)$ by additivity of u on $\approx$ -disjoint pairs. (II): $\eta \approx \xi^\perp$ gives $v(G) = v(E) \Delta 1111$, so exactly one of $m(E), m(G)$ is 1; and $H \approx \Omega$ has vote 1111 , $m(H) = 1$. (III): the countable member has $m = 0$ and H shares the uncountable member's normalised representation (countable perturbation), so $m(H) = m(G) + 0$. (IV) cannot occur. Finally, s_0 is a two-valued state on the $\perp$ -closure $B = \{\emptyset, \Omega, A_i, A_i^\perp\}$ independently of m : each pairwise intersection among $\{A_i, A_i^\perp\}_{i \leq 3}$ ($i \neq j$) is a full fibre $M \times \{f\}$, nonempty, so the only additivity constraints inside B are the complement pairs; and $m \upharpoonright B = s_0$ by Lemma 5.15. $\square$

Remark 5.17 (the σ -failure of m is self-witnessing). Suppose u selected a cell in every row: $\forall \alpha \exists n(\alpha)$ with $u(C_{\alpha, n(\alpha)}) = 1$. Pigeonholing $\alpha \mapsto n(\alpha)$ as in Theorem 5.13 produces two disjoint sets of u -value 1 — impossible for an ultrafilter. So some row α_0 has $u(C_{\alpha_0, n}) = 0$ for all n : there $m(D_{\alpha_0, n}) = 0$ for every n while m of the (co-countable) row union is 1. Every ultrafilter fails σ -additivity on some row, by the same pigeonhole that drives rigidity; no genericity demands on the matrix are needed anywhere in the construction.

5.5 The main theorem

Theorem 5.18 (ZFC witness). *Let $\mathcal{L}$, B , s_0 be as above. Then:*

- (1) s_0 is finitely coherent on $\mathcal{L}$: m extends it (Theorem 5.16);
- (2) no σ -additive two-valued state on $\mathcal{L}$ extends s_0 : by Theorem 5.13 any such extension is some δ_ω , and by Lemma 3.1 this requires $\omega \in K(s_0) = A_1 \cap A_2 \cap A_3 = \emptyset$;
- (3) hence s_0 is a σ -essential contextual state on $\mathcal{L}$ (Definition 1.2), and the target sentence Ψ of §6 is a theorem of ZFC.

Proposition 5.19 (admissibility). *The carrier $\mathcal{L}$ is concrete and σ -complete (by construction); non-Boolean, with the incompatibility of A_1, A_2 robust under countable perturbation (Corollaries 5.9, 5.11); essentially irreducible, with centre exactly the countable/co-countable sets (Corollary 5.12), which m annihilates; non-segregated in the sense of Remark 4.4, as a corollary of the witness property (segregated carriers satisfy Φ , Lemma 4.3 and Remark 4.4; Theorem 5.18); and not Polish-representable, as a corollary of the witness property via Derr–Williamson (§4). It is not a lattice (Corollary 5.11).*

6 Discussion

Write $\text{St}(B)$ for the two-valued states on a sub-orthoposet B , $\text{St}_\sigma(\mathcal{L})$ for the global σ -additive two-valued states on $\mathcal{L}$, $\text{St}_{fa}(\mathcal{L})$ for the global finitely additive ones, $\text{Fin}_\perp(\mathcal{L})$ for the finite $\perp$ -closed sub-orthoposets, and Adm for the admissible carriers: concrete, σ -complete, non-Boolean, essentially irreducible, non-segregated, non-Polish-representable (the last two being consequences of witnesshood, by Lemma 4.3 and Derr–Williamson, not extra demands). The σ -additive lifting property and its negation are

$$\begin{aligned}\Phi(\mathcal{L}) &::= \forall B \in \text{Fin}_\perp(\mathcal{L}) \forall s \in \text{St}(B) \left[\exists \mu \in \text{St}_{fa}(\mathcal{L}) \mu \upharpoonright B = s \implies \exists \nu \in \text{St}_\sigma(\mathcal{L}) \nu \upharpoonright B = s \right], \\ \Psi &::= \exists \mathcal{L} \in \text{Adm} \neg \Phi(\mathcal{L}).\end{aligned}$$

Theorem 5.18 proves Ψ in ZFC. Proposition 2.1 is the statement that Boolean carriers satisfy Φ ; Lemma 4.3 that horizontal σ -sums do; Derr–Williamson that Polish-representable carriers do. The witness therefore inhabits precisely the cell the boundary map of §4 isolated, and each constraint of that map is visible in its anatomy.

Remark 6.1 (consistency strength, resolved). Two routes would assign Ψ large-cardinal strength, and both fail (§4): a measurable-cardinal lower bound, because reduce-to-Dirac is a σ -algebra theorem unavailable on σ -classes; and a Blecher–Weaver upper bound, for want of a masa. Theorem 5.18 settles Ψ deflationarily: a theorem of ZFC, no large-cardinal strength. The two failures survive as the explanation. Reduce-to-Dirac is embraced, not escaped: the carrier imports exactly the σ -algebra fragment Ulam needs (rows, columns, singletons), and Theorem 5.8 certifies it does not Booleanise the carrier. The σ -complete ultrafilter Blecher–Weaver consume is never needed: coherence rides on an arbitrary ultrafilter on $\mathcal{P}(\omega_1)/\text{ctble}$, a BPI-level object of which σ -additivity is never asked (Remark 5.17 shows it necessarily fails).

Remark 6.2 (what witness technology can and cannot import). The construction divides labour between two sources, and the division is forced. *Incompatibility* cannot be imported from a Boolean-ambient combinatorial skeleton: in such an ambient a maximal coherent $\{0, 1\}$ -assignment is an ultrafilter, hence meet-closed, and the pattern globalises. The witness’s incompatibility is instead native, carried by the fibre parity code, and the same absent meet-closure (§4(3)) is what lets the three core assignments cohere — Lemma 5.6(IV) says the linking additivity constraints do not exist in $\mathcal{L}$. What *can* be imported is rigidity, the destruction of

non-Dirac σ -states; Theorem 5.8 is the proof that the imported engine and the native geometry never interact.

Remark 6.3 (a quotient with no σ -additive states at all). In the quotient of $\mathcal{L}$ by the countable ideal, Dirac states die (they concentrate on singletons) and non-Dirac σ -additive states are barred by Theorem 5.13. The quotient is thus a σ -complete non-Boolean orthomodular poset carrying finitely additive two-valued states (the induced $\bar{m}$) but *no* σ -additive two-valued state whatsoever. This artifact is graded modestly: abstract logics with finitely additive but not countably additive states are established art — Navara [17] (a σ -orthocomplete lattice logic, via stateless Greechie components, not set-representable) and Dvurečenskij–Neubrunn–Pulmannová [9] (the splitting-subspace logic of an incomplete inner-product space). The quotient’s residual interest is the conjunction it inherits from the witness: *two-valued* states, σ -completeness, and provenance as a quotient of a *concrete* carrier. It is not itself concrete ($\text{St}_\sigma = \emptyset$, while concrete logics always carry Diracs).

What remains open

Question 6.4 (the lattice-restricted form).

$$\Psi_{\text{lat}} := \exists \mathcal{L} \in \text{Adm} [\mathcal{L} \text{ is a lattice} \wedge \neg \Phi(\mathcal{L})].$$

Theorem 5.18 proves Ψ in ZFC; Ψ_{lat} is open. We conjecture that latticehood forces Φ within Adm , so that Ψ_{lat} fails. The present witness gives limited evidence in that direction: it is concrete but nonlattice, with absent meet-closure permitting its three core assignments to cohere (Remark 6.2). Navara’s σ -orthocomplete lattice example is not set-representable (Remark 6.3), so neither example is both concrete and a lattice. Boolean carriers and horizontal σ -sums are lattices satisfying Φ (Proposition 2.1; Lemma 4.3), but do not decide Ψ_{lat} .

Question 6.5 (cardinality). The witness has $|\Omega| = \aleph_1$. Wright’s theorem bars finite witnesses; can a countable Ω carry one (necessarily without all singletons, or with a different rigidity mechanism)? Is there a ZFC lower bound beyond “uncountable”?

Question 6.6 (characterising Φ). Boolean carriers, Polish-representable carriers, and horizontal σ -sums satisfy Φ ; the product Ulam carrier does not. Is there a structural characterisation of Φ for concrete σ -orthoposets — a gluing-theoretic or exhaustion-theoretic condition separating the two classes, with Theorem 5.18 as the canonical failure?

Acknowledgements. The author thanks the anonymous referees (forthcoming). The construction of §5 was developed jointly with a large language model (Anthropic Claude) in interactive sessions, which also assisted with literature search, verification of cited statements against primary sources, and exposition; OpenAI ChatGPT assisted with earlier literature work. All mathematical claims are being independently verified by the author, including a Lean 4 formalization in progress; see the Verification status paragraph in §1.

References

- [1] Samson Abramsky and Rui Soares Barbosa. The sheaf-theoretic structure of definite causality. *Communications in Mathematical Physics*, 391, 2022. σ -additive sheaf-theoretic contextuality; see also arXiv:1905.08267.
- [2] Samson Abramsky and Adam Brandenburger. The sheaf-theoretic structure of non-locality and contextuality. *New Journal of Physics*, 13:113036, 2011.

- [3] Charles A. Akemann and Nik Weaver. $\mathcal{B}(H)$ has a pure state that is not multiplicative on any masa. *Proceedings of the National Academy of Sciences*, 105(14):5313–5314, 2008.
- [4] Josef Binder and Mirko Navara. Quantum logics with lattice state spaces. *Proc. Amer. Math. Soc.*, 100:688–693, 1987.
- [5] David P. Blecher and Nik Weaver. Quantum measurable cardinals. *Journal of Functional Analysis*, 273(5):1870–1890, 2017.
- [6] Rabanus Derr and Robert C. Williamson. Systems of precision: coherent probabilities on pre-Dynkin-systems and coherent previsions on linear subspaces. *arXiv preprint arXiv:2302.03522*, 2023.
- [7] Jozef Dravecký and Ján Šipoš. On the additivity of Gudder integral. *Math. Slovaca*, 30:299–303, 1980.
- [8] Anatolij Dvurečenskij. Loomis–sikorski theorem for σ -complete MV-algebras and ℓ -groups. *Journal of the Australian Mathematical Society*, 68(2):261–277, 2000.
- [9] Anatolij Dvurečenskij, Tibor Neubrunn, and Sylvia Pulmannová. Regular states and countable additivity on quantum logics. *Proc. Amer. Math. Soc.*, 114:931–938, 1992.
- [10] Andrew M. Gleason. Measures on the closed subspaces of a Hilbert space. *Journal of Mathematics and Mechanics*, 6(6):885–893, 1957.
- [11] Stanley P. Gudder. Quantum probability spaces. *Proceedings of the American Mathematical Society*, 21:296–302, 1969.
- [12] John Harding. Remarks on concrete orthomodular lattices. *International Journal of Theoretical Physics*, 43(10):2149–2168, 2004.
- [13] Thomas Jech. *Set Theory*. Springer, 3rd millennium edition, 2003.
- [14] Gudrun Kalmbach. *Orthomodular Lattices*. Academic Press, 1983.
- [15] Simon Kochen and Ernst P. Specker. The problem of hidden variables in quantum mechanics. *Journal of Mathematics and Mechanics*, 17(1):59–87, 1967.
- [16] Dorothy Maharam. Consistent extensions of linear functionals and of probability measures. In *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability, Vol. II: Probability Theory*, pages 127–147. University of California Press, Berkeley, 1972.
- [17] Mirko Navara. Regularity and σ -additivity of states on quantum logics. *Proc. Amer. Math. Soc.*, 115:427–429, 1992.
- [18] Mirko Navara. State spaces of orthomodular structures. *Rend. Istit. Mat. Univ. Trieste*, 31, suppl. 1:143–201, 2000.
- [19] Mirko Navara and Pavel Pták. Two-valued measures on σ -classes. *Časopis pro pěstování matematiky*, 108(3):225–229, 1983.
- [20] Pavel Pták and Sylvia Pulmannová. *Orthomodular Structures as Quantum Logics*, volume 44 of *Fundamental Theories of Physics*. Kluwer, Dordrecht, 1991.
- [21] Roman Sikorski. *Boolean Algebras*. Springer, 2nd edition, 1964.
- [22] Stanisław Ulam. Zur masstheorie in der allgemeinen mengenlehre. *Fundamenta Mathematicae*, 16:140–150, 1930.

- [23] Ron Wright. The state of the pentagon: a nonclassical example. In A. R. Marlow, editor, *Mathematical Foundations of Quantum Theory*, pages 255–274. Academic Press, New York, 1978.