FATAL LATEX COMPILATION ERROR(s) To view full GitHub Actions output, click on item at top of list here: $https://github.com/pmarcum/SAUNAS_{I}II/actions/runs/11775511169/job/32796099303$ FATALERRORSINmyPaper.tex./myPaper.tex: 132: Undefined control sequence.newule2in0.7ptWARNINGSINmyPaper.texLaTeXWarning: Reference'sec: methods' on page 4 undefined on in put line 142.LaTeXWarning: Reference'sec: DIS' on page 4 undefined on input line 142. $This paper is organized as follows. The datasets and methodology pipeline is described in Sec.~\ref{Sec. 1}. The results are presented in Sec.~\ref{Sec. 2}. The discussion of the properties of$ ule2in0.7ptLa TeXWarning: Reference `tab: Observations' on page 4 undefined on input line 188. $A total of 9.92\ ksof\ observations, using the {\it VFAINT} mode, have been archived for NGC 5084 (ACIS-I, Obs. ID: 12173, PI:\ Stephen\ Murrell Market and Market an$ ule2in0.7ptLaTeXWarning: Reference `tab: Observations' on page 6 undefined on input line 197.Proposal ID: 6785, F702W and F658N, June 1996), WFC3/IR (F160W, Proposal PI: Boizelle, Benjamin, Proposal ID: 15909, and WFC3/IR (F160W, Proposal PI: Boizelle, Benjamin, Proposal ID: 15909, and WFC3/IR (F160W, Proposal PI: Boizelle, Benjamin, Proposal ID: 15909, and WFC3/IR (F160W, Proposal PI: Boizelle, Benjamin, Proposal ID: 15909, and WFC3/IR (F160W, Proposal PI: Boizelle, Benjamin, Proposal ID: 15909, and WFC3/IR (F160W, Proposal PI: Boizelle, Benjamin, Proposal ID: 15909, and WFC3/IR (F160W, Proposal PI: Boizelle, Benjamin, Proposal ID: 15909, and WFC3/IR (F160W, Proposal PI: Boizelle, Benjamin, Proposal ID: 15909, and WFC3/IR (F160W, Proposal PI: Boizelle, Benjamin, Proposal ID: 15909, and WFC3/IR (F160W, Proposal PI: Boizelle, Benjamin, Proposal ID: 15909, and WFC3/IR (F160W, Proposal PI: Boizelle, Benjamin, Benjamin, Benjamin, Benjamin, Benjamin, Benjami $29.91\pm2.12~Mpc~[6.90~arcsec~kpc^{-1},][]koribalski+2004aj128_{1}6, and assuming a Nyquist sampled PSF, the physical spatial resolution scales are also also account to the contraction of the contraction$ 12 pc (WFC3/UVIS) and \sim 36 pc (WFC3/IR), respectively. Table ?? summarizes the available observations. LaTeX Warning: Reference 'fig:NGC5084' on page 8 undefined on input line 222. Line 222 ?? in section ?? ule2in0.7ptLaTeX Warning: Reference 'fig:NGC5084' on page 8 undefined on input line 247. Line 247 — Figure ?? presents the new-processed images in the selected broad X-ray band (0.3-2.0 keV), overlayed on the optical morphology of the galaxy [large FOV optical and near-infrared gri image from Pan-STARRS,][]chambers+2016arXiv1612.05560. Additionally, Fig. ?? in Appendix?? shows the three X-ray (soft: 0.3-1.0 keV, medium: 1.0-2.0 keV, hard: 2.0 - 8.0 keV) bands. ule2in0.7ptLaTeX Warning: Reference 'fig:NGC5084' on page 9 undefined on input line 252. flux inside the 3σ contour in Fig. ??; (2) Core (R < 10"); (3) Extended emission (10" < R < 1.5', and > 3σ); and (4-7) North, West, South, and East lobes. The lobe regions are defined by dividing the extended emission in 4 quadrants (with separations at 45°, 135°, 225°, 315°), excluding the emission from the core $(R < 10^{\circ})$ with a limit at R = 1.5'. The separation between the core and extended regions R = 10" is defined based on the inspection of the X-ray morphology as the maximum radius where the X-ray emission does not show a significant elongation. The results are presented in Table ??. The fluxes take into account PSF deconvolution, and they exclude the emission from point sources identified by the catalog of point sources (XRBs, AGN core). The analysis shows that the integrated emission from the north and east lobes is in excess of a 10σ detection. The south and west are substantially dimmer but significant above the background at 6.6σ and 3.2σ . ule2in0.7ptLaTeX Warning: Reference 'fiq:NGC5084' on page 9 undefined on input line 257.

In order to verify the results from the pipeline, we study the significance of the extended emission in Appendix $\ref{appendix}$ using two different, additional methodologies widely used in the literature. First, we determine if there is an excess of emission around the bright core of the galaxy by comparing the PSF surface brightness profile with the observed profile in the original /ACIS observations, without applying Voronoi binning or PSF deconvolution. This methodology has been extensively used in the literature fabbiano+2017apj8424, fabbiano+2018apj855131, jones+2020apj891133, ma+2020apj900164, ma+2023apj94861toidentifyhotgashalosandotherextendedemissioncompone contours on Fig. $\ref{appendix}$, confirming that the X-ray emission of NGC 5084 is not caused by PSF-scattered light from the bright core and that the extended emission is significant.

$To further investigate the nature of this complex emission, we explored the available HST observations of NGC~5084 (see Table \ref{table}), presented in the latest and the property of the p$
$LaTeXWarning: Reference `fig: NGC 5084_h st' on page 12 undefined on input line 314.$
Line 314
ule2in0.7pt LaTeX Warning: Reference 'fig:ALMA' on page 12 undefined on input line 323.
Line 323 ————— The positions that do not contain emission with an associated intensity higher than the noise at a 95% probability confidence level are masked. The velocity map (moment 1) of the $CO(2-1)$ emission lines is represented in a color scale over the F702W HST/ACS image in the right panel of Fig. ??. The results show an unmistakable edge-on rotation pattern, with the north (south) edge of the disk moving towards (away from) the observer along the line of sight. The peaks of $CO(2-1)$ emission are coincident with the edges of the circumnuclear disk, where the column density is expected to be highest due to projection effects.
ule2in0.7pt
LaTeX Warning: Reference 'fig:ALMA' on page 12 undefined on input line 325. Line 325 ———————
The spectra of the locations with significant emission are combined (summed) to study the line profile and characterize the amplitude of the rotation pattern. The combined spectra is shown in the left panel of Fig.??. We characterize the rotation amplitude using the line width at the half-maximum, following a similar procedure as in $[W_{50},][]$ smith+2021mnras500 ₁ 933.Thevelocities(lowestandhighest)atwhich maximumaremeasuredthroughinterpolationofthecombinedspectra.Thisprocessisrepeatedusing N=1000MonteCarlosimulations, withe 249.1 $^{+12.8}_{-9.6}$ km s ⁻¹ ($W_{50} = 498^{+26}_{-19}$ km s ⁻¹).
$ule2in0.7pt \\ LaTeX\ Warning:\ Reference\ `tab:Radio_Lobes' on page 13 undefined on input line 353. \\ Line 353$
Interestingly, wiegert $+ 2015aj150_81$ identifies extended emission to the east and we sto f the central bright core. This emission is highlighted in level, being located at a symmetric distance from the core. The distance from the east lobe to the core is $R = 31.5^{+4.5}_{-4.5}$ arcsec or $R = 4.52^{+0.66}_{-0.65}$ kpc, while the equivalent from the west lobe is $R = 31.1^{+4.5}_{-4.6}$ arcsec ($R = 4.51^{+0.65}_{-0.67}$ kpc), being compatible at a 3σ confidence level. The position angles from the east and west lobes to the center are also compatible at a 3σ level, $PA_{\rm east-core} = 92.8^{+6.4\circ}_{-7.1\circ}$, $PA_{\rm core-west} = 92.2^{+7.1\circ}_{-7.6\circ}$. Taking into account the inclination of the circumnuclear disk ($i = 71.2^{+1.8\circ}_{-1.7}$), and assuming that the lobes are oriented along the line of the AGN radio jet axis, the deprojected distance to the core is $R = 4.8 \pm 0.70$ kpc, taking into account the uncertainties in the location of the core, radio lobes, and inclination. The results are compatible at the 1.4 and 5 GHz bands. The total luminosity of each lobe in 5 GHz is $L_{\rm 5GHz} = [1.5, 1.8] \times 10^{+19}$ W Hz^{-1} while the core is two orders of magnitude brighter with $L_{\rm 5GHz,core} = 3.60 \pm 0.01 \times 10^{+21}$ W Hz^{-1} . In 1.4 GHz, the lobes are brighter $L_{\rm 1.4GHz} = [4,6] \times 10^{+19}$ W Hz^{-1} , with $L_{\rm 5GHz,core} = 3.60 \pm 0.01 \times 10^{+21}$ W Hz^{-1} in the core. The distances to the core of each lobe and the luminosities of each component are included in Table ??. $L_{\rm 1.4GHz} = 1.000$
LaTeX Warning: Command invalid in math mode on input line 440.
Line 440 —————— using five regions: (1) the central 1.5 arcsec from the core of the galaxy; (2-5) north, south, east, and west sides of the galaxy, resp. (not including the core). The results are shown in Fig.??. The APO/DIS spectra do not reveal any signs of $H\alpha$ [$\lambda = 6562.8$ Å], $H\beta$ [$\lambda = 4861.35$ Å], or $H\gamma$ [$\lambda = 4340.47$ Å] absorption, in any of the regions analyzed, suggesting that the stellar population at the core is not dominated by a classic post-starburst stellar population.
$ule 2in 0.7pt$ $La TeX\ Warning:\ Reference\ `subsec: discussion_faded starburst' on page 18 undefined on in put line 472.$

ule 2in 0.7 pt

ule2in0.7pt

ule 2in 0.7pt

 $LaTeX\ Warning:\ Reference\ 'fig:NGC5084_hst' on page 10 undefined on in put line 293.$

 $NGC\ 5084 color maps. \\ Left\ panel: Ground-based VLT/ATLAS \textit{g-imaps}. Black contours represent the X-rayem is sion (see Fig. \ref{fig. panel}). Darkblue F702W color from Hubble WFPC2/WFC3 photometry. The circumnuclear disk is clearly visible in red. See the color bar for reference.$

These scenarios are idealized; in reality, a mixture of these cases may be responsible for the observed phenomena. In particular, the HI tilt and the observed phenomena and the observed phenomena are idealized. The observed phenomena are idealized and the observed phenomena are idealized and the observed phenomena. The observed phenomena are idealized and the observed phenomena are idealized and the observed phenomena. The observed phenomena are idealized and the observed phenomena are idealized and the observed phenomena are idealized and the observed phenomena. The observed phenomena are idealized and the observed phenomena are idealized and the observed phenomena are idealized and the observed phenomena. The observed phenomena are idealized and the observed phenomena are idealized as a manifest and the observed phenomena are idealized and the observed are idealized and the observed are idealized and the observed are idea

Summary of the formation scenarios for the vertical X-rayem is sion of NGC~5084. Cosmic time increases from left to right. Top~row: (1) Or~row in the contract of the vertical X-rayem is sion of NGC~5084. Cosmic time increases from left to right. Top~row: (1) Or~row in the vertical X-rayem is sion of NGC~5084. Cosmic time increases from left to right. Top~row: (1) Or~row in the vertical X-rayem is sion of NGC~5084. Cosmic time increases from left to right. Top~row: (1) Or~row in the vertical X-rayem is sion of NGC~5084. Cosmic time increases from left to right. Top~row in the vertical X-rayem is sion of NGC~5084. Cosmic time increases from left to right. Top~row in the vertical X-rayem is sion of NGC~5084. Cosmic time increases from left to right. Top~row in the vertical X-rayem is sion of NGC~5084. Cosmic time increases from left to right. Top~row in the vertical X-rayem is sion of NGC~5084. Cosmic time increases from left to right. Top~row in the vertical X-rayem is sion of NGC~5084. Cosmic time increase from left to right. Top~row in the vertical X-rayem is sion of NGC~5084. Cosmic time increase from the vertical X-rayem is sion of NGC~5084. Cosmic time increase from the vertical X-rayem is sion of NGC~5084. Cosmic time increase from the vertical X-rayem is sion of NGC~5084. Cosmic time increase from the vertical X-rayem is sion of NGC~5084. Cosmic time increase from the vertical X-rayem is sion of NGC~5084. Cosmic time increase from the vertical X-rayem is sion of NGC~5084. Cosmic time increase from the vertical X-rayem is sion of NGC~5084. Cosmic time increase from the vertical X-rayem is sion of NGC~5084. Cosmic time increase from the vertical X-rayem is sion of NGC~5084. Cosmic time increase from the vertical X-rayem is sion of NGC~5084. Cosmic time increase from the vertical X-rayem is sion of NGC~5084. Cosmic time increase from the vertical X-rayem is sion of NGC~5084. Cosmic time increase from the vertical X-rayem is sion of NGC~5084. Cosmic time increase from the vertical X-rayem is sion of NGC~5084

 $La TeXWarning: Reference ``subsec: discussion_f a ded starburst' on page 18 undefined on input line 480.$

La TeXWarning: Reference `fig: NGC 5084' on page 20 undefined on input line 523.

$ule2in0.7pt$ $L_{a}T_{a}YW_{a}min_{a}$, $P_{a}f_{a}min_{a}c^{2}m$
$LaTeXWarning: Reference ``subsec: results_Optical_spectra' on page 20 undefined on in put line 529.$ $Line 529$
$Secondly, the optical spectra (see Sec. \ref{Secondly}, Fig. \ref{Secondly}) do not present any Balmer absorption line stypical of post-starburs to bjects [also called E+1] and the optical spectra (see Sec. \ref{Secondly}, Fig. \ref{Secondly}) do not present any Balmer absorption line stypical of post-starburs to bjects [also called E+1] and the optical spectra (see Sec. \ref{Secondly}, Fig. \ref{Secondly}) do not present any Balmer absorption line stypical of post-starburs to bjects [also called E+1] and the optical spectra (see Sec. \ref{Secondly}, Fig. \ref{Secondly}) do not present any Balmer absorption line stypical of post-starburs to bjects [also called E+1] and the optical spectra (see Sec. \ref{Secondly}, Fig. \ref{Secondly}) do not present any Balmer absorption line stypical of post-starburs to bjects [also called E+1] and the optical spectra (see Sec. \ref{Secondly}, Fig. \ref{Secondly}) do not present any Balmer absorption line stypical of post-starburs to bject spectra (see Sec. \ref{Secondly}, Fig. \ref{Secondly}) do not present any Balmer absorption line stypical of post-starburs to bject spectra (see Sec. \ref{Secondly}, Fig. \ref{Secondly}) do not present any Balmer absorption line stypical of post-starburs to bject spectra (see Sec. \ref{Secondly}, Fig. \ref{Secondly}) do not present any Balmer absorption line stypical of post-starburs to bject spectra (see Sec. \ref{Secondly}, Fig. \ref{Secondly}) do not present any Balmer absorption line stypical of post-starburs to bject spectra (see Sec. \ref{Secondly}, Fig. \ref{Secondly}) do not present any Balmer absorption line stypical of post-starburs to bject spectra (see Sec. \ref{Secondly}, Fig. \ref{Secondly}) do not present any Balmer absorption line stypical of post-starburs to bject spectra (see Sec. \ref{Secondly}, Fig. \ref{Secondly}) do not present any Balmer absorption line stypical of post-starburs to bject spectra (see Sec. \ref{Secondly}, Fig. \ref{Secondly}) do not present any Balmer absorption line stypical of post-starburs to bject spectra (see Sec. Secon$
ule2in0.7pt
$La TeXWarning: Reference `tab: Radio_Lobes' on page 22 undefined on input line 537.$
Line 537
$Unfortunately, the systematic effects observed in radio observations along the vertical direction (Sec.~\ref{Sec. systematic}) prevent the identification of additional content of the co$
$5\ orders\ of\ magnitude\ less\ than\ the\ X-shaped\ radio\ galaxies\ from\ cheung+2009apj181_548. This classification correlates with the prediction from the prediction of the prediction of$
$shaped sources should be decaying AGN jets. \ $
ule2in0.7pt
La TeXWarning: Command invalid in math mode on input line 567.
Line 567
Another example of such phenomena is the lenticular galaxy NGC 5252. Classified as a Sey fert 1.9 galaxy argyle + 1990 mnras 243504, oster brown 5007 Å that extends for $R \sim 20$ kpc. Hubble Space Telescope and Fabry-Pérot spectrograph observations of NGC 5252 by morse+1998 apj 505120°) misalignment between the X-ray emission in soft bands $(0.3-2.0~{\rm keV})$ and the optical major axis of the galaxy. The case of NGC 5252 shows that the combination of observational evidence such as the presence of off-axis morphological components and AGN-related features is one strategy to shed light on the formation pathways of specific galaxies. $ule 2in 0.7pt$
LaTeX Warning: Reference 'sec:DIS' on page 25 undefined on input line 591. Line 591 ——————
The horizontal (in-plane) component of the X-ray emission is aligned with both the rotation axis of the newly discovered circumnuclear
$disk\ and\ the\ symmetric\ radio\ lobes,\ identified\ by\ irwin+2019aj158_21aspartofaradioAGN\ jet. Taking into account all available observationale$
shaped X-ray emission: (1) it is the remnant of a re-oriented AGN; (2) it is an outflow generated by an overpressured cocoon of hot gas powered and the remnant of the re
going star formation in the core. Combining the new observational evidence presented in this paper with previous analysis based on environment of the core of th
ule 2in 0.7 pt
$La TeXWarning: Reference ``eq: F_lim_X ray' on page 27 undefined on input line 655.$
Line 655
At $E=1$ ke V and $t=10$ ks, the equivalent exposure $^2(Mt)$ is approximately 3.8×10^6 cm 2 s 1 . For a point source detection of $\sigma=3$, with flux integration over an area associated with the PSF at the center of the ACIS detector (FWHM ~ 1.1 "), Eqn. ?? implies a limiting sensitivity of $f_{lim}=4\times10^{-15}$ erg cm $^{-2}$ s $^{-1}$, equal to the reference instrument point source sensitivity limits.
ule 2in 0.7 pt
$La TeX\ Warning:\ Reference\ `tab: Xray_flux' on page 29 undefined on input line 699.$
$La TeXWarning: Reference `fig: NGC 5084_A perture_H istograms' on page 29 undefined on input line 699.$
Line 699
In this section we assess the statistical significance of the detection of the four X-rayem is similarly assess the statistical significance of the detection of the four X-rayem is similarly assess the statistical significance of the detection of the four X-rayem is similarly assess the statistical significance of the detection of the four X-rayem is similarly as a similar of the four X-rayem is similar of the four X-rayem
$ule 2in 0.7pt$ $LaTeXWarning: Reference `tab: Xray_flux' on page 30 undefined on input line 701.$
Line 701
The analysis indicates that the surface brightness X-rayem is sion in the aperture sissignificantly higher than that of the background. The number of the background of the
$north(p=6.8\times10^{-7}),\ east\ (p=1.1\times10^{-6}),\ south\ (p=1.3\times10^{-4})\ and\ west\ (p\sim0.01).$ The fluxes integrated over the different aper-
tures are tabulated in Table ??. This result supports and verifies the findings described in Sec. ?? and Appendix ??. We conclude that
the extended X-ray emission detected by around NGC 5084 is (1) statistically significant; (2) independent of the PSF deconvolution
process; and (3) independent of the Voronoi binning methodology applied.
ule 2in 0.7 pt
LaTeX Warning: Reference 'sec:methods' on page 30 undefined on input line 711.
Line 711 —————
Detection of the X-ray extended lobes around NGC 5084 without Voronoi binning. Left: /ACIS X-ray flux (0.3–2.0 keV) map rebinned to
12.2×12.2 arcsec (25×25 pixels) for visualization purposes. PSF deconvolution was applied [see Sec. ??, and][[borlaff+2024apj967 ₁ 69Color properties is print to North (blue) West (next) South (arcsec) and Fact (company) labely Vellow containing the North (blue) West (next) South (arcsec) and Fact (company) labely Vellow containing the North (blue) West (next) South (arcsec) and Fact (company) labely Vellow containing the North (blue) West (next) South (arcsec) and Fact (company) labely Vellow containing the North (blue) West (next) South (arcsec) and Fact (company) labely Vellow containing the North (blue) West (next) South (arcsec) and Fact (company) labely Vellow containing the North (blue) West (next) South (arcsec) and Fact (company) labely Vellow containing the North (blue) West (next) South (arcsec) and Fact (company) labely Vellow containing the North (blue) West (next) South (arcsec) and Fact (company) labely Vellow containing the North (blue) West (next) South (arcsec) and Fact (company) labely Vellow containing the North (blue) West (next) South (arcsec) and Fact (company) labely Vellow containing the North (blue) West (next) South (arcsec) and Fact (company) labely Vellow containing the North (blue) West (next) South (arcsec) and Fact (company) labely Vellow containing the North (blue) West (next) South (arcsec) and Fact (company) labely Vellow containing the North (blue) West (next) South (arcsec) and Fact (company) labely South (blue)
$rayemission in the North(blue), West(red), South(green), and East(orange) lobes. Yellow contours: [2,3] \sigma (dotted, dashed respectively) de-$

Detection of the X-ray extended lobes around NGC 5084 without Voronoi binning. Left: /ACIS X-ray flux (0.3-2.0 keV) map rebinned to $12.2 \times 12.2 \text{ arcsec } (25 \times 25 \text{ pixels})$ for visualization purposes. PSF deconvolution was applied [see Sec. ??, and][]borlaff+2024apj967₁69Color rayemissionintheNorth(blue), West(red), South(green), andEast(orange)lobes.Yellow contours:[2,3] σ (dotted, dashed respectively) detection limits as estimated by (see Sec. ??, for reference. Right: Color coded histograms represent the event probability distributions for the background (grey) and the four lobe apertures. The p-values for the null hypothesis that the flux distributions in the lobe apertures are compatible with the background are represented in the legend. ule2in0.7pt

 $NGC~5084 is one of the most massive lenticular galaxies in the Local Universe [] ohls on + 2024 a j 167_31, with a total dynamical mass of approxime koribalski+2004 a j 128_16 and a mass - to - lightratio <math>\Upsilon \geq 65$. Several features of this galaxy make it an interesting case study.

²/ exposure maps: https://cxc.cfa.harvard.edu/ciao/threads/expmap_acis_single/

```
ule2in0.7pt
Citation\ `jones+2009mnras399_{6}83' on page 3 undefined on input line 135.
effective radius R_e. For this galaxy, R_{25} is 20 times larger than R_e, in contrast to R_{25} = 3.6 \times R_e, the nominal average for spirals and
Sos\ measured\ by\ [[][williams+2009mnras400_1665.zaw+2019apj872_134 classified NGC\ 5084 as an AGN according to the kewley+2001apj550_2000]
truncated surface brightness profile comeron + 2012 apj 759_9 8, a potential sign of past gravitational interactions, such a sminor and major measurements of the surface brightness profile comeron + 2012 apj 759_9 8, a potential sign of past gravitational interactions, such a sminor and major measurements and the surface brightness profile comeron + 2012 apj 759_9 8, a potential sign of past gravitational interactions, such a sminor and major measurements and the surface brightness profile comeron + 2012 apj 759_9 8, a potential sign of past gravitational interactions, such as minor and major measurements and the surface brightness profile comeron + 2012 apj 759_9 8.
ule2in0.7pt
Citation'borlaff + 2024apj967_169'onpage4undefinedoninputline141.
This project is the second publication in a series that will study the hot qasha losar ound qalaxies usin qX-ray observations from the X-ray observations from X-ray observations from the X-ray observations from X-ray observations from the X-ray observations from X-ray observations from the X-ray observations from X-ray observations fr
ule2in0.7pt
Citation \ 'borlaff+2024apj967_169' on page 4 undefined on input line 186.
The two main products of\ are surface brightness and signal-to-noise ratio maps, allowing the observer to identify potential sources and the surface brightness and signal-to-noise ratio maps, allowing the observer to identify potential sources and the surface brightness and signal-to-noise ratio maps, allowing the observer to identify potential sources and the surface brightness and signal-to-noise ratio maps, allowing the observer to identify potential sources and the surface brightness and signal-to-noise ratio maps, allowing the observer to identify potential sources and the surface brightness and signal-to-noise ratio maps, allowing the observer to identify potential sources and the surface brightness and surface bright
ule2in0.7pt
Citation'e beling + 2007 apj 661_3 3' on page 6 undefined on input line 192.
Citation'ebeling + 2010mnras407_83' on page 6 undefined on input line 192.
Citation' xue + 2011 apj 195_10' on page 6 undefined on in put line 192.
Adaptive spatial binning techniques (Voronoi, CSMOOTH) cappellari + 2003 mnras 342, 45, ebeling + 2006 mnras 368, 5 have been widely applications and the contraction of the contracti
ule2in0.7pt
LISTOFREFERENCESINmyPaper.tex
Number of citations: 151
2000 ascl. soft 03002S
2012 ascl. soft 08017R
2021pdaa.book....N
an an thar a mai a h + 1996 a p j 466_1 3
antonucci 1993 araa 31_473
argyle + 1990mnras 243_504
athanas soula + 2016 apj 821_9 0
baldi2023aap31_3
barway + 2009mnras 394_1991
beck + 2013 inbook_6 41
beckman 2021 book
blandford + 1974mnras169_395
borlaff + 2014aap570_103
borlaff + 2024apj967_169
boyle + 1998mnras 293_49
bridle + 1984araa22_319
britzen + 2018mnras478_3199
britzen + 2023apj951_106
broos + 2010apj714_1582
byrnemamahit + 2024arXiv2402.05196
capetti + 2002aap394_39
cappellari + 2003mnras 342_3 45
carignan + 1997aj113_1585
cerulo + 2017mnras472_254
chambers + 2016 arXiv1612.05560 \\
chen + 2020apj897_102
cheung + 2009apj181_548
clarke + 1998apj495_189
collaboration + 2013aap558_33
collaboration + 2018aj156_123
collaboration + 2022apj935_167
combes 2017 in proceedings_D i
comeron + 2010mnras402_2462
comeron + 2012apj759_98
davis + 2013nat494_328
```

```
davis + 2022mnras512_{1}522
del vecchio + 2014mnras 439_2736
devaucouleurs + 1991book
diehl + 2006mnras 368_497
dimatteo + 2005nat433_604
donath + 2022 in proceedings_9 8
dressler + 1983apj270_7
dressler + 1997apj490_577
ebeling + 2006mnras368_65
ebeling + 2007apj661_{3}3
ebeling + 2010mnras407_83
elichemoral + 2018aap617_113
emsellem + 2015mnras446_2468
evans + 2010apj189_37
fabbiano + 2017apj842_4
fabbiano + 2018apj855_131
ferrarese + 1996apj470_444
frasermckelvie + 2018mnras481_5580
freeman + 2001 in proceedings_76
fruscione + 2006 in proceedings_6 2701 V
gallimore + 2006aj132_546
qarciaburillo + 2021aap65298
giri + 2024sci11_1371101
gonzalez martin + 2009aap 506_1 107
gottesman + 1986mnras 219_7 59
hardcastle + 2020na88_101539
haverkorn + 2004aap421_1011
heckman + 1993 in proceedings_455
heckman + 2017 incollection_2 431
hodgeskluck + 2010apj710_1205
hodgeskluck + 2010apj717_37
hodgeskluck + 2012apj746_167
hodgeskluck + 2020apj903_35
hodgeskluck 2011 thesis
hopkins + 2009apj691_1168
hopkins + 2013mnras430_1901
hunter 2007 sci 9_9 0
irwin + 2012aj144_43
irwin + 2019aj158_21
jaffe + 1993nat364_213
jiang + 2023apj959_11
jones + 2004mnras 355747
jones + 2009mnras 399_6 83
jones + 2020apj891_133
juravnova + 2019mnras484_2886
keel + 2012mnras 420_878
kewley + 2001apj556_121
khim + 2015apj220_3
kim + 2015apj814_8
kim + 2017apj844_21
kinney + 2000apj537_152
koribalski + 2004aj128_16
kormendy + 1994 in proceedings_147
kormendy + 2004araa42_603
kormendy + 2013araa51_511
KRISHNAMOORTHY 200423
larson + 1980apj237_692
laurikainen + 2010mnras 405_1089
lawrence + 2010apj714_561
leroy + 2021apj257_43
ma + 2020apj900_164
ma + 2023apj948_61
machacek + 2010apj711_1316
madau + 2014araa52_415
```

 $marin 2014 mnras 441_551$ $mcquinn + 2018mnras477_3164$ $moellenhoff + 1987aap174_63$ $moore + 1996nat379_613$ $morse + 1998apj505_159$ $moustakas + 2006apj164_81$ $nagar + 1999apj516_97$ $nandi + 2021apj908_178$ $netzer 2015 araa 53_3 65$ $ohlson + 2024aj167_31$ $oke1971apj170_193$ $onishi + 2015apj806_39$ $onishi + 2017mnras468_4663$ $osterbrock + 1993apj414_552$ $osullivan + 2017mnras472_1482$ $osullivan + 2018aap618_{1}26$ $pan + 2019apj881_119$ $pasetto + 2018aap613_74$ $peschken + 2017mnras468_994$ $peterson 2006 in collection_7 7$ $ramosalmeida + 2017nat1_679$ $renaud + 2015mnras454_3299$ $rubin + 2014apj794_156$ $sanchez almeida + 2012 apj 756_163$ $saripalli + 2013mnras 436_690$ $schmitt + 2002apj575_150$ $sebastian + 2019apj883_189$ $sebastian + 2020mnras499_334$ $shankar + 2009apj690_20$ $shopbell + 1998apj493_129$ $silchenko 1998aap 330_412$ $smith + 2021mnras 500_1933$ $springel + 2005mnras361_776$ $tadhunter + 1989nat341_422$ $tully 1982 apj 257_3 89$ $urry + 1995pasp107_803$ $vandermarel + 1998aj116_2220$ $vika + 2009mnras400_1451$ $vollmer + 2012aap537_143$ $wang + 2019apj870_132$ $wang + 2024apj962_188$ wang + 2024 ar Xiv 2401.09172 $wiegert + 2015aj150_81$ $williams + 2009mnras400_1665$ $xue + 2011apj195_10$ $yang + 2017mnras464_70$ $younger + 2007apj670_269$ $yu2002mnras331_{9}35$ $zaw + 2019apj872_134$ $zeilinger + 1990mnras 246_3 24$ $zheng + 2022afz22_085004$

 $zubovas + 2022mnras515_1705$