

To view full GitHub Actions output, click on item at top of list here:
https://github.com/pmarcum/SAUNAS_III/actions/runs/11789342737/job/32837946188

[illegible]

LaTeX Warning: Reference `tab:Observations' on page 4 undefined on input line 188.
Line #188 -----

LaTeX Warning: Reference `tab:Observations' on page 6 undefined on input line 197.
 Line #197 -----

LaTeX Warning: Reference `fig:NGC5084' on page 8 undefined on input line 222.
Line #222 -----
\ref{fig:NGC5084} in section \ref{subsec:results_xray_ima}

LaTeX Warning: Reference `fig:NGC5084' on page 8 undefined on input line 247.

Line #247 -----

Figure \,ref{fig:NGC5084} presents the new \SAUNAS-processed images in the selected broad X-ray band (0.3--2.0 keV), overlayed on the optical morphology of the galaxy \citep[large FOV optical and near-infrared \$gri\$ image from Pan-STARRS,]{chambers+2016arXiv1612.05560}. Additionally, Fig.\,ref{fig:NGC5084_per_band} in Appendix \ref{Appendix:Xray_subbands} shows the three X-ray (\emph{soft:} 0.3-1.0 keV, \emph{medium:} 1.0-2.0 keV, \emph{hard:} 2.0 - 8.0 keV) bands. \par

LaTeX Warning: Reference `fig:NGC5084' on page 9 undefined on input line 252.

Line #252 -----

flux inside the 3σ contour in Fig.\,ref{fig:NGC5084}; (2) Core ($R<10''$); (3) Extended emission ($10''\leq R\leq 30''$); and (4-7) North, West, South, and East lobes. The lobe regions are defined by dividing the extended emission in 4 quadrants (with separations at 45° , 135° , 225° , 315°), excluding the emission from the core ($R<10''$) with a limit at $R=1.5''$. The separation between the core and extended regions $R=10''$ is defined based on the inspection of the X-ray morphology as the maximum radius where the X-ray emission does not show a significant elongation. The results are presented in Table \ref{tab:Xray_flux}. The fluxes take into account PSF deconvolution, and they exclude the emission from point sources identified by the \Chandra catalog of point sources (XRBs, AGN core). The analysis shows that the integrated emission from the north and east lobes is in excess of a 10σ detection. The south and west are substantially dimmer but significant above the background at 6.6σ and 3.2σ .

LaTeX Warning: Reference `fig:NGC5084' on page 9 undefined on input line 257.

Line #257 -----

In order to verify the results from the pipeline, we study the significance of the extended emission in Appendix \ref{Appendix:Xray_noPSFdeco_test} using two different, additional methodologies widely used in the literature. First, we determine if there is an excess of emission around the bright core of the galaxy by comparing the PSF surface brightness profile with the observed profile in the original \Chandra/ACIS observations, without applying Voronoi binning or PSF deconvolution. This methodology has been extensively used in the literature \citep{fabbiano+2017apj842_4,fabbiano+2018apj855_131,jones+2020apj891_133,ma+2020apj900_164,ma+2023apj948_61} to identify hot gas halos and other extended emission components. The results (see Fig.\,ref{fig:NGC5084_psf_profile}) show a clear excess of emission above that expected PSF scattering up to the same radial distance as predicted by the 3σ contours on Fig.\,ref{fig:NGC5084}, confirming that the X-ray emission of NGC\,5084 is not caused by PSF-scattered light from the bright core and that the extended emission is significant.

LaTeX Warning: Reference `fig:NGC5084_hst' on page 10 undefined on input line 293.

Line #293 -----

To further investigate the nature of this complex emission, we explored the available HST observations of NGC\,5084 (see Table \ref{tab:Observations}), presented in Fig.\,ref{fig:NGC5084_hst}.

LaTeX Warning: Reference `fig:NGC5084_hst' on page 12 undefined on input line 314.

Line #314 -----

The results are shown in the right panel of Fig.\,ref{fig:NGC5084_hst}. The model ellipse fit is shown displaced from the observed disk for clarity. The center of the ellipse is $(\alpha, \delta) = (200^{\text{h}}07^{\text{m}}01^{\text{s}}, -21^{\circ}08'27'')$ $\pm 0.1''$ arcsec, compatible with the location of the core of the galaxy. The disk diameter is $D=304^{+10}_{-11}$ pc, with a median inclination of $i=71.2^{+1.8}_{-1.7}$ degrees, and a position angle of $\theta = -5.1^{+1.1}_{-1.0}$ degrees, indicating that the relative orientation of the

circumnuclear disk is perpendicular to the galactic plane.

LaTeX Warning: Reference `fig:ALMA' on page 12 undefined on input line 323.

Line #323 -----

The positions that do not contain emission with an associated intensity higher than the noise at a 95% probability confidence level are masked. The velocity map (moment 1) of the CO(2-1) emission lines is represented in a color scale over the F702W HST/ACS image in the right panel of Fig.\ref{fig:ALMA}. The results show an unmistakable edge-on rotation pattern, with the north (south) edge of the disk moving towards (away from) the observer along the line of sight. The peaks of CO(2-1) emission are coincident with the edges of the circumnuclear disk, where the column density is expected to be highest due to projection effects.\

LaTeX Warning: Reference `fig:ALMA' on page 12 undefined on input line 325.

Line #325 -----

The spectra of the locations with significant emission are combined (summed) to study the line profile and characterize the amplitude of the rotation pattern. The combined spectra is shown in the left panel of Fig.\ref{fig:ALMA}. We characterize the rotation amplitude using the line width at the half-maximum, following a similar procedure as in \citep[W\$_{50}\$,][smith+2021mnras500_1933]. The velocities (lowest and highest) at which the spectrum reaches its half-maximum are measured through interpolation of the combined spectra. This process is repeated using $N=1000$ Monte Carlo simulations, with each data point being displaced by its spectral flux uncertainty (noise level). The measured rotation velocity is $V_{\rm rot} = 249.1^{+12.8}_{-9.6}$ km s⁻¹ ($W_{50}=498^{+26}_{-19}$ km s⁻¹).

LaTeX Warning: Reference `tab:Radio_Lobes' on page 13 undefined on input line 353.

Line #353 -----

Interestingly, \citep{wiegers+2015aj150_81} identifies extended emission to the east and west of the central bright core. This emission is highlighted in Fig.\ref{fig:NGC5084_EVLA}, and was identified as radio lobes in \citep[see their Table 10]{irwin+2019aj158_21}. These two radio sources are detected at a 10 σ level, being located at a symmetric distance from the core. The distance from the east lobe to the core is $R=31.5^{+4.5}_{-4.5}$ arcsec or $R=4.52^{+0.66}_{-0.65}$ kpc, while the equivalent from the west lobe is $R=31.1^{+4.5}_{-4.6}$ arcsec ($R=4.51^{+0.65}_{-0.67}$ kpc), being compatible at a 3 σ confidence level. The position angles from the east and west lobes to the center are also compatible at a 3 σ level, $PA_{\rm east - core} = 92.8^{+6.4}_{-7.1}$, $PA_{\rm core - west} = 92.2^{+7.1}_{-7.6}$. Taking into account the inclination of the circumnuclear disk ($i=71.2^{+1.8}_{-1.7}$), and assuming that the lobes are oriented along the line of the AGN radio jet axis, the deprojected distance to the core is $R=4.8 \pm 0.70$ kpc, taking into account the uncertainties in the location of the core, radio lobes, and inclination. The results are compatible at the 1.4 and 5 GHz bands. The total luminosity of each lobe in 5 GHz is $L_{\rm 5\,GHz} = [1.5, 1.8] \times 10^{+19}$ W Hz⁻¹ while the core is two orders of magnitude brighter with $L_{\rm 5\,GHz, core} = 3.60 \pm 0.01 \times 10^{+21}$ W Hz⁻¹. In 1.4 GHz, the lobes are brighter $L_{\rm 1.4\,GHz} = [4, 6] \times 10^{+19}$ W Hz⁻¹, with $L_{\rm 5\,GHz, core} = 3.60 \pm 0.01 \times 10^{+21}$ W Hz⁻¹ in the core. The distances to the core of each lobe and the luminosities of each component are included in Table \ref{tab:Radio_Lobes}.

LaTeX Warning: Command \r invalid in math mode on input line 440.

Line #440 -----

using five regions: (1) the central 1.5 arcsec from the core of the galaxy; (2-5) north, south, east, and west sides of the galaxy, resp. (not including the core). The results are shown in Fig.\ref{fig:NGC5084_optical_spectra}. The APO/DIS spectra do not reveal any signs of H α [$\lambda=6562.8$ \AA], H β [$\lambda=4861.35$ \AA], or

H γ ($\lambda=4340.47\text{\AA}$) absorption, in any of the regions analyzed, suggesting that the stellar population at the core is not dominated by a classic post-starburst stellar population.

LaTeX Warning: Reference `subsec:discussion_fadedstarburst' on page 18 undefined on input line 472.

Line #472 -----

These scenarios are idealized; in reality, a mixture of these cases may be responsible for the observed phenomena. In particular, the HI tilt and the disk warp of NGC\,5084 suggest that the galaxy suffered a major merger at some point in its past. Rather than directly eliminating other scenarios, mergers serve as a potential triggering factor in two of the other cases (orientation change and faded starburst). The observational evidence for merger activity is gathered in Sec.\,\ref{subsec:discussion_merger}. Secs.\,\ref{subsec:discussion_fadedstarburst}, \ref{subsec:discussion_cocoon}, and \ref{subsec:discussion_realignment} discuss each of the scenarios presented in Fig.\,\ref{fig:NGC5084_scenarios}.

LaTeX Warning: Reference `subsec:discussion_fadedstarburst' on page 18 undefined on input line 480.

Line #480 -----

\caption{Summary of the formation scenarios for the vertical X-ray emission of NGC\,5084. Cosmic time increases from left to right. \emph{Top row:} (1) Orientation change of AGN jet. Mergers, inflows, precession, and SMBH interactions can reorient the jet direction of an AGN over time; \emph{Central row:} (2) Overpressured cocoon. AGN-jet emission directed against the galactic disk can result in an expansion along the minor axis of the galaxy; \emph{Top:} (3) Faded starburst. Circumnuclear starbursts can generate galactic winds, expelling hot gas in the direction vertical to the galactic plane. However, this scenario requires active star formation or at least a relatively young population of stars at the core, which is not observed (see Sec.\,\ref{subsec:discussion_fadedstarburst}).}

LaTeX Warning: Reference `fig:NGC5084' on page 20 undefined on input line 523.

Line #523 -----

\caption{NGC\,5084 color maps. \emph{Left panel:} Ground-based VLT/ATLAS $g-i$ maps. Black contours represent the X-ray emission (see Fig.\,\ref{fig:NGC5084}). Dark blue square in the core represents the FOV in the right panel. \emph{Right panel:} F475W - F702W color from \emph{Hubble} WFPC2/WFC3 photometry. The circumnuclear disk is clearly visible in red. See the colorbar for reference.}

LaTeX Warning: Reference `subsec:results_Optical_spectra' on page 20 undefined on input line 529.

Line #529 -----

Secondly, the optical spectra (see Sec.\,\ref{subsec:results_Optical_spectra}, Fig.\,\ref{fig:NGC5084_optical_spectra}) do not present any Balmer absorption lines typical of post-starburst objects \cite[also called E+A or K+A galaxies,][dressler+1983apj270_7], not even in regions away from the core, where the AGN could dominate the emission. Given that the post-starburst phase lasts about 300\,Myr, the lack of a Balmer absorption line signal in the optical spectra suggest that the age of the stellar population in the core is older than a Gyr.

LaTeX Warning: Reference `tab:Radio_Lobes' on page 22 undefined on input line 537.

Line #537 -----

Unfortunately, the systematic effects observed in radio observations along the vertical direction (Sec.\,\ref{subsec:results_radiopol}) prevent the identification of additional radio lobes along the minor axis of the galaxy, challenging the ability to determine whether NGC\,5084 also has an X-shape radio morphology. Given the detected radio-lobes,

NGC\,5084 would be classified between high-luminosity LINER/Seyferts and FR \sim 0 galaxies \citep[see Table \ref{tab:Radio_Lobes},][{baldi2023aap31_3}], emitting ~ 4 --5 orders of magnitude less than the X-shaped radio galaxies from \citep{cheung+2009apj181_548}. This classification correlates with the prediction from \citep{hodgeskluck2011thesis} that X-shaped sources should be decaying AGN jets.

LaTeX Warning: Command \r invalid in math mode on input line 567.

Line #567 -----

Another example of such phenomena is the lenticular galaxy NGC\,5252. Classified as a Seyfert 1.9 galaxy \citep{argyle+1990mnras243_504, osterbrock+1993apj414_552}, NGC\,5252 presents a large-scale ionization bi-cone \citep{tadhunter+1989nat341_422} detectable in [OIII] emission line at $\lambda=5007\text{\AA}$ that extends for ~ 20 kpc. \emph{Hubble} Space Telescope and Fabry--Pérot spectrograph observations of NGC\,5252 by \citep{morse+1998apj505_159} revealed different kinematic components, including an inclined circumnuclear gas disk with a diameter of 3 kpc, suggesting that NGC\,5252 underwent a galaxy merger in its past history. This scenario is supported by observational evidence of the presence of both a supermassive (main) and an intermediate (accreting) mass black hole in NGC\,5252, both active and emitting in radio \citep{kim+2015apj814_8, kim+2017apj844_21, yang+2017mnras464_70}. Moreover, recent results presented by \citep{wang+2024arXiv2401.09172} show a strong ($\sim 20^\circ$) misalignment between the X-ray emission in soft bands (0.3--2.0 keV) and the optical major axis of the galaxy. The case of NGC\,5252 shows that the combination of observational evidence such as the presence of off-axis morphological components and AGN-related features is one strategy to shed light on the formation pathways of specific galaxies.

LaTeX Warning: Reference `sec:DIS' on page 25 undefined on input line 591.

Line #591 -----

The horizontal (in-plane) component of the X-ray emission is aligned with both the rotation axis of the newly discovered circumnuclear disk and the symmetric radio lobes, identified by \citep{irwin+2019aj158_21} as part of a radio AGN jet. Taking into account all available observational evidence consolidated in this paper (see Fig.\,\ref{fig:NGC5084_scenarios}), we consider three potential hypotheses for the formation of the observed cross-shaped X-ray emission: (1) it is the remnant of a re-oriented AGN; (2) it is an outflow generated by an overpressured cocoon of hot gas powered by the AGN pointing into the dense ISM within midplane of the galactic disk; or (3) it is part of a faded starburst at the core of the galaxy. Spectroscopic observations on the core of NGC\,5084 do not support the latter scenario, given the lack of spectral evidence for recent (massive stars) or on-going star formation in the core. Combining the new observational evidence presented in this paper with previous analysis based on environment and morphology (see Sec.\,\ref{sec:DIS}), we conclude that NGC\,5084 is with high probability the remnant of at least one merger in the past, which is actively accreting its multiple satellites.}

LaTeX Warning: Reference `eq:F_lim_Xray' on page 27 undefined on input line 655.

Line #655 -----

At $E=1$ keV and $t=10$ s, the equivalent exposure\footnote{\ciao/Chandra exposure maps: \url{https://cxc.cfa.harvard.edu/ciao/threads/expmap_acis_single/}} (M) is approximately $3.8 \times 10^6 \text{ cm}^2 \text{ s}$. For a point source detection of $\sigma=3$, with flux integration over an area associated with the PSF at the center of the ACIS detector ($\text{FWHM} \sim 1.1''$), Eqn.\,\ref{eq:F_lim_Xray} implies a limiting sensitivity of $f_{\text{lim}} = 4 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$, equal to the reference instrument point source sensitivity limits.

LaTeX Warning: Reference `tab:Xray_flux' on page 29 undefined on input line 699.

LaTeX Warning: Reference `fig:NGC5084_Aperture_Histograms' on page 29 undefined on input

line 699.

Line #699 -----

In this section we assess the statistical significance of the detection of the four X-ray emission lobes reported in Sec.\,\ref{subsec:results_xray_ima}, using the PSF deconvolved observations but without employing Voronoi binning. First, four simple box regions are defined, based on the contours detected in Sec.\,\ref{subsec:results_xray_ima} to isolate the emission from the brighter galactic core. The region parameters are listed in Table \ref{tab:Xray_flux} and represented in the left panel of Fig.\,\ref{fig:NGC5084_Aperture_Histograms} (see legend in the right panel). Second, a Poisson means test \citep[\$\chi^2\$-test,]{KRISHNAMOORTHY200423} evaluates the null hypothesis that the difference between the observed and background emission is statistically zero. We repeat the test 500 times using Monte Carlo and Bootstrapping simulations in order to obtain the probability distributions for the surface brightness represented in the right panel of Fig.\,\ref{fig:NGC5084_Aperture_Histograms}.

LaTeX Warning: Reference `tab:Xray_flux' on page 30 undefined on input line 701.

Line #701 -----

The analysis indicates that the surface brightness X-ray emission in the apertures is significantly higher than that of the background. The null hypothesis that the surface brightness distributions of lobe emission and the background are sampled from a common parent population is rejected at a p-value of $p < 0.05$ in all four lobe regions. In other words, this analysis provides strong statistical support for the existence of the lobe emission: north ($p = 6.8 \times 10^{-7}$), east ($p = 1.1 \times 10^{-6}$), south ($p = 1.3 \times 10^{-4}$) and west ($p \sim 0.01$). The fluxes integrated over the different apertures are tabulated in Table \ref{tab:Xray_flux}. This result supports and verifies the findings described in Sec.\,\ref{subsec:results_xray_ima} and Appendix \ref{Appendix:Xray_noPSFdeco_test}. We conclude that the extended X-ray emission detected by \textit{SAUNAS} around NGC\,5084 is (1) statistically significant; (2) independent of the PSF deconvolution process; and (3) independent of the Voronoi binning methodology applied.

LaTeX Warning: Reference `sec:methods' on page 30 undefined on input line 711.

Line #711 -----

\caption{Detection of the X-ray extended lobes around NGC\,5084 without Voronoi binning. \emph{Left:} \Chandra/ACIS X-ray flux (\$0.3\$-\$2.0\$ keV) map rebinned to \$12.2\times12.2\$ arcsec (\$25\times25\$ pixels) for visualization purposes. PSF deconvolution was applied \citep[see Sec.\,\ref{sec:methods}, and]{borlaff+2024apj967_169} \emph{Color rectangles:} Fixed apertures to measure X-ray emission in the North (blue), West (red), South (green), and East (orange) lobes. \emph{Yellow contours:} [\$2\sigma, 3\sigma]\$\sigma\$ (dotted, dashed respectively) detection limits as estimated by \code{SAUNAS} (see Sec.\,\ref{subsec:results_xray_ima}, for reference. \emph{Right:} Color coded histograms represent the event probability distributions for the background (grey) and the four lobe apertures. The p -values for the null hypothesis that the flux distributions in the lobe apertures are compatible with the background are represented in the legend.}

[illegible][illegible]

Line #118 -----

NGC\,5084 is one of the most massive lenticular galaxies in the Local Universe \citep{ohlson+2024aj167_31}, with a total dynamical mass of approximately $1.3 \times 10^{12} \text{ M}_{\odot}$ \citep{koribalski+2004aj128_16} and a mass-to-light ratio $\Upsilon \geq 65$. Several features of this galaxy make it an interesting case study.

Citation 'jones+2009mnras399_683' on page 3 undefined on input line 135.

Line #135 -----

effective radius $R_{\rm e}$. For this galaxy, R_{25} is 20 times larger than $R_{\rm e}$, in contrast to $R_{25}=3.6 \times R_{\rm e}$, the nominal average for spirals and S0s measured by \citet{williams+2009mnras400_1665}. \citet{zaw+2019apj872_134} classified NGC\,5084 as an AGN according to the \citet{kewley+2001apj556_121} optical spectral line ratio criteria, based on 6dF Galaxy Survey spectra \citet{jones+2004mnras355_747,jones+2009mnras399_683}, as did \citet{irwin+2019aj158_21}, based on its radio emission (see Sec.\ref{subsec:data_radiopol}). In addition, NGC\,5084 is the host of a large spheroidal bulge that generates an anti-truncated surface brightness profile \citet{comeron+2012apj759_98}, a potential sign of past gravitational interactions, such as minor and major mergers \citet{younger+2007apj670_269,borlaff+2014aap570_103}.

Citation 'borlaff+2024api967_169' on page 4 undefined on input line 141.

Line #141 -----

This project is the second publication in a series that will study the hot gas halos around galaxies using X-ray observations from the \Chandra\ X-ray observatory. The first paper \citep[\texttt{SAUNAS I}, hereafter]{borlaff+2024apj967_169} describes the \SAUNAS\ (Selective Amplification of Ultra Noisy Astronomical Signal) pipeline to detect low surface brightness emission in \Chandra/ACIS observations. See \texttt{SAUNAS I} for the details and description of the X-ray image reduction methodology. \par

Citation 'borlaff+2024apj967_169' on page 4 undefined on input line 186.

Line #186 -----

The two main products of \SAUNAS\ are surface brightness and signal-to-noise ratio maps, allowing the observer to identify potential sources and their extension up to a certain statistical limit. Since the objective is to identify the underlying shape of the X-ray extended emission in NGC\,5084, PSF deconvolution is a particularly critical step. \citet{borlaff+2024apj967_169} provides a full description of the methodology.

Citation 'ebeling+2007apj661_33' on page 6 undefined on input line 192.

Citation 'ebeling+2010mnras407 83' on page 6 undefined on input line 192.

Citation 'xue+2011apj195_10' on page 6 undefined on input line 192.

Line #192 -----

Adaptive spatial binning techniques (Voronoi, CSMOOTH) \citep{cappellari+2003mnras342_345, ebeling+2006mnras368_65} have been widely applied in X-ray astronomy for several decades \citep[see][and references therein]{ebeling+2007apj661_33, gonzalezmartin+2009aap506_1107, broos+2010apj714_1582, ebeling+2010mnras407_83, xue+2011apj195_10, hodgeskluck+2012apj746_167, wang+2024apj962_188} allowing the detection of extended sources below the canonical point source sensitivity of the observations. While \texttt{SAUNAS I} \citep{borlaff+2024apj967_169} is dedicated to the presentation, benchmarking, and testing of the adaptive binning technique used by the pipeline, Appendix \ref{Appendix:xray_tests} in this paper provides a more in-depth discussion of point source and extended source sensitivities, and a series of tests to further verify the significance of the X-ray emission of NGC\,5084 reported in Sec.\,\ref{subsec:results_xray_ima}.

[illegible]

LIST OF REFERENCES IN myPaper.tex

[illegible]

Number of citations: 151

2000ascl.soft03002S

2012ascl.soft08017R
2021pdaa.book.....N
anantharamaiah+1996apj466_13
antonucci1993araa31_473
argyle+1990mnras243_504
athanassoula+2016apj821_90
baldi2023aap31_3
barway+2009mnras394_1991
beck+2013inbook_641
beckman2021book
blandford+1974mnras169_395
borlaff+2014aap570_103
borlaff+2024apj967_169
boyle+1998mnras293_49
bridle+1984araa22_319
britzen+2018mnras478_3199
britzen+2023apj951_106
broos+2010apj714_1582
brynemamahit+2024arXiv2402.05196
capetti+2002aap394_39
cappellari+2003mnras342_345
carignan+1997aj113_1585
cerulo+2017mnras472_254
chambers+2016arXiv1612.05560
chen+2020apj897_102
cheung+2009apj181_548
clarke+1998apj495_189
collaboration+2013aap558_33
collaboration+2018aj156_123
collaboration+2022apj935_167
combes2017inproceedings_Di
comeron+2010mnras402_2462
comeron+2012apj759_98
davis+2013nat494_328
davis+2022mnras512_1522
delvecchio+2014mnras439_2736
devaucouleurs+1991book
diehl+2006mnras368_497
dimatteo+2005nat433_604
donath+2022inproceedings_98
dressler+1983apj270_7
dressler+1997apj490_577
ebeling+2006mnras368_65
ebeling+2007apj661_33
ebeling+2010mnras407_83
elichemoral+2018aap617_113
emsellem+2015mnras446_2468
evans+2010apj189_37
fabbiano+2017apj842_4
fabbiano+2018apj855_131
ferrarese+1996apj470_444
frasermckelvie+2018mnras481_5580
freeman+2001inproceedings_76
fruscione+2006inproceedings_62701V
gallimore+2006aj132_546
garciaburillo+2021aap652_98

giri+2024sci11_1371101
gonzalezmartin+2009aap506_1107
gottesman+1986mnras219_759
hardcastle+2020na88_101539
haverkorn+2004aap421_1011
heckman+1993inproceedings_455
heckman+2017incollection_2431
hodgeskluck+2010apj710_1205
hodgeskluck+2010apj717_37
hodgeskluck+2012apj746_167
hodgeskluck+2020apj903_35
hodgeskluck2011thesis
hopkins+2009apj691_1168
hopkins+2013mnras430_1901
hunter2007sci9_90
irwin+2012aj144_43
irwin+2019aj158_21
jaffe+1993nat364_213
jiang+2023apj959_11
jones+2004mnras355_747
jones+2009mnras399_683
jones+2020apj891_133
juravnova+2019mnras484_2886
keel+2012mnras420_878
kewley+2001apj556_121
khim+2015apj220_3
kim+2015apj814_8
kim+2017apj844_21
kinney+2000apj537_152
koribalski+2004aj128_16
kormendy+1994inproceedings_147
kormendy+2004araa42_603
kormendy+2013araa51_511
KRISHNAMOORTHY200423
larson+1980apj237_692
laurikainen+2010mnras405_1089
lawrence+2010apj714_561
leroy+2021apj257_43
ma+2020apj900_164
ma+2023apj948_61
machacek+2010apj711_1316
madau+2014araa52_415
marin2014mnras441_551
mcquinn+2018mnras477_3164
moellenhoff+1987aap174_63
moore+1996nat379_613
morse+1998apj505_159
moustakas+2006apj164_81
nagar+1999apj516_97
nandi+2021apj908_178
netzer2015araa53_365
ohlson+2024aj167_31
oke1971apj170_193
onishi+2015apj806_39
onishi+2017mnras468_4663
osterbrock+1993apj414_552

osullivan+2017mnras472_1482
osullivan+2018aap618_126
pan+2019apj881_119
pasetto+2018aap613_74
peschken+2017mnras468_994
peterson2006incollection_77
ramosalmeida+2017nat1_679
renaud+2015mnras454_3299
rubin+2014apj794_156
sanchezalmeida+2012apj756_163
saripalli+2013mnras436_690
schmitt+2002apj575_150
sebastian+2019apj883_189
sebastian+2020mnras499_334
shankar+2009apj690_20
shopbell+1998apj493_129
silchenko1998aap330_412
smith+2021mnras500_1933
springel+2005mnras361_776
tadhunter+1989nat341_422
tully1982apj257_389
urry+1995pasp107_803
vandermaarel+1998aj116_2220
vika+2009mnras400_1451
vollmer+2012aap537_143
wang+2019apj870_132
wang+2024apj962_188
wang+2024arXiv2401.09172
wiegert+2015aj150_81
williams+2009mnras400_1665
xue+2011apj195_10
yang+2017mnras464_70
younger+2007apj670_269
yu2002mnras331_935
zaw+2019apj872_134
zeilinger+1990mnras246_324
zheng+2022afz22_085004
zubovas+2022mnras515_1705