



Relativistic time of arrival

(Thesis proposal)

Philip Caesar M. Flores

pmflores2@up.edu.ph

Theoretical Physics Group, National Institute of Physics,
University of the Philippines Diliman, Quezon City

05 October 2021

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

1 Introduction

- Classical time of arrival
- Quantum time of arrival
- Time operators

2 Non-relativistic TOA operators

- Free case
- Interacting case

3 Relativistic TOA operators

- Motivation
- Related studies

4 Research proposal

- Comments
- Goals

Current Physical Theories

Introduction

Classical time of arrival

Quantum time of arrival

Time operators

Non-relativistic TOA operators

Free case

Interacting case

Relativistic TOA operators


Motivation

Related studies

Research proposal

Comments

Goals



- **Classical mechanics** treats time only as a parameter:
 - Newton's equations
 - Maxwell's equations
- Time is **absolute**.

Current Physical Theories

Introduction

Classical time of arrival

Quantum time of arrival

Time operators

Non-relativistic TOA operators

Free case

Interacting case

Relativistic TOA operators


Motivation

Related studies

Research proposal

Comments

Goals



- Time in relativity is **dynamical**
 - SR treats time and space as components of a **4-vector**
 - Time in GR is influenced by the geometry of **spacetime**

Current Physical Theories

Introduction

Classical time of arrival

Quantum time of arrival

Time operators

Non-relativistic TOA operators

Free case

Interacting case

Relativistic TOA operators


Motivation

Related studies

Research proposal

Comments

Goals



- Quantum mechanics treats time as an **external parameter** which governs the evolution of the system
- Quantization schemes lack any fundamental notion of time

Current Physical Theories

Introduction

Classical time of arrival

Quantum time of arrival

Time operators

Non-relativistic TOA operators

Free case

Interacting case

Relativistic TOA operators


Motivation

Related studies

Research proposal

Comments

Goals



- Different treatment of time then poses a problem in attempts to unify general relativity and quantum theory
- This constitutes one aspect of the problem of time in quantum gravity

Time of arrival in classical physics

Introduction

Classical time of arrival

Quantum time of arrival

Time operators

Non-relativistic TOA operators

Free case

Interacting case

Relativistic TOA operators


Motivation

Related studies

Research proposal

Comments

Goals



Method 1:

We invert the classical equation of motion

$$x = q + \frac{p}{\mu}t \Rightarrow t = \mu \frac{(x - q)}{p} \quad (1)$$

where, μ is the mass of the particle and p is the initial momentum

Time of arrival in classical physics

Introduction

Classical time of arrival

Quantum time of arrival

Time operators

Non-relativistic TOA operators

Free case

Interacting case

Relativistic TOA operators


Motivation

Related studies

Research proposal

Comments

Goals



Method 2:

Use a detector to record the instant of time t_q at which the particle leaves q and another detector to record its arrival t_x at x .

The difference $\Delta t = t_x - t_q$ is the time of arrival.

- Methods 1 and 2 will always yield the **same result**.

Time of arrival in classical physics

Introduction

Classical time of arrival

Quantum time of arrival

Time operators

Non-relativistic TOA operators

Free case

Interacting case

Relativistic TOA operators


Motivation

Related studies

Research proposal

Comments

Goals



Interacting case:

Given the Hamiltonian $H(q, p) = \frac{p^2}{2\mu} + V(q)$, the time of arrival is

$$T_x(q, p) = -\text{sgn}(p) \sqrt{\frac{\mu}{2}} \int_x^q \frac{dq'}{\sqrt{H(q, p) - V(q')}} \quad (2)$$

Time of arrival in quantum physics

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

We cannot use method 1

In the Heisenberg picture, the equation of motion of the position operator is

$$\frac{d\hat{x}}{dt} = \frac{i}{\hbar} [\hat{H}, \hat{x}] \quad (3)$$

- We can quantize the classical expression of the time of arrival but this leads to **ordering ambiguity**.
- The classical TOA may be **complex and/or multiple valued**.

Time of arrival in quantum physics

Introduction

Classical time of arrival

Quantum time of arrival

Time operators

Non-relativistic TOA operators

Free case

Interacting case

Relativistic TOA operators

Motivation

Related studies

Research proposal

Comments

Goals

We cannot use method 2

The first detector will collapse the wavefunction and the state of the particle upon arrival at x is no longer causally related to the state during preparation.

- The standard formulation of quantum mechanics has **no standard solution** to the time of arrival of a particle.
- It is **non-sensical** to ask the time of arrival of the particle because time is not an observable in quantum mechanics.

Time operators

Introduction

Classical time of arrival

Quantum time of arrival

Time operators

Non-relativistic TOA operators

Free case

Interacting case

Relativistic TOA operators

Motivation

Related studies

Research proposal

Comments

Goals

Pauli's theorem:¹

There is no self-adjoint time operator that is canonically conjugate with its corresponding bounded system Hamiltonian .

- This has lead to a diverse treatment of time **within and beyond the standard formulation** of quantum mechanics to bypass Pauli's theorem².
- Classical mechanis is always a **fundamental reference** for all approaches.

¹ Wolfgang Pauli et al. "Handbuch der physik". In: *Geiger and scheel 2* (1933), pp. 83–272.

² Juan Gonzalo Muga and C Richard Leavens. "Arrival time in quantum mechanics". In: *Physics Reports* 338.4 (2000), pp. 353–438; Iñigo L. Egusquiza J. Gonzalo Muga Rafael Sala Mayato. *Time in quantum mechanics Volume 1*. 2nd ed. Lecture Notes in Physics 734. Springer-Verlag Berlin Heidelberg, 2007. ISBN: 9783540734727; 3540734724; Adolfo Campo Gonzalo Muga Andreas Ruschhaupt. *Time in Quantum Mechanics - Vol. 2*. 1st ed. Lecture Notes in Physics 789. Springer-Verlag Berlin Heidelberg, 2009. ISBN: 3642031730; 9783642031731.

Time operators

Introduction

Classical time of arrival

Quantum time of arrival

Time operators

Non-relativistic TOA operators

Free case

Interacting case

Relativistic TOA operators

Motivation

Related studies

Research proposal

Comments

Goals

- Pauli's proof was only **formal**, without regard to the domains of the operators involved and to the validity of the operations leading to his conclusion³.
- Several studies on time of arrival operators have also been made.

³Eric Galapon. "Pauli's theorem and quantum canonical pairs: the consistency of a bounded, self-adjoint time operator canonically conjugate to a Hamiltonian with non-empty point spectrum". In: *Proceedings of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences* 458.2018 (2002), pp. 451–472.

Free time of arrival operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

Aharonov-Bohm operator⁴

Symmetric quantization of the classical TOA yields

$$\hat{T}_{AB} = -\frac{\mu}{2} \left(\hat{p}^{-1} \hat{q} + \hat{q} \hat{p}^{-1} \right) \quad (4)$$

- This is a **maximally symmetric** operator.
- The eigenfunctions are **complete but non-orthogonal**⁵

$$\langle p|t, \pm \rangle = \sqrt{\frac{|p|}{\mu \hbar}} \exp\left(\frac{i}{\hbar} \frac{p^2}{2\mu} t\right) \Theta(\pm p) \quad (5)$$

⁴Yakir Aharonov and David Bohm. "Time in the quantum theory and the uncertainty relation for time and energy". In: *Physical Review* 122.5 (1961), p. 1649.

⁵JG Muga, CR Leavens, and JP Palao. "Space-time properties of free-motion time-of-arrival eigenfunctions". In: *Physical Review A* 58.6 (1998), p. 4336.

Free time of arrival operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

Grot-Rovelli-Tate operator⁶

Regularize \hat{T}_{AB} at the point $p = 0$ which yields

$$\hat{T}_{GRT} = -\mu \sqrt{f_{\epsilon}(\hat{p})} \hat{q} \sqrt{f_{\epsilon}(\hat{p})} \quad (6)$$

$$f_{\epsilon}(\hat{p}) = \begin{cases} \hat{p}^{-1}, & |p| > \epsilon \\ \epsilon^2 \hat{p}, & |p| < \epsilon \end{cases} \quad (7)$$

- This is a **self-adjoint** operator.
- **Regularization** introduces negative energies to bypasses Pauli's theorem.

⁶Norbert Grot, Carlo Rovelli, and Ranjeet S Tate. "Time of arrival in quantum mechanics". In: *Physical Review A* 54.6 (1996), p. 4676.

Free time of arrival operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

Grot-Rovelli-Tate operator⁶

Regularize \hat{T}_{AB} at the point $p = 0$ which yields

$$\hat{T}_{GRT} = -\mu \sqrt{f_{\epsilon}(\hat{p})} \hat{q} \sqrt{f_{\epsilon}(\hat{p})} \quad (6)$$

$$f_{\epsilon}(\hat{p}) = \begin{cases} \hat{p}^{-1}, & |p| > \epsilon \\ \epsilon^2 \hat{p}, & |p| < \epsilon \end{cases} \quad (7)$$

- The eigenfunctions are

$$\langle p|t, \pm \rangle_{\epsilon} = \sqrt{\frac{1}{\mu \hbar f_{\epsilon}(p)}} \exp\left(\frac{i}{\hbar} \frac{t}{\mu} \int_{\pm \epsilon}^p \frac{dp'}{f_{\epsilon}(p')}\right) \Theta(\pm p) \quad (8)$$

⁶Norbert Grot, Carlo Rovelli, and Ranjeet S Tate. "Time of arrival in quantum mechanics". In: *Physical Review A* 54.6 (1996), p. 4676.

Free time of arrival operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

Kijowski-Delgado-Muga operator⁷

Self-adjoint extension of \hat{T}_{AB} is obtained using the combination

$$\hat{T}_{KDM} = \hat{T}_{AB}\Theta(\hat{p}) - \hat{T}_{AB}\Theta(-\hat{p}) \quad (9)$$

- Bypasses Pauli's theorem because \hat{T}_{KDM} is not canonically conjugate to \hat{H} , instead

$$\left[\text{sgn}(\hat{p})\hat{H}, \hat{T}_{KDM} \right] = i\hbar \quad (10)$$

⁷ Jerzy Kijowski. "On the time operator in quantum mechanics and the Heisenberg uncertainty relation for energy and time". In: *Reports on Mathematical Physics* 6.3 (1974), pp. 361–386; V Delgado and JG Muga. "Arrival time in quantum mechanics". In: *Physical Review A* 56.5 (1997), p. 3425.

Free time of arrival operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

Confined time of arrival operator⁸

The operator in position space is

$$\hat{T}_\gamma \varphi(q) = \int_{-l}^l dq' T_\gamma(q, q') \varphi(q') \quad (11)$$

$$T_\gamma(q, q') = -\mu \frac{(q + q')}{4\hbar \sin \gamma} \left(e^{i\gamma} \Theta(q - q') + e^{-i\gamma} \Theta(q' - q) \right) \quad (12)$$

- Self adjointness was addressed by **spatial confinement** and imposing **non-vanishing boundary conditions** $\varphi(-l) = e^{-2i\gamma} \varphi(l)$ in the system Hilbert space.

⁸Eric A Galapon, Roland F Caballar, and Ricardo T Bahague Jr. "Confined quantum time of arrivals". In: *Physical review letters* 93.18 (2004), p. 180406; Eric A Galapon, Roland F Caballar, and Ricardo Bahague. "Confined quantum time of arrival for the vanishing potential". In: *Physical Review A* 72.6 (2005), p. 062107.

Confined time of arrival operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

- In the limit of **large confining length**, the eigenfunctions become two-fold degenerate and are related to the Aharonov-Bohm TOA eigenfunctions as

$$\langle p|t\rangle_{non} = \frac{1}{\sqrt{2}} (\langle p|t, +\rangle - \langle p|t, -\rangle) \quad (13)$$

$$\langle p|t\rangle_{nod} = \frac{1}{\sqrt{2}} (\langle p|t, +\rangle + \langle p|t, -\rangle) \quad (14)$$

- Non-nodal: arrival with detection⁹
- Nodal: arrival with non-detection

⁹Denny Lane B Sombillo and Eric A Galapon. "Particle detection and non-detection in a quantum time of arrival measurement". In: *Annals of Physics* 364 (2016), pp. 261–273.

Confined time of arrival operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators



Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals



- The eigenfunctions exhibit **unitary collapse**¹⁰
- The peaks occur at at time equal to the CTOA eigenvalue where the position uncertainty is minimum.

¹⁰Eric A Galapon. "Theory of quantum arrival and spatial wave function collapse on the appearance of particle". In: *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* 465.2101 (2009), pp. 71–86.

TOA operators in the interacting case

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

■ Quantizing the classical TOA

$$T_x(q, p) = -\text{sgn}(p) \sqrt{\frac{\mu}{2}} \int_x^q \frac{dq'}{\sqrt{H(q, p) - V(q')}}$$

has been deemed **not meaningful** because it is not generally real and single valued everywhere¹¹ but can be addressed on physical grounds¹²

- The TOA of a quantum particle is always real and single valued because
 - it can tunnel through the classically forbidden region
 - measuring the first TOA will collapse the wavefunction

¹¹ Asher Peres. *Quantum theory: concepts and methods*. Vol. 57. Springer Science & Business Media, 2006.

¹² Eric A Galapon and John Jaykel P Magadan. "Quantizations of the classical time of arrival and their dynamics". In: *Annals of Physics* 397 (2018), pp. 278–302.

Quantized time of arrival operators

■ Sum of the Bender-Dunne basis operators¹³

$$\hat{T}_{m,n} = \frac{1}{\sum_{k=0}^n a_k^{(n)}} \sum_{k=0}^n a_k^{(n)} \hat{q}^k \hat{p}^m \hat{q}^{n-k} \quad (15)$$

$$= \frac{1}{\sum_{j=0}^m a_j^{(m)}} \sum_{j=0}^m a_j^{(m)} \hat{p}^j \hat{q}^n \hat{p}^{m-j} \quad (16)$$

$$a_k^{(n)} = \begin{cases} \binom{n}{k}, & \text{Weyl ordering} \\ 1, & \text{Born-Jordan ordering} \\ \delta_{n,0} + \delta_{n,n}, & \text{simple symmetric ordering} \end{cases} \quad (17)$$

¹³Carl M. Bender and Gerald V. Dunne. "Polynomials and Operator Orderings". In: *Journal of Mathematical Physics* 29.8 (Aug. 1988), pp. 1727–1731. ISSN: 0022-2488, 1089-7658. DOI: 10.1063/1.527869; Carl M. Bender and Gerald V. Dunne. "Exact Solutions to Operator Differential Equations". In: *Physical Review D* 40.8 (Oct. 15, 1989), pp. 2739–2742. ISSN: 0556-2821. DOI: 10.1103/PhysRevD.40.2739; Carl M. Bender and Gerald V. Dunne. "Integration of Operator Differential Equations". In: *Physical Review D* 40.10 (Nov. 15, 1989), pp. 3504–3511. ISSN: 0556-2821. DOI: 10.1103/PhysRevD.40.3504.

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

Quantized time of arrival operators

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

In position space, the TOA operators are integral operators¹⁴

$$(\hat{T}\phi)(q) = \frac{\mu}{i\hbar} \int_{-\infty}^{\infty} dq' T(q, q') \text{sgn}(q - q') \phi(q') \quad (18)$$

where,

$$T_W(q, q') = \frac{1}{2} \int_0^{\frac{q+q'}{2}} ds {}_0F_1 \left[; 1; \frac{\mu}{2\hbar^2} (q - q')^2 V \left(\frac{q + q'}{2}, s \right) \right]$$

¹⁴Eric A Galapon and John Jaykel P Magadan. "Quantizations of the classical time of arrival and their dynamics". In: *Annals of Physics* 397 (2018), pp. 278–302.

Quantized time of arrival operators

Introduction

Classical time of arrival

Quantum time of arrival

Time operators

Non-relativistic TOA operators

Free case

Interacting case

Relativistic TOA operators

Motivation

Related studies

Research proposal

Comments

Goals

$$\begin{aligned} T_{BJ}(q, q') = & \frac{1}{2(q - q')} \int_0^q ds \int_0^s du \\ & \times {}_0F_1 \left[; 1; \frac{\mu}{2\hbar^2} (q - q')^2 V(s, u) \right] \\ & - \frac{1}{2(q - q')} \int_0^{q'} ds \int_0^s du \\ & \times {}_0F_1 \left[; 1; \frac{\mu}{2\hbar^2} (q - q')^2 V(s, u) \right] \end{aligned}$$

$$\begin{aligned} T_{SS}(q, q') = & \frac{1}{4} \int_0^q ds {}_0F_1 \left[; 1; \frac{\mu}{2\hbar^2} (q - q')^2 V(q, s) \right] \\ & - \frac{1}{4} \int_0^{q'} ds {}_0F_1 \left[; 1; \frac{\mu}{2\hbar^2} (q - q')^2 V(q', s) \right] \end{aligned}$$

where, $V(x, y) = V(x) - V(y)$.

Supraquantized time of arrival operators

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies


Research proposal

Comments
Goals

Supraquantization

Method of constructing quantum observables without quantization.

- Process of quantization is **circular**



- No quantization exists such that for all classical observables f and g , the **Dirac condition** is satisfied

$$\{f, g\} \Rightarrow [\hat{Q}_f, \hat{Q}_g] = i\hbar$$

Supraquantized time of arrival operators

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

Supraquantized kernel¹⁵

The time kernel function must satisfy the PDE

$$-\frac{\hbar^2}{2\mu} \left(\frac{\partial^2}{\partial q^2} - \frac{\partial^2}{\partial q'^2} \right) T(q, q') + (V(q) - V(q')) T(q, q') = 0 \quad (19)$$

with the boundary conditions $T(q, q) = \frac{q}{2}$ and $T(q, -q) = 0$

- The supraquantized kernel is **unique**, in contrast to the quantized kernel.
- Supraquantization uses the classical observable as a boundary condition.

¹⁵Eric A Galapon. "Shouldn't there be an antithesis to quantization?" In: *Journal of mathematical physics* 45.8 (2004), pp. 3180–3215.

Supraquantized time of arrival operators

Introduction

- Classical time of arrival
- Quantum time of arrival
- Time operators

Non-relativistic TOA operators

- Free case
- Interacting case

Relativistic TOA operators

- Motivation
- Related studies

Research proposal

- Comments
- Goals

Linear systems

For potentials of the form $V(q) = aq^2 + bq + c$, the supraquantized kernel is equal to the Weyl-ordered kernel

Non-linear systems

The supraquantized kernel is equal to the Weyl-ordered kernel plus corrections terms which arise due to the obstruction to quantization.

Why relativistic TOA operators?

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

- If position can be treated as an observable then it is **natural to promote time**, e.g. TOA, as an observable with a corresponding operator¹⁶.
- There are ‘negative energies’ and ‘negative probabilities’

Additional problem

Although the presence of anti-particles naturally bypasses Pauli’s theorem, we are now faced with the problem that we do not know if the particle that arrived is the same particle we started with.

¹⁶ Joseph Bunao and Eric A Galapon. “A one-particle time of arrival operator for a free relativistic spin-0 charged particle in $(1+1)$ dimensions”. In: *Annals of Physics* 353 (2015), pp. 83–106.

Razavi TOA operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

- Earliest construction of a relativistic free TOA operator was done by Razavi¹⁷

Construction

The classical TOA is first calculated

$$t = \left\{ \int dq \left(\left. \frac{\partial H}{\partial p} \right|_{p=p(q,H)} \right)^{-1} \right\}_{H=H(p,q)} \quad (20)$$

then quantized using simple-symmetric quantization.

¹⁷ M Razavy. "Quantum-mechanical conjugate of the hamiltonian operator". In: *Il Nuovo Cimento B* (1965-1970) 63.1 (1969), pp. 271–308.

Razavi TOA operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

Spin-0

The positive-energy solution of the Klein-Gordon equation is generated by $H = \sqrt{p^2 c^2 + \mu^2 c^4}$ which leads to the operator

$$\hat{T}_{\text{Ra}}^{(0)} = \frac{1}{2} \left(\frac{\sqrt{\hat{p}^2 c^2 + \mu^2 c^4}}{\hat{p} c^2} \hat{q} + \hat{q} \frac{\sqrt{\hat{p}^2 c^2 + \mu^2 c^4}}{\hat{p} c^2} \right) \quad (21)$$

- This looks like the relativistic version of the Aharonov-Bohm TOA operator

Razavi TOA operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

Spin-0

The eigenfunction is given as

$$\varphi_{\tau}(p) = N \sqrt{\frac{|p|c}{\sqrt{p^2 c^2 + \mu^2 c^4}}} \exp\left(\frac{i}{\hbar} \tau \sqrt{p^2 c^2 + \mu^2 c^4}\right) \quad (22)$$

- Not square integrable unless τ is complex where $\text{Im}(\tau) < 0$
- If τ is real then the eigenfunction becomes

$$\varphi_{\tau}(p, \epsilon) = \int_{\tau-\epsilon}^{\tau+\epsilon} d\tau' \varphi_{\tau'}(p) \quad (23)$$

Razavi TOA operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

Spin-1/2

Apply a Foldy-Wouthuysen transformation to $H = \alpha \cdot \hat{p} + \beta \mu c^2$ where

$$H \rightarrow \hat{U} H \hat{U}^{-1} = \beta \sqrt{p^2 c^2 + \mu^2 c^4} \quad (24)$$

where, $\hat{U}^{\pm 1} = \cos \theta \pm \beta \alpha \cdot \hat{p} \sin \theta$ such that $\tan 2\theta = \frac{|p|}{\mu c}$.
Since the unitary transform preserves the commutation relation, then

$$\hat{T}_{\text{Ra}}^{(1/2)} = \hat{U} \hat{T}_{\text{Ra}}^{(0)} \hat{U}^{-1} \quad (25)$$

León TOA operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

- Extended the Grot-Rovelli-Tate TOA operator to the positive energy solutions of the Klein-Gordon equation¹⁸

Spin-0

The operator is given as

$$\hat{T}_{\text{Le}} = -\frac{\sqrt{E_p}}{c} \hat{p}^{-1/2} \hat{Q}_{NW} \hat{p}^{1/2} \frac{\sqrt{E_p}}{c} \quad (26)$$

where $E_p = \sqrt{p^2 c^2 + \mu^2 c^4}$ and \hat{Q}_{NW} is the Newton-Wigner position operator¹⁹

¹⁸ Juan León. "Time-of-arrival formalism for the relativistic particle". In: *Journal of Physics A: Mathematical and General* 30.13 (1997), p. 4791.

¹⁹ Theodore Duddell Newton and Eugene P Wigner. "Localized states for elementary systems". In: *Reviews of Modern Physics* 21.3 (1949), p. 400.

León TOA operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

- The Newton-Wigner position operator is

$$\hat{Q}_{NW} = \hat{x} - i\hbar \frac{pc^2}{2E_p^2} \quad (27)$$

- By construction, its eigenfunctions are **not Lorentz covariant**
- The localized state in position space is **not a Dirac delta**

Wang-Xiong TOA operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

- Direct extension of the non-relativistic TOA operator²⁰

$$t = -\mu \frac{q}{p} \Rightarrow -\frac{E_p}{c^2} \frac{q}{p} \quad (28)$$

Spin-1/2

Total symmetrization of the TOA yields the operator

$$\hat{T}_{WX} = -\frac{1}{4c^2}(\hat{H}\hat{T}_{AB} + \hat{T}_{AB}\hat{H}) = -\alpha_1 \frac{\hat{q}}{c^2} + \beta \hat{T}_{AB} \quad (29)$$

where $\hat{H} = \alpha_1 \hat{p} + \beta \mu c^2$

²⁰Zhi-Yong Wang and Cai-Dong Xiong. "Relativistic free-motion time-of-arrival". In: *Journal of Physics A: Mathematical and Theoretical* 40.8 (2007), p. 1897.

Wang-Xiong TOA operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

- The proper-time eigenfunctions are solved by

$$\hat{T}_{WX}\varphi(p) = t\varphi(p) \quad (30)$$

$$t = -\lambda \frac{E_p}{pc^2} q \quad \text{and} \quad \tau = \mu \frac{q}{p} \quad (31)$$

which yields

$$\varphi(p) = N \left(\frac{q^2/c^2}{q^2/c^2 + \tau^2} \right)^{1/4} \xi(x) e^{-ipx/\hbar} \quad (32)$$

$$\xi(p) = \sqrt{\frac{\tau + bt_x}{2bt_x}} \left[\frac{\eta_s}{\sigma_1 x/c^2} \right]_{\tau + bt_x} \eta_s \quad (33)$$

where $b = \pm 1$ and $t_x = \sqrt{x^2/c^2 + \tau^2}$

Wang-Xiong TOA operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

■ Dual relations

- $t^2 = q^2/c^2 + \tau^2 \leftrightarrow E_p^2 = p^2 c^2 + \mu^2 c^4$
- $\hat{T}_{WX} = -\alpha_1 \hat{q}/c^2 + \beta \hat{T}_{AB} \leftrightarrow \hat{H} = \alpha_1 \hat{p} + \beta \mu c^2$
- $-i\hbar \partial_E \varphi = \hat{T}_{WX} \varphi \leftrightarrow i\hbar \partial_t \psi = \hat{H} \psi$

■ They have also applied this operator to quantum field theory²¹

²¹ ZY Wang, B Chen, and CD Xiong. "Time in quantum mechanics and quantum field theory". In: *Journal of Physics A: Mathematical and General* 36.18 (2003), p. 5135; Zhi-Yong Wang, Cai-Dong Xiong, and Bing He. "Arrival time in quantum field theory". In: *Physics Letters B* 666.4 (2008), pp. 382–385; Zhi-Yong Wang, Qi Qiu, and Cai-Dong Xiong. "Time operator in QFT with Virasoro constraints". In: *Physics Letters B* 718.4-5 (2013), pp. 1515–1518.

Bauer-Aguillon TOA operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

- Applies Lorentz and Born's reciprocity^{22,23} invariance to the canonical quantization of special relativity

- $p_\mu p^\mu = p_o^2 - \vec{p}^2 = (m_o c)^2$ leads to

$$(\hat{p}_\mu \hat{p}^\mu - (m_o c)^2) |\psi\rangle = 0 \Rightarrow c \hat{p}_o |\psi\rangle = (c\alpha \cdot \hat{\mathbf{p}} + \beta m_o c^2) |\psi\rangle$$

provided $[\hat{p}_\mu, \hat{p}_\nu] = 0$

- $x_\mu x^\mu = x_o^2 - \vec{r}^2 = s_o^2$ leads to

$$(\hat{x}_\mu \hat{x}^\mu - s_o^2) |\psi\rangle = 0 \Rightarrow \frac{\hat{x}_o}{c} |\psi\rangle = \left(\frac{\alpha \cdot \hat{\mathbf{r}}}{c} + \beta \tau_o \right) |\psi\rangle$$

provided $[\hat{x}_\mu, \hat{x}_\nu] = 0$

- $\mathcal{O}^- = x_\mu p^\mu - p^\mu x_\mu \rightarrow [\hat{x}_\mu, \hat{p}_\nu] = i\hbar \eta_{\mu\nu} \infty$

²²Max Born. "A suggestion for unifying quantum theory and relativity". In: *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences* 165.921 (1938), pp. 291–303;
Max Born. "Reciprocity theory of elementary particles". In: *Reviews of Modern Physics* 21.3 (1949), p. 463.

²³The transformation $x \rightarrow p$ and $p \rightarrow -x$ leaves the Hamilton equations invariant

Bauer-Aguillon TOA operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

Spin-1/2

The time operator is

$$\hat{T}_{BA} = \frac{\alpha \cdot \hat{\mathbf{r}}}{c} + \beta \tau_0 \quad (34)$$

where τ_0 is a real-valued constant²⁴

- This time operator is not canonically conjugate to the Hamiltonian.

²⁴ **Mariano Bauer**. "A dynamical time operator in Dirac's relativistic quantum mechanics". In: *International Journal of Modern Physics A* 29.06 (2014), p. 1450036; **M Bauer**. "A time operator in the simulations of the Dirac equation". In: *International Journal of Modern Physics A* 34.22 (2019), p. 1950114; **CA Aguillón, M Bauer, and GE García**. "Time and energy operators in the canonical quantization of special relativity". In: *European Journal of Physics* 41.3 (2020), p. 035601; **M Bauer, CA Aguillón, and GE García**. "Conditional interpretation of time in quantum gravity and a time operator in relativistic quantum mechanics". In: *International Journal of Modern Physics A* 35.21 (2020), p. 2050114.

Bunao-Galapon one-particle TOA operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

- A TOA operator was constructed by solving the commutation relation $[\hat{H}, \hat{T}] = i\hbar$
- The one-particle TOA operator was constructed by taking transform $\hat{T}_\Phi = \hat{U}\hat{T}\hat{U}^{-1}$ where²⁵

$$\hat{U}^\pm = \frac{(\mu c^2 + E_p)\hat{\sigma}_0 \mp (\mu c^2 - E_p)\hat{\sigma}_1}{\sqrt{4\mu c^2 E_p}} \quad (35)$$

²⁵ Herman Feshbach and Felix Villars. "Elementary relativistic wave mechanics of spin 0 and spin 1/2 particles". In: *Reviews of Modern Physics* 30.1 (1958), p. 24; Walter Greiner et al. *Relativistic quantum mechanics*. Vol. 2. Springer, 2000.

Bunao-Galapon one-particle TOA operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

Spin-0

The one particle operator is²⁶

$$\hat{T}_{BG}^{(\Phi)} = \hat{\sigma}_3 \left[-\frac{1}{2E_p} \left(\frac{\hat{p}\hat{q} + \hat{q}\hat{p}}{2} \right) - \left(\frac{\mu^2 c^2}{E_p} + \frac{p^2}{2E_p} \right) \left(\frac{\hat{p}^{-1}\hat{q} + \hat{q}\hat{p}^{-1}}{2} \right) \right] \quad (36)$$

- Reduces to the $\hat{\sigma}_3 \hat{T}_{AB}$ as $c \rightarrow \infty$
- Not self-adjoint but maximally symmetric

²⁶ Joseph Bunao and Eric A Galapon. "A one-particle time of arrival operator for a free relativistic spin-0 charged particle in (1+ 1) dimensions". In: *Annals of Physics* 353 (2015), pp. 83–106.

Bunao-Galapon one-particle TOA operator

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

Spin-1/2

The one particle operator is²⁷ the same as that for the spin-0

- There is an extra term but it was dropped to impose **parity inversion symmetry** to the operator
- The extra term **commutes** with the Hamiltonian

²⁷ Joseph Bunao and Eric A Galapon. "A relativistic one-particle Time of Arrival operator for a free spin-1/2 particle in (1+ 1) dimensions". In: *Annals of Physics* 356 (2015), pp. 369–382.

Comments

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

- There is still **no consensus** on the status of time in non-relativistic quantum mechanics
- **Additional problems** arise because of special relativity
 - Lorentz invariance: Hamiltonian formulation of only works in some specific Lorentz frame and the TOA is also measured in that frame²⁸
 - Identity of the particle upon arrival
 - Klein-Gordon and Dirac equation cannot provide a well-defined local probability distribution²⁹

²⁸ Herbert Goldstein, Charles Poole, and John Safko. *Classical mechanics*. 2002.

²⁹ AJ Kálnay. *The localization problem*. in "Problems in the Foundations of Physics"(M. Bunpe. Ed.). Vol. 4. pp. 93-J 10. 1971; Donald Reed. "Lawrence P. Horwitz: Relativistic Quantum Mechanics". In: *Foundations of Physics* 47.11 (2017), pp. 1498–1502; Theodore Duddell Newton and Eugene P Wigner. "Localized states for elementary systems". In: *Reviews of Modern Physics* 21.3 (1949), p. 400.

Problem

Introduction

Classical time of arrival
Quantum time of arrival
Time operators

Non-relativistic TOA operators

Free case
Interacting case

Relativistic TOA operators

Motivation
Related studies

Research proposal

Comments
Goals

- Given a relativistic particle with a certain initial state, what is the TOA probability distribution?
- Does this TOA distribution have an underlying ideal distribution generated by a corresponding TOA operator?
- How do we construct the TOA operator?



Thank You