

#10: DIFFERENTIAL EQUATIONS

14. A large spherical balloon is deflating.

At time t seconds the balloon has radius r cm and volume V cm³

The volume of the balloon is modelled as decreasing at a constant rate.

(a) Using this model, show that

$$\frac{dr}{dt} = -\frac{k}{r^2}$$

where k is a positive constant.

(3)

Given that

- the initial radius of the balloon is 40 cm
- after 5 seconds the radius of the balloon is 20 cm
- the volume of the balloon continues to decrease at a constant rate until the balloon is empty

(b) solve the differential equation to find a complete equation linking r and t .

(5)

(c) Find the limitation on the values of t for which the equation in part (b) is valid.

(2)

a) $V = \frac{4}{3}\pi r^3$

$$\frac{dV}{dr} = 4\pi r^2$$

$$\begin{aligned} \frac{dr}{dt} &= \frac{dr}{dV} \times \frac{dV}{dt} \\ &= \frac{1}{4\pi r^2} \times -c \end{aligned}$$

$$= -\frac{c}{4\pi} \times \frac{1}{r^2} \quad \text{Let } k = \frac{c}{4\pi}$$

$$= -\frac{k}{r^2}$$

2020 PAPER 1
23.6%

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 14 continued

$$b) \frac{dr}{dt} = -\frac{k}{r^2}$$

$\leftarrow r^2$

$$r^2 dr = -k dt$$

$$\int r^2 dr = \int -k dt$$

$$\frac{1}{3}r^3 = -kt + c$$

$$\text{When } t=0, r=40 \rightarrow \frac{1}{3}(40)^3 = c \quad c = \frac{64000}{3}$$

$$t=5, r=20 \rightarrow \frac{1}{3}(20)^3 = -5k + \frac{64000}{3}$$

$$k = \frac{11200}{3}$$

$$\text{Hence } \frac{1}{3}r^3 = -\frac{11200}{3}t + \frac{64000}{3}$$

$$r^3 = 64000 - 11200t$$

$$c) r>0, \text{ hence } 64000 - 11200t > 0$$

$$\begin{aligned} \frac{64000}{11200} &> t \\ \frac{40}{7} &> t \end{aligned}$$

$$\text{Hence } 0 \leq t < \frac{40}{7}$$



#9: DIFFERENTIATION

13. The function g is defined by

$$g(x) = \frac{3\ln(x) - 7}{\ln(x) - 2} \quad x > 0 \quad x \neq k$$

where k is a constant.

(a) Deduce the value of k .

(1)

(b) Prove that

$$g'(x) > 0$$

for all values of x in the domain of g .

(3)

(c) Find the range of values of a for which

$$g(a) > 0$$

(2)

a) $\ln k - 2 = 0$
 $\ln k = 2$
 $k = e^2$

b) Quotient Rule

$$u = 3\ln x - 7 \quad v = \ln x - 2$$

$$u' = \frac{3}{x} \quad v' = \frac{1}{x}$$

$$g'(x) = \frac{vu' - uv'}{v^2}$$

$$g'(x) = \frac{\frac{3}{x}(\ln x - 2) - \frac{1}{x}(3\ln x - 7)}{(\ln x - 2)^2}$$

$$= \frac{\frac{3}{x}\ln x - \frac{6}{x} - \frac{3}{x}\ln x + \frac{7}{x}}{(\ln x - 2)^2}$$

$$= \frac{\frac{1}{x}}{(\ln x - 2)^2}$$

As $x > 0$, $\frac{1}{x} > 0$, and $\ln x - 2$ is squared, it is also ≥ 0 ,

hence $g'(x) > 0$

2020 PAPER 2
23.3%

Question 13 continued

c) $g(a) > 0$

$$\frac{3\ln a - 7}{\ln a - 2} > 0$$

$$3\ln a - 7 > 0$$

$$\ln a > \frac{7}{3}$$

$$a > e^{\frac{7}{3}}$$



9.




Figure 4




Figure 5

Figure 4 shows a sketch of a Ferris wheel.

The height above the ground, H m, of a passenger on the Ferris wheel, t seconds after the wheel starts turning, is modelled by the equation

$$H = |A \sin(bt + \alpha)|$$

where A , b and α are constants.

Figure 5 shows a sketch of the graph of H against t , for one revolution of the wheel.

Given that

- the maximum height of the passenger above the ground is 50 m
- the passenger is 1 m above the ground when the wheel starts turning
- the wheel takes 720 seconds to complete one revolution

- (a) find a complete equation for the model, giving the exact value of A , the exact value of b and the value of α to 3 significant figures. (4)

- (b) Explain why an equation of the form

$$H = |A \sin(bt + a)| + d$$

where d is a positive constant, would be a more appropriate model. (1)

**#8: MODELLING
WITH TRIGONOMETRY**

2022 PAPER 2
22.4%

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 9 continued

$$H = |A \sin(bt + \alpha)|$$


shifts α to the left
 $f(x+\alpha)$

scales the amplitude
 $af(x)$

stretches by factor $\frac{1}{b}$
 $f(bx)$

$$A = 50 \quad (\text{max height})$$

720 for one revolution



When $t=0$, $H=1$

$$1 = |50 \sin(\frac{1}{4}t + \alpha)|$$

$$1 = 50 \sin \alpha$$

$$\alpha = 1.15 \quad (3sf)$$

$$\text{Hence } H = |50 \sin(\frac{1}{4}t + 1.15)|$$

b) In this second model the passenger now does not touch the ground which is more realistic

(Total for Question 9 is 5 marks)



#7: IMPLICIT DIFFERENTIATION, POINTS OF INFLECTION

15. The curve C has equation

$$x^2 \tan y = 9 \quad 0 < y < \frac{\pi}{2}$$

2020 PAPER 1
21.7%

(a) Show that

$$\frac{dy}{dx} = \frac{-18x}{x^4 + 81} \quad (4)$$

(b) Prove that C has a point of inflection at $x = \sqrt[4]{27}$

Product Rule

$$a) \quad x^2 \tan y = 9 \quad u = x^2 \quad v = \tan y$$

$$u' = 2x \quad v' = \sec^2 y \quad \frac{dy}{dx}$$

$$x^2 \sec^2 y \frac{dy}{dx} + 2x \tan y = 0$$

$$\frac{dy}{dx} = \frac{-2x \tan y}{x^2 \sec^2 y}$$

$$\tan y = \frac{9}{x^2}$$

$$\frac{dy}{dx} = \frac{-2x \times \frac{9}{x^2}}{x^2 \left(1 + \frac{81}{x^4}\right)}$$

$$1 + \tan^2 y = \sec^2 y$$

$$1 + \left(\frac{9}{x^2}\right)^2 = \sec^2 y$$

$$1 + \frac{81}{x^4} = \sec^2 y$$

$$\frac{dy}{dx} = \frac{-\frac{18}{x}}{x^2 + \frac{81}{x^2}}$$

$$\frac{dy}{dx} = \frac{-18x}{x^4 + 81}$$

Question 15 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

b) Quotient Rule

$$u = -18x \quad v = x^4 + 81$$

$$u' = -18 \quad v' = 4x^3$$

$$\frac{d^2y}{dx^2} = \frac{-18(x^4 + 81) + 18x(4x^3)}{(x^4 + 81)^2}$$

$$= \frac{-18x^4 - 1458 + 72x^4}{(x^4 + 81)^2}$$

$$\frac{d^2y}{dx^2} = \frac{54x^4 - 1458}{(x^4 + 81)^2}$$

Point of inflection if: $\frac{d^2y}{dx^2} = 0$

and there is a change of sign of $\frac{d^2y}{dx^2}$ either side of that solution

$$\frac{d^2y}{dx^2} = 0$$

$$54x^4 - 1458 = 0$$

$$x^4 = 27$$

$$x = \sqrt[4]{27}$$



When $x < \sqrt[4]{27}$, $\frac{d^2y}{dx^2} < 0$ and when $x > \sqrt[4]{27}$, $\frac{d^2y}{dx^2} > 0$

hence there is a point of inflection when $x = \sqrt[4]{27}$



#6: PARAMETRIC INTEGRATION


2022 PAPER 1
21.6%

Figure 6

Figure 6 shows a sketch of the curve C with parametric equations

$$x = 8 \sin^2 t \quad y = 2 \sin 2t + 3 \sin t \quad 0 \leq t \leq \frac{\pi}{2}$$

The region R , shown shaded in Figure 6, is bounded by C , the x -axis and the line with equation $x = 4$

(a) Show that the area of R is given by

$$\int_0^a (8 - 8 \cos 4t + 48 \sin^2 t \cos t) dt \quad (5)$$

where a is a constant to be found.

(b) Hence, using algebraic integration, find the exact area of R . (4)

a) Find a

$$\begin{aligned} x &= 4 \\ 4 &= 8 \sin^2 t \\ \frac{1}{2} &= \sin^2 t \\ \pm \frac{\sqrt{2}}{2} &= \sin t \quad (\text{But } 0 \leq t \leq \frac{\pi}{2}, \text{ so } \sin t = \frac{\sqrt{2}}{2}) \end{aligned}$$

$$t = \frac{\pi}{4}$$

$$\text{Hence } a = \frac{\pi}{4}$$



Question 16 continued

$$\begin{aligned} x &= 8(\sin t)^2 \\ \frac{dx}{dt} &= 16 \sin t \cos t \end{aligned}$$

$$\begin{aligned} R &= \int_0^{\frac{\pi}{4}} y \frac{dx}{dt} dt = \int_0^{\frac{\pi}{4}} (2\sin 2t + 3\sin t) 16 \sin t \cos t dt \\ &= \int_0^{\frac{\pi}{4}} (32\sin 2t \sin t \cos t + 48\sin^2 t \cos t) dt \\ \sin 2t &= 2\sin t \cos t \\ 2\sin^2 t &= 1 - \cos^2 t \\ &= \int_0^{\frac{\pi}{4}} (16 \sin^2 t \sin 2t + 48\sin^2 t \cos t) dt \\ &= \int_0^{\frac{\pi}{4}} (8(1 - \cos 4t) + 48\sin^2 t \cos t) dt \\ &= \int_0^{\frac{\pi}{4}} (8 - 8\cos 4t + 48\sin^2 t \cos t) dt \end{aligned}$$

$$b) \quad R = \left[8t - 2\sin 4t + 16 \sin^3 t \right]_0^{\frac{\pi}{4}}$$

$$\begin{aligned} &= 8\left(\frac{\pi}{4}\right) - 2\sin \pi + 16 \sin^3\left(\frac{\pi}{4}\right) - 0 - 0 - 0 \\ &= 2\pi - 0 + 16\left(\frac{\sqrt{2}}{2}\right)^3 \\ &= 2\pi + 4\sqrt{2} \end{aligned}$$

DO NOT WRITE IN THIS AREA



#5: CIRCLES

14. A circle C with radius r

- lies only in the 1st quadrant
- touches the x -axis and touches the y -axis


The line l has equation $2x + y = 12$

(a) Show that the x coordinates of the points of intersection of l with C satisfy

$$5x^2 + (2r - 48)x + (r^2 - 24r + 144) = 0 \quad (3)$$

Given also that l is a tangent to C ,

(b) find the two possible values of r , giving your answers as fully simplified surds.



Intersection with $2x + y = 12$

$$y = 12 - 2x$$

$$(x-r)^2 + (12-2x-r)^2 = r^2$$

$$\begin{array}{r} 12 & -2x & -r \\ \hline 144 & -24x & -12r \\ -2x & -24x & +4x^2 + 2xr \\ -r & -12r & +2xr + r^2 \end{array}$$

$$x^2 - 2xr + r^2 + 144 - 48x - 24r + 4x^2 + 4xr + r^2 = r^2$$

$$5x^2 + 2xr - 48x + r^2 - 24r + 144 = 0$$

$$5x^2 + (2r - 48)x + (r^2 - 24r + 144) = 0$$

b) tangent, $b^2 - 4ac = 0$

$$a=5, b=2r-48, c=r^2-24r+144$$

$$(2r-48)^2 - 4(5)(r^2 - 24r + 144) = 0$$

$$4r^2 - 192r + 2304 - 20r^2 + 480r - 2880 = 0$$

$$-16r^2 + 288r - 576 = 0$$

$$r = 9 + 3\sqrt{5}, r = 9 - 3\sqrt{5}$$

2020 PAPER 2
20.0%

Question 14 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

#4: PARAMETRIC EQUATIONS

13. A curve C has parametric equations

$$x = \frac{t^2 + 5}{t^2 + 1} \quad y = \frac{4t}{t^2 + 1} \quad t \in \mathbb{R}$$

Show that all points on C satisfy

$$(x - 3)^2 + y^2 = 4$$

**2021 PAPER 1
19.7%**

(3)

$$x - 3 = \frac{t^2 + 5}{t^2 + 1} - \frac{3(t^2 + 1)}{t^2 + 1}$$

$$= \frac{t^2 + 5 - 3t^2 - 3}{t^2 + 1}$$

$$= \frac{2 - 2t^2}{t^2 + 1}$$

$$(2 - 2t^2)^2 =$$

$$(x - 3)^2 = \frac{4 - 8t^2 + 4t^4}{(t^2 + 1)^2}$$

$$y^2 = \frac{16t^2}{(t^2 + 1)^2}$$

$$\begin{aligned} \text{Hence } (x - 3)^2 + y^2 &= \frac{4 - 8t^2 + 4t^4}{(t^2 + 1)^2} + \frac{16t^2}{(t^2 + 1)^2} \\ &= \frac{4t^4 + 8t^2 + 4}{(t^2 + 1)^2} \\ &= \frac{4(t^4 + 2t^2 + 1)}{(t^2 + 1)^2} = \frac{4(t^2 + 1)^2}{(t^2 + 1)^2} \\ &= 4 \end{aligned}$$

Question 13 continued

DO NOT WRITE IN THIS AREA

(Total for Question 13 is 3 marks)



#3: DIFFERENTIAL EQUATIONS

2021 PAPER 2
14.3%




Figure 5

Water flows at a constant rate into a large tank.

The tank is a cuboid, with all sides of negligible thickness.

The base of the tank measures 8 m by 3 m and the height of the tank is 5 m.

There is a tap at a point T at the bottom of the tank, as shown in Figure 5.

At time t minutes after the tap has been opened

- the depth of water in the tank is h metres
- water is flowing into the tank at a constant rate of 0.48 m^3 per minute
- water is modelled as leaving the tank through the tap at a rate of $0.1h \text{ m}^3$ per minute

(a) Show that, according to the model,

$$1200 \frac{dh}{dt} = 24 - 5h \quad (4)$$

$$\begin{aligned} \frac{dV}{dt} &= 0.48 \\ \frac{dV}{dt} &= 0.48 - 0.1h \\ \frac{dh}{dt} &= -0.1h \end{aligned}$$

Given that when the tap was opened, the depth of water in the tank was 2 m,

(b) show that, according to the model,

$$h = A + Be^{-kt} \quad (6)$$

where A , B and k are constants to be found.

Given that the tap remains open,

(c) determine, according to the model, whether the tank will ever become full, giving a reason for your answer.

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 14 continued

$$\begin{aligned} V &= 8 \times 3 \times h \\ V &= 24h \end{aligned}$$

$$\frac{dV}{dh} = 24$$

$$\frac{dh}{dt} = \frac{dV}{dt} = \frac{dV}{dh} \frac{dh}{dt}$$

$$\frac{dh}{dt} = \frac{1}{24} (0.48 - 0.1h)$$

$$24 \frac{dh}{dt} = 0.48 - 0.1h$$

$\downarrow \times 50$

$$1200 \frac{dh}{dt} = 24 - 5h$$

$$b) \frac{1200}{24-5h} \frac{dh}{dt} = 1$$

$$\int \frac{1200}{24-5h} dh = \int 1 dt$$

$$-240 \ln |24-5h| = t + C$$

$$\begin{aligned} \text{When } t=0, h &= 2 \\ -240 \ln 14 &= C \end{aligned}$$

$$\text{Hence } -240 \ln |24-5h| = t - 240 \ln 14$$

$$240(\ln 14 - \ln |24-5h|) = t$$

$$\ln \left(\frac{14}{24-5h} \right) = \frac{t}{240}$$

$$\frac{14}{24-5h} = e^{t/240}$$

$$\frac{24-5h}{14} = e^{-t/240}$$

$$24 - 14e^{-t/240} = 5h$$

$$h = 4.8 - 2.8e^{-t/240}$$

$$c) \text{ As } t \rightarrow \infty \\ -2.8e^{-t/240} \rightarrow 0$$

So $h \rightarrow 4.8$
This is less than
5m so
never fills up.

16. Use algebra to prove that the square of any natural number is either a multiple of 3 or one more than a multiple of 3

(4)

#2: PROOF

2020 PAPER 2

6.0%

Cannot exhaust using evens and odds...

Instead:

$$\begin{aligned}3k &\cdot \text{multiples of 3} & 24 \\3k+1 &\cdot \text{one more than a multiple of 3} & 31 \\3k-1 &\cdot \text{one less than a multiple of 3} & 17\end{aligned}$$

So if k is a positive integer...

$$\begin{aligned}(3k)^2 &= 9k^2 \\&= 3(3k^2) \text{ so is a multiple of 3}\end{aligned}$$

$$\begin{aligned}(3k+1)^2 &= 9k^2 + 6k + 1 \\&= 3(3k^2 + 2k) + 1 \text{ so is one more than a multiple of 3}\end{aligned}$$

$$\begin{aligned}(3k-1)^2 &= 9k^2 - 6k + 1 \\&= 3(3k^2 - 2k) + 1 \text{ so is one more than a multiple of 3}\end{aligned}$$

Hence we have proven the statement.

Question 16 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



16. Prove by contradiction that there are no positive integers p and q such that

#1: PROOF BY CONTRADICTION

2020 PAPER 1

4.3%

Assume, for contradiction, that there are positive integers p and q such that

$$4p^2 - q^2 = 25 \quad (4)$$

$$\begin{array}{l} 4p^2 - q^2 = 25 \\ \text{D.O.T.S.} \end{array}$$

$$(2p+q)(2p-q) = 25$$

$$\begin{array}{ll} 5 \times 5 = 25 & \text{are the only integer factor} \\ 25 \times 1 = 25 & \text{pairs for 25} \end{array}$$

$$\text{Hence } 2p+q = 5$$

$$2p-q = 5$$

$$p = 2.5, q = 0$$

Neither p and q are positive integers

$$\text{or } 2p+q = 25$$

$$2p-q = 1$$

$$p = 6.5, q = 12$$

p is not an integer.

Hence we have contradicted our assumption, and

so there are no positive integers p and q such that

$$4p^2 - q^2 = 25$$

Question 16 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

