



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

# Aiming for A\*: ‘in between’ Pure 1 and Pure 2 [Edexcel]

Bicen Maths

**Note:**

- I have used questions from the international A-Level, old specification A-Level, and other exam boards – I only used ‘recent’ years of the old specification, as this is when examiners were gearing up for the new style of questions, so they’re a good fit!
- This is to ensure the questions are all ‘fresh’, as at this stage I know that exam question fatigue is real.
- I’ve used my knowledge to select ones that match Edexcel’s style as much as possible.



- 1 (a) For a small angle  $\theta$ , where  $\theta$  is in radians, show that  $2\cos\theta + (1 - \tan\theta)^2 \approx 3 - 2\theta$ . [3]
- (b) Hence determine an approximate solution to  $2\cos\theta + (1 - \tan\theta)^2 = 28\sin\theta$ . [2]

## OCR Pure/Mech 3 2020



- 2 A sequence of transformations maps the curve  $y = e^x$  to the curve  $y = e^{2x+3}$ .  
Give details of these transformations. [3]

## OCR Pure 1 2018



- 4 Prove algebraically that  $n^3 + 3n - 1$  is odd for all positive integers  $n$ . [4]

## OCR MEI Pure 1 2021



- 1 Beth states that for all real numbers  $p$  and  $q$ , if  $p^2 > q^2$  then  $p > q$ .  
Prove that Beth is **not** correct. [2]

DO NOT WRITE IN THIS AREA

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



1	(a)	$2\left(1 - \frac{1}{2}\theta^2\right) + (1 - \theta)^2$ $2 - \theta^2 + 1 - 2\theta + \theta^2$ $= 3 - 2\theta \quad \text{A.G.}$	B1   [3]	2.1	Correct statement
	(b)	$3 - 2\theta = 28\theta$ $\theta = 0.1$	M1   [2]	1.1a A1	Attempt to expand and simplify given expression Obtain given answer Use $28\sin\theta \approx 28\theta$ and attempt to solve Obtain 0.1 oe

2		Refers to translation and stretch  State translation $\begin{pmatrix} -3 \\ 0 \end{pmatrix}$  State stretch by scale factor 0.5 parallel to the $x$ -axis	M1   A1   A1  [3]	1.1   1.1   1.1	In either order; ignore details here; allow any equivalent wording (such as move or shift for translation) to describe geometrical transformations but not statements such as add $-3$ to $x$ (do not accept 'enlargement' or 'shear' for stretch) Or state translation in $x$ -direction by $-3$ (units); accept horizontal to indicate direction or parallel to the $x$ -axis; term 'translate' or 'translation' needed for award of A1 Or in the $x$ direction or horizontally; term 'stretch' needed for award of A1; these two transformations must be in this order – if details correct for M1A1A1 but order wrong, award M1A1A0
---	--	---	---	-----------------------------------	---

<b>4</b>		If $n$ is even then $n$ can be written as $2m$ . $n^3 + 3n - 1 = 8m^3 + 6m - 1$	<b>E1</b>	<b>2.1</b>	Consider when $n$ is even
		$= 2(4m^3 + 3m) - 1$ For all $m$ , $2(4m^3 + 3m)$ is even, hence $2(4m^3 + 3m) - 1$ is odd	<b>E1</b>	<b>2.4</b>	Conclude from useable form
		If $n$ is odd then $n$ can be written as $2m + 1$ $n^3 + 3n - 1 = 8m^3 + 12m^2 + 6m + 1 + 6m + 3 - 1$ $= 8m^3 + 12m^2 + 12m + 3$	<b>E1</b>	<b>2.1</b>	Consider when $n$ is odd
		$= 2(4m^3 + 6m^2 + 6m) + 3$ For all $m$ , $2(4m^3 + 6m^2 + 6m)$ is even, hence $2(4m^3 + 6m^2 + 6m) + 3$ is odd	<b>E1</b>	<b>2.4</b>	Conclude from useable form
<b>1</b>		For example $(-3)^2 = 9 > 2^2 = 4$ and $(-3) < 2$ So Beth is not correct		<b>M1</b>	
				<b>E1</b>	[2]



# What came up in Paper 1?

- factor theorem
- binomial expansion
- numerical methods, Newton-Raphson
- differentiation first principles
- quotient rule/decreasing functions
- modulus graphs
- simple differential equation (tank)
- functions - composite, inverse, domain, range
- geometric series (included indices)
- area integration/differentiation problem
- sector/segment areas
- harmonic identity ( $R\cos \theta R\sin \theta$ )
- integration by substitution
- differential equation - set up and solve (balloon)
- proof, inc. proof by contradiction

## Key

- Doesn't usually appear in both
- Could potentially appear again
- Could easily appear again

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



# What should come up in Paper 2?

- Small angle approximations
- Graph transformations
- Proof (again!)
- Sequences + series – arithmetic and/or sigma notation
- Circles
- Vectors
- Trigonometric identities equations
- Implicit differentiation,  $dx/dy$
- Connected rates of change + optimisation problems
- More integration - limit of a sum, by parts, partial fractions
- Parametric differentiation + integration
- Exponential + logarithmic equations
- Modelling - quadratics, linear, exp & logs
- Trapezium rule
- Inequalities? Regions?
- Numerical methods?
- Surds?
- Pseudo-quadratics?

## Key

- Not covered in this session
- Covered in the 'starter' – use textbook and Exam Qs PDF for more practice questions
- Covered in the main session or Your Turn questions

After the session, make sure you:

- Review anything you didn't understand
- Complete the Your Turn questions
- Review the red and blue topics
- Look over your notes for topics which have already come up, just in case

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



# Sequences and Series

- They usually have at least 2 questions on this topic

- $S_n = \frac{n}{2}(2a + (n - 1)d)$

- $u_n = a + (n - 1)d$

- $S_n = \frac{a(1-r^n)}{1-r}$

- $u_n = ar^{n-1}$

## Arithmetic series

$$S_n = \frac{1}{2} n(a + l) = \frac{1}{2} n[2a + (n - 1)d]$$

## Geometric series

$$S_n = \frac{a(1 - r^n)}{1 - r}$$

$$S_{\infty} = \frac{a}{1 - r} \text{ for } |r| < 1$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



- 3 An arithmetic progression has first term 2 and common difference  $d$ , where  $d \neq 0$ . The first, third and thirteenth terms of this progression are also the first, second and third terms, respectively, of a geometric progression.

By determining  $d$ , show that the arithmetic progression is an increasing sequence.

[5]

DO NOT WRITE IN THIS AREA



- 3 A particular phone battery will last 10 hours when it is first used. Every time it is recharged, it will only last 98% of its previous time.

Find the maximum total length of use for the battery.

[3]

DO NOT WRITE IN THIS AREA

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



3	$2+2d=2r$ $2+12d=2r^2$ $1+6d=(1+d)^2 \text{ or } 2+12d=2(1+d)^2$ $d^2 - 4d = 0 \Rightarrow d = \dots$ <p><math>d = 4</math> and as the common difference is positive the progression is an increasing sequence</p>	<b>B1</b> <b>B1</b> <b>M1*</b> <b>M1dep*</b> <b>A1</b> <b>[5]</b>	<b>1.1</b> <b>1.1</b> <b>1.1</b> <b>1.1</b> <b>2.4</b>	Or for $a + 2d = ar$ Or for $a + 12d = ar^2$ Setting up an equation in $d$ or $r$ only – dependent on one <b>B</b> mark Solving their two-term quadratic equation in $d$ (or three-term quadratic in $r$ ) Correct value for $d$ and link to increasing sequence – must either say that $d$ is positive (oe) or state at least the correct first four terms and comment that they are increasing	$2+12(r-1)=2r^2$ $r^2 - 6r + 5 = 0$ $(r-5)(r-1) = 0$ $\Rightarrow r = \dots$ Condone no mention of $d \neq 0$
---	--	--	--	--	---

3	$10 + 10 \times 0.98 + 10 \times 0.98^2 \text{ or } 10 + 9.8 + 9.604$ $\frac{10}{1-0.98}$ $500$	<b>M1</b> <b>M1</b> <b>A1</b> <b>[3]</b>	<b>3.1b</b> <b>1.1</b> <b>3.2a</b>	Use of GP with common ratio 0.98 Sum to infinity
---	---	---	--	---



2. A sequence  $u_1, u_2, u_3 \dots$  is defined by

$$u_1 = 20$$

$$u_{n+1} = u_n + 5 \sin\left(\frac{n\pi}{2}\right) - 3(-1)^n$$

- (a) (i) Show that  $u_2 = 28$   
(ii) Find the value of  $u_3$  and the value of  $u_4$

(3)

Given that the sequence is periodic with order 4

- (b) (i) write down the value of  $u_5$

(ii) find the value of  $\sum_{r=1}^{25} u_r$

(3)

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Question	Scheme	Marks	AOs
2(a)(i) (ii)	e.g. $(u_2 =) 20 + 5\sin\left(\frac{\pi}{2}\right) - 3(-1)^1 = 28 *$	B1*	2.1
	$u_3 = 28 + 5\sin\left(\frac{2\pi}{2}\right) - 3(-1)^2 (= 25)$ or $u_4 = "25" + 5\sin\left(\frac{3\pi}{2}\right) - 3(-1)^3 (= 23)$	M1	1.1b
	$u_3 = 25$ and $u_4 = 23$	A1	1.1b
		(3)	
(b)(i) (ii)	$(u_5 =) 20$	B1	2.2a
	e.g. $\sum_{r=1}^{25} u_r = 6(20 + 28 + "25" + "23") + 20$	M1	3.1a
	$= 596$	A1	1.1b
		(3)	
(6 marks)			



7. A sequence is defined by

$$\begin{aligned} u_1 &= 3 \\ u_{n+1} &= u_n - 5, \quad n \geq 1 \end{aligned}$$

Find the values of

(a)  $u_2, u_3$  and  $u_4$

(2)

(b)  $u_{100}$

(3)

(c)  $\sum_{i=1}^{100} u_i$

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



7.

(a)  $u_2 = -2, u_3 = -7 \text{ and } u_4 = -12$

M1, A1 [2]

(b)  $d = -5$  and arithmetic

B1  
M1, A1 [3]Uses  $a + (n - 1)d$  with  $a = 3$  and  $n = 100$ , to give  $-492$ 

(c)  $S_{100} = \frac{n}{2}(2a + (n-1)d) \text{ or } \frac{n}{2}(a + l)$

M1

$$S_{100} = \frac{100}{2}(6 + 99 \times -5) \text{ or } \frac{100}{2}(3 + -492)$$

dM1

$$= -24\ 450$$


A1 [3]

8 marks

# Circles

## Tips:

- Use completing the square to get to standard form
- Use discriminant to decide how many intersections





14. The circle  $C$  has equation

$$x^2 + y^2 + 16y + k = 0$$

where  $k$  is a constant.

- (a) Find the coordinates of the centre of  $C$ .

(2)

Given that the radius of  $C$  is 10

- (b) find the value of  $k$ .

(2)

The point  $A(a, -16)$ , where  $a > 0$ , lies on the circle  $C$ . The tangent to  $C$  at the point  $A$  crosses the  $x$ -axis at the point  $D$  and crosses the  $y$ -axis at the point  $E$ .



- (c) Find the exact area of triangle  $ODE$ .

(7)

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



14	Mark (a) and (b) together		
(a)	$(0, -8)$	$x = 0 \text{ or } y = -8$ (May be seen on a sketch)	B1
		$x = 0 \text{ and } y = -8$ (May be seen on a sketch)	B1
			(2)
(b)	Uses 64, 100 and $k$ (not $k^2$ ) to obtain a value for $k$		M1
	$k = -36$	cao	A1
	$k = -36$ scores both marks		(2)
14(c)	$y = -16 \Rightarrow a = 6$	Correct $x$ -coordinate. Allow $x = 6$ or just sight of 6. May be seen on a sketch.	B1
	$m_N = \frac{-16+8}{6-0} \left( = -\frac{4}{3} \right)$ or $m_N = \frac{-16+8}{a-0} \left( = -\frac{8}{a} \right)$	Correct attempt at gradient using the centre and their $A$ . Allow one sign slip. If they use $O$ for the centre, this is M0. Allow if in terms of $a$ i.e. if they haven't found or can't find $a$ .	M1
	$m_T = -1 \div -\frac{4}{3} = \dots$ or $m_T = -1 \div -\frac{8}{a} = \dots$	Correct use of perpendicular gradient rule. Allow if in terms of $a$ .	M1
	$y + 16 = \frac{3}{4}(x - "6")$ or $y + 16 = \frac{a}{8}(x - "6")$	Correct straight line method using a gradient which is <b>not</b> the radius gradient and their $A$ or $(a, -16)$ . Allow a gradient in terms of $a$ .	M1
	$x = 0 \Rightarrow y = -\frac{41}{2}$ , $y = 0 \Rightarrow x = \frac{82}{3}$	Correct values	A1
	$\text{Area} = \frac{1}{2} \times \frac{41}{2} \times \frac{82}{3}$	Correct method for area using vertices of the form $(0, 0)$ , $(X, 0)$ and $(0, Y)$ where $X$ and $Y$ are numeric and have come from the intersections of their tangent with the axes. Allow negative lengths here. <b>Dependent on the previous M mark.</b>	dM1
	$= \frac{1681}{6}$ or $280\frac{1}{6}$ or $280.1\dot{6}$ (clear dot over 6)	Cao. Must be <b>positive</b> and may be recovered from sign errors on $-\frac{41}{2}$ and/or $\frac{82}{3}$ <b>but must be from a correct tangent equation.</b>	A1



2 A circle with centre  $C$  has equation  $x^2 + y^2 - 6x + 4y + 4 = 0$ .

(a) Find

(i) the coordinates of  $C$ , [2]

(ii) the radius of the circle. [1]

(b) Determine the set of values of  $k$  for which the line  $y = kx - 3$  does not intersect or touch the circle. [5]

DO NOT WRITE IN THIS AREA

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



2	(a)	(i)	$(x-3)^2 - 9 + (y+2)^2 - 4 + 4 = 0 \Rightarrow (x-3)^2 + (y+2)^2 = 9$  $C(3, -2)$	M1  A1 [2]	1.1  1.1	$(x \pm 3)^2$ and $(y \pm 2)^2$ seen (or implied by correct answer) or one correct coordinate  Accept $x = 3$ and $y = -2$
2	(a)	(ii)	$r = 3$	B1  [1]	1.1	Allow if stated explicitly in (a)(i) but not written down in (a)(ii) www for $r$
2	(b)		$(x-3)^2 + (kx-3+2)^2 = 9$ or $x^2 + (kx-3)^2 - 6x + 4(kx-3) + 4 = 0$ $(1+k^2)x^2 + (-6-2k)x + 1 = 0$ $(-6-2k)^2 - 4(1+k^2)(1)$ $36+24k+4k^2 - 4 - 4k^2 < 0 \Rightarrow 32+24k < 0$ $k < -\frac{4}{3}$	M1*  A1  M1dep*  M1dep*  A1 [5]	3.1a  1.1  3.1a  2.1  2.2a	Substitutes the correct equation of the line into any form of their equation of the circle  oe (all terms on the same side – may not be factorised but should be simplified to 5 terms)  Correct explicit use of discriminant on their 3TQ to get an expression in $k$ only  Discriminant $< 0$ and simplify to the form $ak+b < 0$ (oe)  Fully correct (no additional values)



# Vectors


- I'm going to concentrate on **geometric** problems here – *but of course, you should learn the basics for magnitude, angles with axes, etc.*
- They've not asked one of these style of questions for a long time, so I think they're worth revisiting

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



- 9 Points  $A$ ,  $B$  and  $C$  have position vectors  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{c}$  relative to an origin  $O$  in 3-dimensional space. Rectangles  $OADC$  and  $BEFG$  are the base and top surface of a cuboid.



- The point  $M$  is the midpoint of  $BC$ .
- The point  $X$  lies on  $AM$  such that  $AX = 2XM$ .

- (a) Find  $\overrightarrow{OX}$  in terms of  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{c}$ , simplifying your answer. [4]
- (b) Hence show that the lines  $OF$  and  $AM$  intersect. [2]

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---




9	(a)	<b>Summary method:</b> $\overrightarrow{OM} = \frac{1}{2}(\mathbf{b} + \mathbf{c}) \text{ or } \mathbf{b} + \frac{1}{2}(-\mathbf{b} + \mathbf{c}) \text{ oe}$ $\overrightarrow{AM}$ or $\overrightarrow{MA}$ attempted in terms of $\mathbf{a}$ , $\mathbf{b}$ and $\mathbf{c}$ $(= \pm(\frac{1}{2}(\mathbf{b} + \mathbf{c}) - \mathbf{a}) \text{ oe})$ $\overrightarrow{OX} = \mathbf{a} + \frac{2}{3}\overrightarrow{AM} \text{ or } \overrightarrow{OM} + \frac{1}{3}\overrightarrow{MA} \text{ oe}$ attempted in terms of $\mathbf{a}$ , $\mathbf{b}$ and $\mathbf{c}$ $\overrightarrow{OX} = \frac{1}{3}(\mathbf{a} + \mathbf{b} + \mathbf{c})$	<b>B1</b> <b>M1</b> <b>M1</b> <b>A1</b>	Can be implied May be included in working, eg $\overrightarrow{AX} = \frac{2}{3}(\frac{1}{2}(\mathbf{b} + \mathbf{c}) - \mathbf{a})$ Not necessarily correct Not necessarily correct or equivalent simplified form
---	-----	--	--	--

9	(b)	$\overrightarrow{OF} = \mathbf{a} + \mathbf{b} + \mathbf{c}$ Hence $X$ lies on $OF$ , so $AM$ and $OF$ intersect	<b>B1*</b> <b>B1<sub>dep</sub></b> <b>[2]</b>	Both statements needed. NB dep on B1
---	-----	---	---	--------------------------------------



5



The diagram shows points  $A$  and  $B$ , which have position vectors  $\mathbf{a}$  and  $\mathbf{b}$  with respect to an origin  $O$ .  $P$  is the point on  $OB$  such that  $OP : PB = 3:1$  and  $Q$  is the midpoint of  $AB$ .

- (a) Find  $\vec{PQ}$  in terms of  $\mathbf{a}$  and  $\mathbf{b}$ . [2]

The line  $OA$  is extended to a point  $R$ , so that  $PQR$  is a straight line.

- (b) Explain why  $\vec{PR} = k(2\mathbf{a} - \mathbf{b})$ , where  $k$  is a constant. [2]
- (c) Hence determine the ratio  $OA : AR$ . [4]

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



<b>5</b>	<b>(a)</b>	$\overline{BQ} = \frac{1}{2}(\mathbf{a} - \mathbf{b})$ $\overline{PQ} = \frac{1}{4}\mathbf{b} + \frac{1}{2}(\mathbf{a} - \mathbf{b}) = \frac{1}{2}\mathbf{a} - \frac{1}{4}\mathbf{b}$	<b>B1</b>  <b>[2]</b>	<b>1.1a</b>  <b>1.1</b>	Correct $\overline{BQ}$ or $\overline{QB}$ Correct $\overline{PQ}$
	<b>(b)</b>	$\overline{PR}$ has the same direction as $\overline{PQ}$ , so vector must be a multiple of $\overline{PQ}$ So $\overline{PR} = \lambda(\frac{1}{2}\mathbf{a} - \frac{1}{4}\mathbf{b}) = \frac{1}{4}\lambda(2\mathbf{a} - \mathbf{b}) = k(2\mathbf{a} - \mathbf{b})$ A.G.	<b>B1</b>  <b>B1</b>  <b>[2]</b>	<b>2.4</b>  <b>2.1</b>	Explain parallel (or collinear) vectors have direction vectors that are multiples of each other Show given answer convincingly
	<b>(c)</b>	$\overline{AR} = -\mathbf{a} + \frac{3}{4}\mathbf{b} + k(2\mathbf{a} - \mathbf{b})$ $\overline{AR}$ multiple of $\mathbf{a}$ only, $\frac{3}{4}\mathbf{b} - k\mathbf{b} = 0$ Obtain $k = \frac{3}{4}$ ratio $OA : AR = 2:1$	<b>B1</b>  <b>M1</b>  <b>A1</b>  <b>A1</b>  <b>[4]</b>	<b>1.1</b>  <b>3.1a</b>  <b>1.1</b>  <b>1.1</b>	Correct expression for $\overline{AR}$ (or $\overline{OR}$ ), in terms of $k$ Use coefficient of $\mathbf{b} = 0$ Obtain correct value for $k$ Correct ratio (allow $1: \frac{1}{2}$ ) oe

# Trigonometry

- In my view, the following are essential to know by heart:

**Simple:**

$$\tan x = \frac{\sin x}{\cos x}$$

$$\sec x = \frac{1}{\cos x}$$

$$\operatorname{cosec} x = \frac{1}{\sin x}$$

**Pythagorean:**

$$\sin^2 x + \cos^2 x = 1$$

$$1 + \tan^2 x = \sec^2 x$$

$$1 + \cot^2 x = \operatorname{cosec}^2 x$$

**Double angle:**

$$\sin 2x = 2 \sin x \cos x$$

$$\cos 2x = \cos^2 x - \sin^2 x$$

$$\cos 2x = 2 \cos^2 x - 1$$

$$\cos 2x = 1 - 2 \sin^2 x$$

- Use the shortcuts for finding the solutions:

$$\sin x = \sin(180 - x) \text{ then } \pm 360$$

$$\cos x = \cos(360 - x) \text{ then } \pm 360$$

$$\tan x = \tan(x + 180)$$

- Work on the ‘messy’ side, or one where can add fractions



9. (a) Prove that

$$\sin 2x - \tan x \equiv \tan x \cos 2x, \quad x \neq (2n + 1)90^\circ, \quad n \in \mathbb{Z} \quad (4)$$

(b) Given that  $x \neq 90^\circ$  and  $x \neq 270^\circ$ , solve, for  $0^\circ \leq x < 360^\circ$ ,

$$\sin 2x - \tan x = 3 \tan x \sin x$$

Give your answers in degrees to one decimal place where appropriate.

*(Solutions based entirely on graphical or numerical methods are not acceptable.)*

(5)

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



<b>9(a)</b>	$\begin{aligned}\sin 2x - \tan x &= 2\sin x \cos x - \tan x \\ &= \frac{2\sin x \cos^2 x}{\cos x} - \frac{\sin x}{\cos x} \\ &= \frac{\sin x}{\cos x} \times (2\cos^2 x - 1) \\ &= \tan x \cos 2x\end{aligned}$	M1  M1  dM1 A1*  <b>(4)</b>
<b>(b)</b>	$\begin{aligned}\tan x \cos 2x &= 3 \tan x \sin x \Rightarrow \tan x(\cos 2x - 3 \sin x) = 0 \\ \cos 2x - 3 \sin x &= 0 \\ \Rightarrow 1 - 2 \sin^2 x - 3 \sin x &= 0 \\ \Rightarrow 2 \sin^2 x + 3 \sin x - 1 &= 0 \Rightarrow \sin x = \frac{-3 \pm \sqrt{17}}{4} \Rightarrow x = \dots\end{aligned}$ <p style="margin-left: 100px;">Two of    <math>x = 16.3^\circ, 163.7^\circ, 0, 180^\circ</math></p> <p style="margin-left: 100px;">All four of    <math>x = 16.3^\circ, 163.7^\circ, 0, 180^\circ</math></p>	M1  M1  M1  A1  A1  <b>(5)</b> <b>(9 marks)</b>



Hint: (a) is in tan and cot... keep it that way!



8. (a) Prove that

$$2 \cot 2x + \tan x \equiv \cot x \quad x \neq \frac{n\pi}{2}, n \in \mathbb{Z}$$

(4)

- (b) Hence, or otherwise, solve, for  $-\pi \leq x < \pi$ ,

$$6 \cot 2x + 3 \tan x = \operatorname{cosec}^2 x - 2$$

Give your answers to 3 decimal places.

*(Solutions based entirely on graphical or numerical methods are not acceptable.)*

(6)

DO NOT WRITE IN THIS AREA

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



8 (a)	$\begin{aligned}2 \cot 2x + \tan x &\equiv \frac{2}{\tan 2x} + \tan x \\&\equiv \frac{(1 - \tan^2 x)}{\tan x} + \frac{\tan^2 x}{\tan x} \\&\equiv \frac{1}{\tan x} \\&\equiv \cot x\end{aligned}$	B1 M1 M1 A1*
(b)	$\begin{aligned}6 \cot 2x + 3 \tan x = \operatorname{cosec}^2 x - 2 &\Rightarrow 3 \cot x = \operatorname{cosec}^2 x - 2 \\&\Rightarrow 3 \cot x = 1 + \cot^2 x - 2 \\&\Rightarrow 0 = \cot^2 x - 3 \cot x - 1 \\&\Rightarrow \cot x = \frac{3 \pm \sqrt{13}}{2} \\&\Rightarrow \tan x = \frac{2}{3 \pm \sqrt{13}} \Rightarrow x = .. \\&\Rightarrow x = 0.294, -2.848, -1.277, 1.865\end{aligned}$	(4) M1 A1 M1 M1 A2,1,0



Hint: Can  $1 = \sin^2 x + \cos^2 x$  help us at some point?



8. (a) Prove that

$$\sec 2A + \tan 2A \equiv \frac{\cos A + \sin A}{\cos A - \sin A}, \quad A \neq \frac{(2n+1)\pi}{4}, \quad n \in \mathbb{Z}$$

(5)

- (b) Hence solve, for  $0 \leq \theta < 2\pi$ ,

$$\sec 2\theta + \tan 2\theta = \frac{1}{2}$$

Give your answers to 3 decimal places.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



8(a)	$\begin{aligned}\sec 2A + \tan 2A &= \frac{1}{\cos 2A} + \frac{\sin 2A}{\cos 2A} \\&= \frac{1 + \sin 2A}{\cos 2A} \\&= \frac{1 + 2\sin A \cos A}{\cos^2 A - \sin^2 A} \\&= \frac{\cos^2 A + \sin^2 A + 2\sin A \cos A}{\cos^2 A - \sin^2 A} \\&= \frac{(\cos A + \sin A)(\cos A + \sin A)}{(\cos A + \sin A)(\cos A - \sin A)} \\&= \frac{\cos A + \sin A}{\cos A - \sin A}\end{aligned}$	B1 M1 M1 M1 A1* (5)
(b)	<p> <math>\sec 2\theta + \tan 2\theta = \frac{1}{2}</math> p <math>\frac{\cos \theta + \sin \theta}{\cos \theta - \sin \theta} = \frac{1}{2}</math></p> <p> <b>P</b> <math>2\cos \theta + 2\sin \theta = \cos \theta - \sin \theta</math> </p> <p> <b>P</b> <math>\tan \theta = -\frac{1}{3}</math> </p> <p> <b>P</b> <math>\theta = \text{arctan } -\frac{1}{3}</math> or <math>\theta = 2.820, 5.961</math> </p>	M1 A1 dM1A1 (4) (9 marks)




# Implicit Differentiation

- If you differentiate  $y$  with respect to  $x$ , it becomes  $\frac{dy}{dx}$
- If you differentiate a function in  $y$  with respect to  $x$ , do as you expect, then multiply by  $\frac{dy}{dx}$
- I recommend writing out the product rule carefully for any product expressions including a  $y$ 
  - e.g.  $4x^2y^3$
- If you have  $\frac{dy}{dx} = \frac{f(x,y)}{g(x,y)}$ , then  $f(x,y) = 0$  for zero (horizontal) gradient, and  $g(x,y) = 0$  for undefined (vertical) gradient

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**6 In this question you must show detailed reasoning.**

The diagram shows the curve with equation  $4xy = 2(x^2 + 4y^2) - 9x$ .

(a) Show that  $\frac{dy}{dx} = \frac{4x - 4y - 9}{4x - 16y}$ . [3]

At the point  $P$  on the curve the tangent to the curve is parallel to the  $y$ -axis and at the point  $Q$  on the curve the tangent to the curve is parallel to the  $x$ -axis.

(b) Show that the distance  $PQ$  is  $k\sqrt{5}$ , where  $k$  is a rational number to be determined. [8]

I've pre-done some of (b) so as not to waste your time in the session

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



6	(a)	<b>DR</b> $4y + 4x \frac{dy}{dx} = 4x + 16y \frac{dy}{dx} - 9$ $4x \frac{dy}{dx} - 16y \frac{dy}{dx} = 4x - 4y - 9 \Rightarrow \frac{dy}{dx} = \frac{4x - 4y - 9}{4x - 16y}$	<b>M1</b> <b>A1</b>	<b>1.1</b> <b>1.1</b>	$4xy = 2(x^2 + 4y^2) - 9x$ For correct differentiation of either LHS or RHS, even if not in an equation
6	(b)	<b>DR</b> (At P) $4x - 16y = 0$ $x = 4y \Rightarrow 16y^2 = 2(16y^2 + 4y^2) - 36y$ $24y^2 - 36y = 0$ $y(2y - 3) = 0 \Rightarrow y = \frac{3}{2}$ $P\left(6, \frac{3}{2}\right)$ (At Q) $4x - 4y - 9 = 0$ $\Rightarrow 4x\left(x - \frac{9}{4}\right) = 2x^2 + 8\left(x - \frac{9}{4}\right)^2 - 9x$ $4x^2 - 24x + 27 = 0$ $Q\left(\frac{3}{2}, -\frac{3}{4}\right)$ only $PQ^2 = \left(6 - \frac{3}{2}\right)^2 + \left(\frac{3}{2} - \left(-\frac{3}{4}\right)\right)^2$ $PQ = \frac{9}{4}\sqrt{5}$	<b>M1*</b> <b>M1dep*</b> <b>A1</b> <b>M1*</b> <b>M1dep*</b> <b>A1</b> <b>M1</b> <b>A1</b>	<b>3.1a</b> <b>2.1</b> <b>1.1</b> <b>3.1a</b> <b>2.1</b> <b>3.2a</b> <b>1.1</b> <b>2.2a</b> <b>[8]</b>	Forms two-term quadratic equation in y or x (if correct $x^2 - 6x = 0$ ) Forms three-term quadratic equation in y or x (if correct $16y^2 - 24y - 27 = 0$ ) Correct implies distance formula for their P and Q www



6. A curve has equation

$$4y^2 + 3x = 6ye^{-2x}$$

- (a) Find  $\frac{dy}{dx}$  in terms of  $x$  and  $y$ .

(5)

The curve crosses the  $y$ -axis at the origin and at the point  $P$ .

- (b) Find the equation of the normal to the curve at  $P$ , writing your answer in the form  $y = mx + c$  where  $m$  and  $c$  are constants to be found.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



6(a)	$4y^2 + 3x = 6ye^{-2x}$	
	$4y^2 + 3x \rightarrow 8y \frac{dy}{dx} + 3$	B1
	$6ye^{-2x} \rightarrow -12ye^{-2x} + 6e^{-2x} \frac{dy}{dx}$	M1 A1
	$8y \frac{dy}{dx} + 3 = -12ye^{-2x} + 6e^{-2x} \frac{dy}{dx} \Rightarrow \frac{dy}{dx} = \frac{12ye^{-2x} + 3}{6e^{-2x} - 8y}$ oe	M1 A1
		(5)
(b)	Sets $x = 0$ in $4y^2 + 3x = 6ye^{-2x} \Rightarrow y = \frac{3}{2}$ oe	B1
	Substitutes $\left(0, \frac{3}{2}\right)$ in their $\frac{dy}{dx} = \frac{12ye^{-2x} + 3}{6e^{-2x} - 8y} = \left(\frac{7}{-2}\right)$	M1
	$m_N = -1 \div \frac{7}{-2} \Rightarrow y = \frac{2}{7}x + \frac{3}{2}$	dM1
	$y = \frac{2}{7}x + \frac{3}{2}$ oe e.g. $y = \frac{6}{21}x + \frac{3}{2}$	A1
		(4)



(ii) Given  $x = \sin^2 2y$ ,  $0 < y < \frac{\pi}{4}$ , find  $\frac{dy}{dx}$  as a function of  $y$ .

Write your answer in the form

$$\frac{dy}{dx} = p \operatorname{cosec}(qy), \quad 0 < y < \frac{\pi}{4}$$

where  $p$  and  $q$  are constants to be determined.

(5)

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



(5)

(ii)

$$x = \sin^2 2y \Rightarrow \frac{dx}{dy} = 2 \sin 2y \times 2 \cos 2y$$

M1A1

Uses  $\sin 4y = 2 \sin 2y \cos 2y$  in their expression

M1

$$\frac{dx}{dy} = 2 \sin 4y \Rightarrow \frac{dy}{dx} = \frac{1}{2 \sin 4y} = \frac{1}{2} \operatorname{cosec} 4y$$

M1A1

(5)  
**(10 marks)**



5. The point  $P$  lies on the curve with equation

$$x = (4y - \sin 2y)^2$$

Given that  $P$  has  $(x, y)$  coordinates  $\left( p, \frac{\pi}{2} \right)$ , where  $p$  is a constant,

- (a) find the exact value of  $p$ . (1)

The tangent to the curve at  $P$  cuts the  $y$ -axis at the point  $A$ .

- (b) Use calculus to find the coordinates of  $A$ . (6)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



<b>5.(a)</b> $p = 4\pi^2$ or $(2\pi)^2$	B1   <b>(1)</b>
<b>(b)</b> $x = (4y - \sin 2y)^2 \Rightarrow \frac{dx}{dy} = 2(4y - \sin 2y)(4 - 2\cos 2y)$ Sub $y = \frac{\pi}{2}$ into $\frac{dx}{dy} = 2(4y - \sin 2y)(4 - 2\cos 2y)$ $p \quad \frac{dx}{dy} = 24\pi \quad (= 75.4) \quad / \quad \frac{dy}{dx} = \frac{1}{24\pi} \quad (= 0.013)$ Equation of tangent $y - \frac{\pi}{2} = \frac{1}{24\pi}(x - 4\pi^2)$ Using $y - \frac{\pi}{2} = \frac{1}{24\pi}(x - 4\pi^2)$ with $x = 0$ p $y = \frac{\pi}{3}$ cso	M1A1        M1        M1        M1, A1        <b>(6)</b>
	<b>(7 marks)</b>



8. Given that

$$y = 8 \tan(2x), \quad -\frac{\pi}{4} < x < \frac{\pi}{4}$$

show that

$$\frac{dx}{dy} = \frac{A}{B + y^2}$$

where  $A$  and  $B$  are integers to be found.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



8

 Differentiates wrt  $x$ 

$$\text{Inverts to get } \frac{dx}{dy} = \frac{1}{16 \sec^2 2x}$$

$$= \frac{1}{16(1 + \tan^2 2x)}$$

$$\frac{dy}{dx} = 16 \sec^2(2x) \text{ oe}$$

$$\begin{aligned}\frac{dy}{dx} &= 16(1 + \tan^2(2x)) \\ &= 16 \left( 1 + \left( \frac{y}{8} \right)^2 \right)\end{aligned}$$

$$\begin{aligned}\frac{dx}{dy} &= \frac{A}{B + y^2} \\ &= \frac{4}{64 + y^2}\end{aligned}$$

M1

dM1

ddM1

A1

(4)

(4 marks)



# Modelling with differentiation

- Connected rates of change
  - Write out what you are looking for, then split the derivative up:
    - $\frac{dh}{dt} = \frac{dh}{dx} \times \frac{dx}{dt}$
- Optimisation problems
  - Setup by eliminating variables, differentiate and set to 0, then solve

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



4.




Diagram not  
drawn to scale

**Figure 1**

A water container is made in the shape of a hollow inverted right circular cone with semi-vertical angle of  $30^\circ$ , as shown in Figure 1. The height of the container is 50 cm.

When the depth of the water in the container is  $h$  cm, the surface of the water has radius  $r$  cm and the volume of water is  $V$  cm<sup>3</sup>.

(a) Show that  $V = \frac{1}{9}\pi h^3$

[You may assume the formula  $V = \frac{1}{3}\pi r^2 h$  for the volume of a cone.]

(2)

Given that the volume of water in the container increases at a constant rate of  $200 \text{ cm}^3 \text{ s}^{-1}$ ,

- (b) find the rate of change of the depth of the water, in  $\text{cm s}^{-1}$ , when  $h = 15$   
Give your answer in its simplest form in terms of  $\pi$ .

(4)

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---


---



4. (a)	$\frac{r}{h} = \tan 30 \Rightarrow r = h \tan 30 \left\{ \Rightarrow r = \frac{h}{\sqrt{3}} \text{ or } r = \frac{\sqrt{3}}{3} h \right\}$ <b>or</b> $\frac{h}{r} = \tan 60 \Rightarrow r = \frac{h}{\tan 60} \left\{ \Rightarrow r = \frac{h}{\sqrt{3}} \text{ or } r = \frac{\sqrt{3}}{3} h \right\}$ <b>or</b> $\frac{r}{\sin 30} = \frac{h}{\sin 60} \Rightarrow r = \frac{h \sin 30}{\sin 60} \left\{ \Rightarrow r = \frac{h}{\sqrt{3}} \text{ or } r = \frac{\sqrt{3}}{3} h \right\}$ <b>or</b> $h^2 + r^2 = (2r)^2 \Rightarrow r^2 = \frac{1}{3}h^2$	Correct use of trigonometry to find $r$ in terms of $h$ <b>or</b> correct use of Pythagoras to find $r^2$ in terms of $h^2$	M1
	$\left\{ V = \frac{1}{3}\pi r^2 h \Rightarrow \right\} V = \frac{1}{3}\pi \left( \frac{h}{\sqrt{3}} \right)^2 h \Rightarrow V = \frac{1}{9}\pi h^3 *$	Correct proof of $V = \frac{1}{9}\pi h^3$ or $V = \frac{1}{9}h^3\pi$ Or shows $\frac{1}{9}\pi h^3$ or $\frac{1}{9}h^3\pi$ with some reference to $V =$ in their solution	A1 *
		[2]	
(b) Way 1	$\frac{dV}{dt} = 200$		
	$\frac{dV}{dh} = \frac{1}{3}\pi h^2$	$\frac{1}{3}\pi h^2$ o.e.	B1
	Either <ul style="list-style-type: none"> <li>• <math>\left\{ \frac{dV}{dh} \times \frac{dh}{dt} = \frac{dV}{dt} \Rightarrow \right\} \left( \frac{1}{3}\pi h^2 \right) \frac{dh}{dt} = 200</math></li> <li>• <math>\left\{ \frac{dh}{dt} = \frac{dV}{dt} \div \frac{dV}{dh} \Rightarrow \right\} \frac{dh}{dt} = 200 \div \left( \frac{1}{3}\pi h^2 \right)</math></li> </ul>	either $\left( \text{their } \frac{dV}{dh} \right) \times \frac{dh}{dt} = 200$ or $200 \div \left( \text{their } \frac{dV}{dh} \right)$	M1
	When $h = 15, \frac{dh}{dt} = 200 \div \frac{1}{3}\pi(15)^2 \quad \left\{ = \frac{200}{75\pi} = \frac{600}{225\pi} \right\}$	<b>dependent on the previous M mark</b>	dM1
	$\frac{dh}{dt} = \frac{8}{3\pi} (\text{cms}^{-1})$	$\frac{8}{3\pi}$ A1 cao	[4] 6



5



For a cone with base radius  $r$ , height  $h$  and slant height  $l$ , the following formulae are given.

$$\text{Curved surface area, } S = \pi r l$$

$$\text{Volume, } V = \frac{1}{3}\pi r^2 h$$

A container is to be designed in the shape of an inverted cone with no lid. The base radius is  $rm$  and the volume is  $V\text{m}^3$ .

The area of the material to be used for the cone is  $4\pi\text{ m}^2$ .

(a) Show that  $V = \frac{1}{3}\pi \sqrt{16r^2 - r^6}$ . [4]

(b) In this question you must show detailed reasoning.

It is given that  $V$  has a maximum value for a certain value of  $r$ .

Find the maximum value of  $V$ , giving your answer correct to 3 significant figures. [5]

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



5	(a)		An example of a correct method: $l = \frac{4}{r} \text{ or } l = \frac{4\pi}{r\pi} \text{ exactly (not } lr = 4\text{)}$ $(h = \sqrt{l^2 - r^2})$ $h = \sqrt{\frac{16}{r^2} - r^2} \text{ or } \frac{\sqrt{16-r^4}}{r} \text{ oe}$ $V = \frac{1}{3}\pi r^2 \sqrt{\frac{16}{r^2} - r^2} \text{ or } \frac{1}{3}\pi r^2 \frac{\sqrt{16-r^4}}{r} \text{ oe}$ $(\text{=} \frac{\pi}{3} \sqrt{16r^2 - r^6} \text{ AG})$	<b>B1</b>	<b>3.1a</b>	Other correct methods may be seen eg $lr = 4$ : B1, find $h$ into $l$ & $r$ : B1, Subst $h$ & $lr$ into $V$ : M1, convincing: A1
					<b>1.1</b>	Express $l$ correctly in terms of $r$ May be implied
5	(b)	<b>DR</b>	$\frac{d}{dr} \left( \frac{\pi}{3} \sqrt{16r^2 - r^6} \right)$ $\frac{\pi(32r - 6r^5)}{3 \times 2\sqrt{16r^2 - r^6}} = 0 \text{ oe}$ (Their derivative = 0) $r = \frac{2}{\sqrt[4]{3}} \text{ or } \sqrt[4]{\frac{16}{3}} \text{ oe or } 1.52 \text{ (3 sf)} \text{ Allow 1.5}$ $\text{or } r^2 = \frac{4}{\sqrt{3}}$ $r = -\frac{2}{\sqrt[4]{3}} \text{ or } -1.52 \text{ invalid OR } r = 0 \text{ invalid or } r > 0$ $(V_{\max} = \frac{\pi}{3} \sqrt{16 \times 1.51967^2 - 1.51967^6})$ $\text{Max } V = 5.20 \text{ (3 sf)} \text{ Allow 5.2 or a.r.t. 5.2}$	<b>M1</b>	<b>1.1a</b>	Attempt differentiate $V$ or $\frac{V}{\pi}$ or $3V$
					<b>A1</b>	<b>2.1</b>
			$r = \frac{2}{\sqrt[4]{3}}$ $\text{or } r^2 = \frac{4}{\sqrt{3}}$ $r = -\frac{2}{\sqrt[4]{3}} \text{ or } -1.52 \text{ invalid OR } r = 0 \text{ invalid or } r > 0$ $(V_{\max} = \frac{\pi}{3} \sqrt{16 \times 1.51967^2 - 1.51967^6})$ $\text{Max } V = 5.20 \text{ (3 sf)} \text{ Allow 5.2 or a.r.t. 5.2}$	<b>A1</b>	<b>1.1</b>	Lose this mark if incorrect values of $r$ also given, eg $r = \pm 2$ obtained from $(16r^2 - r^6)^{-\frac{1}{2}} = 0$
					<b>B1f</b>	<b>3.2a</b>
				<b>A1</b>	<b>1.1</b>	Condone $V = 5.20 \text{ m}^3$
					<b>[5]</b>	



# Integration


- By parts
- Partial fractions
- ... more in the parametric section...

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



8



The diagram shows the curve  $M$  with equation  $y = xe^{-2x}$ .

- (a) Show that  $M$  has a point of inflection at the point  $P$  where  $x = 1$ . [5]

The line  $L$  passes through the origin  $O$  and the point  $P$ . The shaded region  $R$  is enclosed by the curve  $M$  and the line  $L$ .

- (b) Show that the area of  $R$  is given by

$$\frac{1}{4}(a + be^{-2}),$$

where  $a$  and  $b$  are integers to be determined.

[6]

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



8	(a)	$y' = e^{-2x}(1 - 2x)$ $y'' = e^{-2x}(-4 + 4x)$  $y'' = 0$ at $x = 1$ and $y''(0.5) = -2e^{-1} < 0$ , $y''(1.5) = 2e^{-3} > 0$ (so change of sign indicates a point of inflection at $x = 1$ )	<b>M1*</b> <b>A1</b> <b>A1ft</b> <b>M1dep*</b> <b>A1 [5]</b>	<b>2.1</b> <b>1.1</b> <b>1.1</b> <b>3.1a</b> <b>2.2a</b>	Differentiates $y$ with respect to $x$ – answer of the form $\pm e^{-2x} \pm \lambda x e^{-2x}$ Follow through their first derivative Solves $y'' = 0$ (or attempts to verify $y'' = 0$ by substituting $x = 1$ ) or considers sign change either side of $y'' = 0$ Conclusion not required for this mark
8	(b)	$\int xe^{-2x} dx = -\frac{1}{2}xe^{-2x} + \frac{1}{2}\int e^{-2x} dx$  $\int xe^{-2x} dx = -\frac{1}{2}xe^{-2x} - \frac{1}{4}e^{-2x}$  $\begin{aligned} \int_0^1 xe^{-2x} dx &= \left[ -\frac{1}{2}xe^{-2x} - \frac{1}{4}e^{-2x} \right]_0^1 \\ &= \left( -\frac{1}{2}e^{-2} - \frac{1}{4}e^{-2} \right) - \left( 0 - \frac{1}{4} \right) \end{aligned}$ $\frac{1}{4} - \frac{3}{4}e^{-2}$ Area of triangle below $OP = \frac{1}{2}e^{-2}$ $= \frac{1}{4}(1 - 5e^{-2})$	<b>M1*</b> <b>A1</b> <b>M1dep*</b> <b>A1</b> <b>B1</b> <b>A1</b>	<b>2.1</b> <b>1.1</b> <b>1.1</b> <b>1.1</b> <b>1.1</b> <b>2.2a</b>	Integration by parts – of the form $\pm \alpha xe^{-2x} \pm \beta \int e^{-2x} dx$ Use of correct limits in their fully integrated expression – need not be simplified (or equivalent) Allow unsimplified Or by correctly evaluating $\int_0^1 e^{-2x} dx$ $a = 1, b = -5$ (must be in this form)



This one is a little weird – so I hope by showing you, you'll be ready for it in the exam!  
It's where you have an exponential and trig product integral.




Figure 2

(a) Find  $\int e^{2x} \sin x \, dx$  (5)

Figure 2 shows a sketch of part of the curve with equation

$$y = e^{2x} \sin x \quad x \geq 0$$

The finite region  $R$  is bounded by the curve and the  $x$ -axis and is shown shaded in Figure 2.

(b) Show that the exact area of  $R$  is  $\frac{e^{2\pi} + 1}{5}$

*(Solutions relying on calculator technology are not acceptable.)*

(2)

DO NOT WRITE IN THIS AREA

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Question Number	Scheme	Marks
7(a) Way 1	$\int e^{2x} \sin x \, dx = \frac{1}{2} e^{2x} \sin x - \int \frac{1}{2} e^{2x} \cos x \, dx$	M1
	$= \dots - \frac{1}{4} e^{2x} \cos x - \int \frac{1}{4} e^{2x} \sin x \, dx$	dM1
	$\int e^{2x} \sin x \, dx = \frac{1}{2} e^{2x} \sin x - \frac{1}{4} e^{2x} \cos x - \int \frac{1}{4} e^{2x} \sin x \, dx$	A1
	$\frac{5}{4} \int e^{2x} \sin x \, dx = \frac{1}{2} e^{2x} \sin x - \frac{1}{4} e^{2x} \cos x \Rightarrow \int e^{2x} \sin x \, dx = \dots$	ddM1
	$= \frac{2}{5} e^{2x} \sin x - \frac{1}{5} e^{2x} \cos x + c$	A1
		(5)
7(a) Way 2	$\int e^{2x} \sin x \, dx = -e^{2x} \cos x + \int 2e^{2x} \cos x \, dx$	M1
	$= \dots + 2e^{2x} \sin x - \int 4e^{2x} \sin x \, dx$	dM1
	$\int e^{2x} \sin x \, dx = -e^{2x} \cos x + 2e^{2x} \sin x - \int 4e^{2x} \sin x \, dx$	A1
	$5 \int e^{2x} \sin x \, dx = -e^{2x} \cos x + 2e^{2x} \sin x \Rightarrow \int e^{2x} \sin x \, dx = \dots$	ddM1
	$= \frac{2}{5} e^{2x} \sin x - \frac{1}{5} e^{2x} \cos x + c$	A1
		(5)
(b)	$\left( \frac{2}{5} e^{2\pi} \sin \pi - \frac{1}{5} e^{2\pi} \cos \pi \right) - \left( \frac{2}{5} e^0 \sin 0 - \frac{1}{5} e^0 \cos 0 \right) = \dots$	M1
	$= \frac{1}{5} e^{2\pi} + \frac{1}{5} = \frac{e^{2\pi} + 1}{5} *$	A1*
		(2)
		(7 marks)



(3)

5. (a) Express  $\frac{9(4+x)}{16-9x^2}$  in partial fractions.

Given that

$$f(x) = \frac{9(4+x)}{16-9x^2}, \quad x \in \mathbb{R}, \quad -\frac{4}{3} < x < \frac{4}{3}$$

- (b) express  $\int f(x) dx$  in the form  $\ln(g(x))$ , where  $g(x)$  is a rational function.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



**5(a)**

$$\frac{9(4+x)}{16-9x^2} \equiv \frac{A}{(4-3x)} + \frac{B}{(4+3x)} \Rightarrow A \text{ or } B$$

$A = 6$  or  $B = 3$  obtained at any point of the solution

$$\frac{9(4+x)}{16-9x^2} \equiv \frac{6}{(4-3x)} + \frac{3}{(4+3x)}$$

M1

A1

A1

(3)

**(b)**

$$\begin{aligned} \int \frac{9(4+x)}{16-9x^2} dx &\equiv \int \frac{A}{(4-3x)} + \frac{B}{(4+3x)} dx \\ &= -\frac{A}{3} \ln(4-3x) + \frac{B}{3} \ln(4+3x) (+c) \end{aligned}$$

$$(-2 \ln(4-3x) + \ln(4+3x) (+c))$$

$$= \ln \frac{(4+3x)}{(4-3x)^2} + c, = \ln \frac{k(4+3x)}{(4-3x)^2} \quad \text{or} \quad \ln \left| \frac{k(4+3x)}{(4-3x)^2} \right|$$

M1 A1ft

M1, A1

(7 marks)

(4)



# Parametrics

- ‘Parametric world’ is easier to work in than ‘Cartesian world’ – i.e. only create a Cartesian equation if they ask for one
- Differentiation:  $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$
- Integration:  $\int y \frac{dx}{dt} dt$  *Make sure the limits are in t*
- Be prepared to use a lot of different techniques in these questions – differentiation, integration, exponentials, logarithms, trigonometric identities...

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



5.




Figure 2

Figure 2 shows a sketch of the curve  $C$  with parametric equations

$$x = 4 \tan t, \quad y = 5\sqrt{3} \sin 2t, \quad 0 \leq t < \frac{\pi}{2}$$

The point  $P$  lies on  $C$  and has coordinates  $\left(4\sqrt{3}, \frac{15}{2}\right)$ .

- (a) Find the exact value of  $\frac{dy}{dx}$  at the point  $P$ .

Give your answer as a simplified surd.

(4)

The point  $Q$  lies on the curve  $C$ , where  $\frac{dy}{dx} = 0$

- (b) Find the exact coordinates of the point  $Q$ .

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



5.	$x = 4 \tan t, \quad y = 5\sqrt{3} \sin 2t, \quad 0 \leq t < \frac{\pi}{2}$		
(a) Way 1	$\frac{dx}{dt} = 4 \sec^2 t, \quad \frac{dy}{dt} = 10\sqrt{3} \cos 2t$ $\Rightarrow \frac{dy}{dx} = \frac{10\sqrt{3} \cos 2t}{4 \sec^2 t} \quad \left\{ = \frac{5}{2}\sqrt{3} \cos 2t \cos^2 t \right\}$ $\left\{ \text{At } P\left(4\sqrt{3}, \frac{15}{2}\right), \quad t = \frac{\pi}{3} \right\}$	<b>Either both</b> $x$ <b>and</b> $y$ <b>are differentiated correctly with respect to</b> $t$ <b>or</b> their $\frac{dy}{dt}$ <b>divided by</b> their $\frac{dx}{dt}$ <b>or</b> applies $\frac{dy}{dt}$ <b>multiplied by</b> their $\frac{dt}{dx}$ Correct $\frac{dy}{dx}$ (Can be implied)	M1 A1 oe
	$\frac{dy}{dx} = \frac{10\sqrt{3} \cos\left(\frac{2\pi}{3}\right)}{4 \sec^2\left(\frac{\pi}{3}\right)}$	<b>dependent on the previous M mark</b> <i>Some evidence</i> of substituting $t = \frac{\pi}{3}$ or $t = 60^\circ$ into their $\frac{dy}{dx}$	dM1
	$\frac{dy}{dx} = -\frac{5}{16}\sqrt{3}$ or $-\frac{15}{16\sqrt{3}}$	$-\frac{5}{16}\sqrt{3}$ or $-\frac{15}{16\sqrt{3}}$ <b>from a correct solution only</b>	A1 cso
(b)	$\left\{ 10\sqrt{3} \cos 2t = 0 \Rightarrow t = \frac{\pi}{4} \right\}$		[4]
	$\text{So } x = 4 \tan\left(\frac{\pi}{4}\right), \quad y = 5\sqrt{3} \sin\left(2\left(\frac{\pi}{4}\right)\right)$	At least one of either $x = 4 \tan\left(\frac{\pi}{4}\right)$ or $y = 5\sqrt{3} \sin\left(2\left(\frac{\pi}{4}\right)\right)$ or $x = 4$ or $y = 5\sqrt{3}$ or $y = \text{awrt } 8.7$	M1
	Coordinates are $(4, 5\sqrt{3})$	$(4, 5\sqrt{3})$ or $x = 4, y = 5\sqrt{3}$	A1
			[2]
			6



The curve  $C$  shown in Figure 3 has parametric equations

$$x = 3 \cos t, \quad y = 9 \sin 2t, \quad 0 \leq t \leq 2\pi$$

The curve  $C$  meets the  $x$ -axis at the origin and at the points  $A$  and  $B$ , as shown in Figure 3.

- (a) Write down the coordinates of  $A$  and  $B$ .

(2)

- (b) Find the values of  $t$  at which the curve passes through the origin.

(2)

- (c) Find an expression for  $\frac{dy}{dx}$  in terms of  $t$ , and hence find the gradient of the curve

when  $t = \frac{\pi}{6}$

(4)

- (d) Show that the cartesian equation for the curve  $C$  can be written in the form

$$y^2 = ax^2(b - x^2)$$

where  $a$  and  $b$  are integers to be determined.

(4)




Figure 3



**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---






11 (a)	(3,0) and (-3, 0)	B1, B1  M1 A1	(2)  (2)
(b)	$\frac{\pi}{2}$ and $\frac{3\pi}{2}$		
(c)	$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{\frac{18 \cos 2t}{-3 \sin t}}{\frac{18 \times \frac{1}{2}}{-3 \times \frac{1}{2}}} = -6$	M1 A1  dM1 A1	(4)
(d)	$y^2 = 81 \times 4 \sin^2 t \cos^2 t$  Attempts to replace $\cos^2 t = \frac{x^2}{9}$ and $\sin^2 t = 1 - \frac{x^2}{9}$  Correct eqn $y^2 = 81 \times 4 \times \left(1 - \frac{x^2}{9}\right) \times \frac{x^2}{9}$  Obtain $y^2 = 4x^2(9 - x^2)$	$y = 9 \times 2 \sin t \cos t$  Attempts to replace $\cos t = \frac{x}{3}$ and $\sin t = \sqrt{1 - \frac{x^2}{9}}$  Correct eqn $y = 9 \times 2 \times \sqrt{1 - \frac{x^2}{9}} \times \frac{x}{3}$  Obtain $y^2 = 4x^2(9 - x^2)$	M1  M1  A1  A1  (4)  <b>(12 marks)</b>



5



The diagram shows the curve  $C$  with parametric equations

$$x = \frac{3}{t}, \quad y = t^3 e^{-2t}, \quad \text{where } t > 0.$$

The maximum point on  $C$  is denoted by  $P$ .

- (a) Determine the exact coordinates of  $P$ .

[4]

The shaded region  $R$  is enclosed by the curve, the  $x$ -axis and the lines  $x = 1$  and  $x = 6$ .

- (b) Show that the area of  $R$  is given by

$$\int_a^b 3te^{-2t} dt,$$

where  $a$  and  $b$  are constants to be determined.

[3]

- (c) Hence determine the exact area of  $R$ .

[5]

DO NOT WRITE IN THIS AREA

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA



5	(a)	$\frac{dy}{dt} = 3t^2 e^{-2t} + t^3 (-2e^{-2t})$ $\frac{dy}{dt} = 0 \Rightarrow t^2 e^{-2t} (3 - 2t) = 0 \Rightarrow t = \dots$ $t = \frac{3}{2}$ $P\left(2, \frac{27}{8} e^{-3}\right)$	<b>M1*</b>  <b>M1dep*</b>  <b>A1</b> <b>A1</b>	<b>2.1</b>  <b>1.1</b>  <b>1.1</b> <b>2.2a</b> <b>[4]</b>	<p>Attempts to differentiate <math>y</math> with respect to <math>t</math> using the product rule – answer of the form <math>\frac{dy}{dt} = \lambda t^2 e^{-2t} + \mu t^3 e^{-2t}</math> or <math>y' = \alpha x^{-5} e^{-6x^{-1}} (\beta x + \gamma)</math></p> <p>Sets their derivative equal to zero and solves for <math>t</math></p> <p>From correct working only (or for <math>x = 2</math>)</p> <p>From correct working only y-coordinate must be exact but ISW</p>
5	(b)	$\frac{dx}{dt} = -3t^{-2}$ and $\int y \frac{dx}{dt} dt$  $x = 6 \Rightarrow t = 0.5$ and $x = 1 \Rightarrow t = 3$  $\text{Area} = \int_3^{0.5} t^3 e^{-2t} \left(-\frac{3}{t^2}\right) dt = \int_3^{0.5} -3te^{-2t} dt = \int_{0.5}^3 3te^{-2t} dt$	<b>M1</b>  <b>B1</b>  <b>A1</b>	<b>2.1</b>  <b>1.1</b>  <b>2.2a</b> <b>[3]</b>	<p>Differentiates <math>x</math> with respect to <math>t</math> and attempts to set up integral for the required area</p> <p>Stating 0.5 and 3 is sufficient for this mark</p> <p>Must be correctly shown</p>
5	(c)	$u = 3t$ , and $dv$ or $\frac{dy}{dt} = e^{-2t}$  $\int 3te^{-2t} dt = -\frac{3}{2}te^{-2t} + \frac{3}{2} \int e^{-2t} dt$ $= \dots -\frac{3}{4}e^{-2t} (+c)$ $\left[ -\frac{3}{2}te^{-2t} - \frac{3}{4}e^{-2t} \right]_{0.5}^3 = \left( -\frac{3}{2}(3)e^{-6} - \frac{3}{4}e^{-6} \right) - \left( -\frac{3}{2}(0.5)e^{-1} - \frac{3}{4}e^{-1} \right)$ $\text{Area} = -\frac{21}{4}e^{-6} + \frac{3}{2}e^{-1}$	<b>M1*</b>  <b>A1</b>  <b>A1</b>  <b>M1dep*</b>  <b>A1</b>	<b>1.1</b>  <b>1.1</b>  <b>1.1</b>  <b>1.1</b> <b>2.2a</b> <b>[5]</b>	<p>Integrating by parts as far as <math>f(t) \pm \int g(t) dt</math></p> <p>Allow correct un-simplified for both A marks</p> <p>Use of their <math>t</math>-limits (so not 1 and 6) in fully integrated expression (must subtract bottom limit from top limit)</p> <p>ISW once correct exact answer seen</p>



Figure 4 shows a sketch of part of the curve  $C$  with parametric equations

$$x = 3\theta \sin \theta, \quad y = \sec^3 \theta, \quad 0 \leq \theta < \frac{\pi}{2}$$

The point  $P(k, 8)$  lies on  $C$ , where  $k$  is a constant.

- (a) Find the exact value of  $k$ .

(2)

The finite region  $R$ , shown shaded in Figure 4, is bounded by the curve  $C$ , the  $y$ -axis, the  $x$ -axis and the line with equation  $x = k$ .

- (b) Show that the area of  $R$  can be expressed in the form

$$\lambda \int_a^\beta (\theta \sec^2 \theta + \tan \theta \sec^2 \theta) d\theta$$

where  $\lambda$ ,  $\alpha$  and  $\beta$  are constants to be determined.

(4)

- (c) Hence use integration to find the exact value of the area of  $R$ .

(6)




Figure 4

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



8.	$x = 3\theta \sin \theta, \quad y = \sec^3 \theta, \quad 0 \leq \theta < \frac{\pi}{2}$		
(a)	$\{ \text{When } y = 8, \} \quad 8 = \sec^3 \theta \Rightarrow \cos^3 \theta = \frac{1}{8} \Rightarrow \cos \theta = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{3}$ $k \text{ (or } x) = 3\left(\frac{\pi}{3}\right) \sin\left(\frac{\pi}{3}\right)$ $\text{so } k \text{ (or } x) = \frac{\sqrt{3}\pi}{2}$	Sets $y = 8$ to find $\theta$ and attempts to substitute their $\theta$ into $x = 3\theta \sin \theta$	M1
		$\frac{\sqrt{3}\pi}{2} \text{ or } \frac{3\pi}{2\sqrt{3}}$	A1
	<b>Note:</b> Obtaining two value for $k$ without accepting the correct value is final A0		
(b)	$\frac{dx}{d\theta} = 3\sin \theta + 3\theta \cos \theta$ $\left\{ \int y \frac{dx}{d\theta} \{ d\theta \} \right\} = \int (\sec^3 \theta)(3\sin \theta + 3\theta \cos \theta) \{ d\theta \}$ $= 3 \int \sec^2 \theta + \tan \theta \sec^2 \theta \, d\theta$ $x = 0 \text{ and } x = k \Rightarrow \underline{\alpha = 0} \text{ and } \underline{\beta = \frac{\pi}{3}}$	$3\theta \sin \theta \rightarrow 3\sin \theta + 3\theta \cos \theta$ Can be implied by later working Applies $(\pm K \sec \theta) \left( \text{their } \frac{dx}{d\theta} \right)$ Ignore integral sign and $d\theta$ ; $K \neq 0$ Achieves the correct result no errors in their working, e.g. bracketing or manipulation errors. <b>Must have</b> integral sign and $d\theta$ in their final answer.	B1 M1 A1 * B1
	<b>Note:</b> The work for the final B1 mark must be seen in part (b) only.		
(c) Way 1	$\left\{ \int \sec^2 \theta \, d\theta \right\} = \theta \tan \theta - \int \tan \theta \{ d\theta \}$ $= \theta \tan \theta - \ln(\sec \theta)$ or $= \theta \tan \theta + \ln(\cos \theta)$	$\theta \sec^2 \theta \rightarrow A\theta g(\theta) - B \int g(\theta), A > 0, B > 0,$ where $g(\theta)$ is a trigonometric function in $\theta$ and $g(\theta) = \text{their } \int \sec^2 \theta \, d\theta$ . [Note: $g(\theta) \neq \sec^2 \theta$ ] <b>dependent on the previous M mark</b> Either $\lambda \theta \sec^2 \theta \rightarrow A\theta \tan \theta - B \int \tan \theta, A > 0, B > 0$ or $\theta \sec^2 \theta \rightarrow \theta \tan \theta - \int \tan \theta$ $\theta \sec^2 \theta \rightarrow \theta \tan \theta - \ln(\sec \theta)$ or $\theta \tan \theta + \ln(\cos \theta)$ or $\lambda \theta \sec^2 \theta \rightarrow \lambda \theta \tan \theta - \lambda \ln(\sec \theta)$ or $\lambda \theta \tan \theta + \lambda \ln(\cos \theta)$	M1 dM1 A1
	<b>Note:</b> Condone $\theta \sec^2 \theta \rightarrow \theta \tan \theta - \ln(\sec x)$ or $\theta \tan \theta + \ln(\cos x)$ for A1		
	$\left\{ \int \tan \theta \sec^2 \theta \, d\theta \right\}$ $= \frac{1}{2} \tan^2 \theta \text{ or } \frac{1}{2} \sec^2 \theta$ or $\frac{1}{2u^2}$ where $u = \cos \theta$ or $\frac{1}{2} u^2$ where $u = \tan \theta$	$\tan \theta \sec^2 \theta \text{ or } \lambda \tan \theta \sec^2 \theta \rightarrow \pm C \tan^2 \theta \text{ or } \pm C \sec^2 \theta$ or $\pm Cu^{-2}$ , where $u = \cos \theta$ $\tan \theta \sec^2 \theta \rightarrow \frac{1}{2} \tan^2 \theta \text{ or } \frac{1}{2} \sec^2 \theta \text{ or } \frac{1}{2cos^2\theta} \text{ or } \tan^2 \theta - \frac{1}{2} \sec^2 \theta$ or $0.5u^{-2}$ , where $u = \cos \theta$ or $0.5u^2$ , where $u = \tan \theta$ or $\lambda \tan \theta \sec^2 \theta \rightarrow \frac{\lambda}{2} \tan^2 \theta \text{ or } \frac{\lambda}{2} \sec^2 \theta \text{ or } \frac{\lambda}{2cos^2\theta}$ or $0.5\lambda u^{-2}$ , where $u = \cos \theta$ or $0.5\lambda u^2$ , where $u = \tan \theta$	M1 A1
	$\{ \text{Area}(R) \} = \left[ 3\theta \tan \theta - 3 \ln(\sec \theta) + \frac{3}{2} \tan^2 \theta \right]_0^{\frac{\pi}{3}}$ or $\left[ 3\theta \tan \theta - 3 \ln(\sec \theta) + \frac{3}{2} \sec^2 \theta \right]_0^{\frac{\pi}{3}}$ $= \left( 3\left(\frac{\pi}{3}\right) \sqrt{3} - 3 \ln 2 + \frac{3}{2}(3) \right) - (0) \text{ or } \left( 3\left(\frac{\pi}{3}\right) \sqrt{3} - 3 \ln 2 + \frac{3}{2}(4) \right) - \left( \frac{3}{2} \right)$ $= \frac{9}{2} + \sqrt{3}\pi - 3 \ln 2 \text{ or } \frac{9}{2} + \sqrt{3}\pi + 3 \ln\left(\frac{1}{2}\right) \text{ or } \frac{9}{2} + \sqrt{3}\pi - \ln 8 \text{ or } \ln\left(\frac{1}{8}e^{\frac{9}{2}+\sqrt{3}\pi}\right)$		
			[6]
			12



# Exponentials and Logarithms - modelling

- Remember – modelling can only ask you to substitute, solve, or comment – don't reinvent maths just because it is in context!
- If it asks for rate, differentiate
- For non-linear to linear, take logs of both sides – I've done loads of these before, so only covered in a Your Turn question.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



9. The amount of an antibiotic in the bloodstream, from a given dose, is modelled by the formula

$$x = D e^{-0.2t}$$

where  $x$  is the amount of the antibiotic in the bloodstream in milligrams,  $D$  is the dose given in milligrams and  $t$  is the time in hours after the antibiotic has been given.

A first dose of 15 mg of the antibiotic is given.

- (a) Use the model to find the amount of the antibiotic in the bloodstream 4 hours after the dose is given. Give your answer in mg to 3 decimal places. (2)

A second dose of 15 mg is given 5 hours after the first dose has been given. Using the same model for the second dose,

- (b) show that the **total** amount of the antibiotic in the bloodstream 2 hours after the second dose is given is 13.754 mg to 3 decimal places. (2)

No more doses of the antibiotic are given. At time  $T$  hours after the second dose is given, the total amount of the antibiotic in the bloodstream is 7.5 mg.

- (c) Show that  $T = a \ln\left(b + \frac{b}{e}\right)$ , where  $a$  and  $b$  are integers to be determined. (4)

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



9(a)	Subs $D = 15$ and $t = 4$ $x = 15e^{-0.2 \times 4} = 6.740$ (mg)	M1A1	
(b)	$15e^{-0.2 \times 7} + 15e^{-0.2 \times 2} = 13.754$ (mg)	M1A1*	(2)
(c)	$15e^{-0.2 \times T} + 15e^{-0.2 \times (T+5)} = 7.5$ $15e^{-0.2 \times T} + 15e^{-0.2 \times T} e^{-1} = 7.5$ $15e^{-0.2 \times T} (1+e^{-1}) = 7.5 \Rightarrow e^{-0.2 \times T} = \frac{7.5}{15(1+e^{-1})}$ $T = -5 \ln \left( \frac{7.5}{15(1+e^{-1})} \right) = 5 \ln \left( 2 + \frac{2}{e} \right)$	M1  dM1  A1, A1	(2)  (4)  (8 marks)



- 6 A mobile phone company records their annual sales on 31<sup>st</sup> December every year.

Paul thinks that the annual sales,  $S$  million, can be modelled by the equation  $S = ab^t$ , where  $a$  and  $b$  are both positive constants and  $t$  is the number of years since 31<sup>st</sup> December 2015.

Paul tests his theory by using the annual sales figures from 31<sup>st</sup> December 2015 to 31<sup>st</sup> December 2019. He plots these results on a graph, with  $t$  on the horizontal axis and  $\log_{10}S$  on the vertical axis.

- (a) Explain why, if Paul's model is correct, the results should lie on a straight line of best fit on his graph. [3]

The results lie on a straight line of best fit which has a gradient of 0.146 and an intercept on the vertical axis of 0.583.

- (b) Use these values to obtain estimates for  $a$  and  $b$ , correct to 2 significant figures. [2]

- (c) Use this model to predict the year in which, on the 31<sup>st</sup> December, the annual sales would first be recorded as greater than 200 million. [3]

- (d) Give a reason why this prediction may not be reliable. [1]

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



6	(a)	$\log_{10}S = \log_{10}(ab^t)$	<b>M1</b>	<b>2.1</b>	Attempt to show reduction to linear form
		$\log_{10}S = \log_{10}a + \log_{10}b^t$			
		$\log_{10}S = t\log_{10}b + \log_{10}a$			Obtain correct equation
		which is of the form $Y = mX + c$	<b>A1</b>	<b>2.4</b>	Link to equation of straight line
				[3]	
	(b)	$\log_{10}a = 0.583 \Rightarrow a = 10^{0.583} = 3.8$	<b>B1</b>	<b>1.1</b>	Obtain $a = 3.8$ , or better, from either eqn
		$\log_{10}b = 0.146 \Rightarrow b = 10^{0.146} = 1.4$	<b>B1</b>	<b>1.1</b>	Obtain $b = 1.4$ , or better, from either eqn
				[2]	
	(c)	$3.8 \times 1.4^t = 200$ $1.4^t = 52.63$	<b>M1</b>	<b>3.1a</b>	Link their model to 200 and attempt to solve for $t$
		$t = 11.8$	<b>A1</b>	<b>1.1</b>	Obtain $t = 11.8$ , or better, www (allow $t = 12$ )
		so year is 2027	<b>A1FT</b>	<b>3.2a</b>	FT their value for $t$
	(d)	Unlikely that sales will continue at same rate Finite market	<b>B1</b>	<b>3.5b</b>	Any sensible reason – eg pattern not necessarily continuing or the market being limited by no. of customers
				[1]	



Sorry if another rabbit population question is a little triggering...

The number of rabbits on an island is modelled by the equation

$$P = \frac{100e^{-0.1t}}{1 + 3e^{-0.9t}} + 40, \quad t \in \mathbb{R}, t \geq 0$$

where  $P$  is the number of rabbits,  $t$  years after they were introduced onto the island

A sketch of the graph of  $P$  against  $t$  is shown in Figure 3.

- (a) Calculate the number of rabbits that were introduced onto the island.

(1)

(b) Find  $\frac{dP}{dt}$

(3)

The number of rabbits initially increases, reaching a maximum value  $P_T$  when  $t = T$

- (c) Using your answer from part (b), calculate

(i) the value of  $T$  to 2 decimal places,

(ii) the value of  $P_T$  to the nearest integer.

*(Solutions based entirely on graphical or numerical methods are not acceptable.)*

(4)

For  $t > T$ , the number of rabbits decreases, as shown in Figure 3, but never falls below  $k$ , where  $k$  is a positive constant.

- (d) Use the model to state the maximum value of  $k$ .

(1)



Figure 3

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



8 (a)	$P_0 = \frac{100}{1+3} + 40 = 65$ $\frac{d}{dt} e^{kt} = C e^{kt}$	B1 (1) M1 M1 A1 (3)
(b)	$\frac{dP}{dt} = \frac{(1+3e^{-0.9t}) \times -10e^{-0.1t} - 100e^{-0.1t} \times -2.7e^{-0.9t}}{(1+3e^{-0.9t})^2}$	
(c)(i)	<p>At maximum <math>-10e^{-0.1t} - 30e^{-0.1t} \times e^{-0.9t} + 270e^{-0.1t} \times e^{-0.9t} = 0</math></p> $e^{-0.1t} (-10 + 240e^{-0.9t}) = 0$ $e^{-0.9t} = \frac{10}{240} \quad \text{or } e^{0.9t} = 24$	M1
(c) (ii)	$-0.9t = \ln\left(\frac{1}{24}\right) \Rightarrow t = \frac{10}{9} \ln(24) = 3.53$	M1, A1
(d)	$\text{Sub } t = 3.53 \Rightarrow P_T = 102$	A1 (4)
	40	B1 (1)
		<b>9 marks</b>



5. A bath is filled with hot water. The temperature,  $\theta^\circ\text{C}$ , of the water in the bath,  $t$  minutes after the bath has been filled, is given by

$$\theta = 20 + Ae^{-kt}$$

where  $A$  and  $k$  are positive constants.

Given that the temperature of the water in the bath is initially  $38^\circ\text{C}$ ,

- (a) find the value of  $A$ .

(2)

The temperature of the water in the bath 16 minutes after the bath has been filled is  $24.5^\circ\text{C}$ .

- (b) Show that  $k = \frac{1}{8} \ln 2$

(4)

Using the values for  $k$  and  $A$ ,

- (c) find the temperature of the water 40 minutes after the bath has been filled, giving your answer to 3 significant figures.

(2)

- (d) Explain why the temperature of the water in the bath cannot fall to  $19^\circ\text{C}$ .

(1)

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---





DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

<b>5(a)</b>	$t = 0, \theta = 38 \Rightarrow 38 = 20 + Ae^{-k \cdot 0}$	For substituting $t = 0$ and $\theta = 38$ into $\theta = 20 + Ae^{-kt}$	M1
	$\Rightarrow A = 18$	Correct value for $A$	A1
	<b><math>A = 18</math> with no working scores both marks</b>		(2)
<b>(b)</b>	$t = 16, \theta = 24.5 \Rightarrow 24.5 = 20 + "18"e^{-k \cdot 16}$	For substituting $t = 16$ and $\theta = 24.5$ into $\theta = 20 +$ their " $A$ " $e^{-kt}$	M1
	$\Rightarrow 18e^{-k \cdot 16} = 4.5$ or $e^{-k \cdot 16} = \frac{1}{4}$	This mark is for a correct equation with the <b>constants combined</b> . Allow equivalent correct equations e.g. $e^{16k} = 4$	A1
	$\Rightarrow e^{16k} = 4 \Rightarrow 16k = \ln 4$ or $\Rightarrow \ln 18e^{-k \cdot 16} = \ln 4.5 \Rightarrow \ln 18 + \ln e^{-k \cdot 16} = \ln 4.5 \Rightarrow \ln e^{-k \cdot 16} = \ln \frac{1}{4}$ $\Rightarrow -16k = \ln \frac{1}{4}$		M1
	Uses correct log or exponential work to move from: $e^{\pm nk} = C$ to $\pm nk = \alpha \ln C$ or $pe^{\pm nk} = q$ to $\pm nk = \alpha \ln \beta$		<b>5(d)</b>
	$-16k = \ln \frac{1}{4} \Rightarrow k = -\frac{1}{16} \ln \frac{1}{4} = \frac{1}{8} \ln 2 *$ Shows that $k = \frac{1}{8} \ln 2$		
<b>(b)</b>	There must be <b>at least one intermediate line</b> between their $\pm nk = \alpha \ln C$ or their $\pm nk = \alpha \ln \beta$ and the printed answer.	A1*	
	So for example $-16k = \ln \frac{1}{4} \Rightarrow k = \frac{1}{8} \ln 2 *$ scores A0 as there is no intermediate line.		
	Note: The marks in part (b) can be scored by using $\theta = 20 + Ae^{-kt}$ and substituting 2 out of: $A = 18$ , $\theta = 24.5$ , $k = \frac{1}{8} \ln 2$ to show that the 3rd variable is correct followed by a conclusion e.g. so $k = \frac{1}{8} \ln 2$		
<b>(c)</b>			(4)
	$t = 40 \Rightarrow \theta = 20 + "18"e^{-\frac{1}{8} \ln 2 \cdot 40}$	Substitutes $t = 40$ into the given equation with their $A$ and the given value of $k$ to obtain a value for $\theta$	M1
	$\Rightarrow \theta = \text{awrt } 20.6 (\text{ }^\circ\text{C})$	Awrt 20.6	A1
	<b>Correct answer only scores both marks</b>		(2)

- Examples:**
  - The lower limit is 20
  - $\theta > 20$
  - As  $t$  tends to infinity temperature tends to 20
  - The temperature cannot go below 20
  - $e^{-kt}$  tends towards zero so the temperature tends to 20
  - $e^{-kt}$  is always positive so the temperature is always bigger than 20
  - Substitutes  $\theta = 19$  in  $\theta = 20 + "18"e^{-kt}$  (may be implied by e.g.  $e^{-kt} = -\frac{1}{18}$ ) and states e.g. that you cannot find the log of a negative number or "which is not possible"

Do not accept  $e^{-kt}$  cannot be negative without reference to the "20"

(1)

B1



3. The value of a car is modelled by the formula

$$V = 16000e^{-kt} + A, \quad t \geq 0, t \in \mathbb{R}$$

where  $V$  is the value of the car in pounds,  $t$  is the age of the car in years, and  $k$  and  $A$  are positive constants.

Given that the value of the car is £17 500 when new and £13 500 two years later,

- (a) find the value of  $A$ ,

(1)

- (b) show that  $k = \ln\left(\frac{2}{\sqrt{3}}\right)$

(4)

- (c) Find the age of the car, in years, when the value of the car is £6000

Give your answer to 2 decimal places.

(4)

DO NOT WRITE IN THIS AREA

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



<b>3(a)</b> $A = 1500$	B1 <b>(1)</b>
<b>(b)</b> Sub $t = 2, V = 13500 \Rightarrow 16000e^{-2k} = 12000$ $\Rightarrow e^{-2k} = \frac{3}{4}$ 0.75 oe $\Rightarrow k = -\frac{1}{2} \ln \frac{3}{4}, = \ln \sqrt{\frac{4}{3}} = \ln \left( \frac{2}{\sqrt{3}} \right)$	M1 A1 dM1, A1* <b>(4)</b>
<b>(c)</b> Sub $6000 = 16000e^{-\ln(\frac{2}{\sqrt{3}})T} + 1500 \Rightarrow e^{-\ln(\frac{2}{\sqrt{3}})T} = C$ $\Rightarrow e^{-\ln(\frac{2}{\sqrt{3}})T} = \frac{45}{160} = 0.28125$ $\Rightarrow T = -\frac{\ln(\frac{45}{160})}{\ln(\frac{2}{\sqrt{3}})} = 8.82$	M1 A1 M1 A1 <b>(4)</b> <b>(9 marks)</b>



- 11 In a science experiment a substance is decaying exponentially. Its mass,  $M$  grams, at time  $t$  minutes is given by  $M = 300e^{-0.05t}$ .

- (i) Find the time taken for the mass to decrease to half of its original value. [3]

A second substance is also decaying exponentially. Initially its mass was 400 grams and, after 10 minutes, its mass was 320 grams.

- (ii) Find the time at which both substances are decaying at the same rate. [8]

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



11	(i)	When $t = 0, M = 300$	B1	2.2a	Identify that the initial mass is 300g
		$300e^{-0.05t} = 150$ $e^{-0.05t} = 0.5$ $-0.05t = \ln 0.5$	M1	3.1a	Equate to 150 and attempt to solve
		$t = 13.9$ (minutes)	A1	1.1	Obtain 13.86, or better
			[3]		
	(ii)	$M_2 = 400e^{kt}$	B1	2.2a	State or imply $400e^{kt}$
		$320 = 400e^{10k}$ $k = 0.1\ln 0.8$	M1	1.1a	Attempt to find $k$
		$M_2 = 400e^{-0.0223t}$	A1	1.1	Obtain correct expression for mass of second substance
		Substance 1: $\frac{dM_1}{dt} = -15e^{-0.05t}$ Substance 2: $\frac{dM_2}{dt} = -8.93e^{-0.0223t}$	M1	3.1a	Attempt differentiation at least once
			A1ft	1.1	Both derivatives correct
		$-15e^{-0.05t} = -8.93e^{-0.0223t}$ $e^{0.0277t} = 1.681$	M1	3.1a	Equate derivatives and rearrange as far as $e^{f(t)} = c$
		$0.0277t = 0.519$	M1	1.1	Attempt to solve equation of form $e^{f(t)} = c$
		time = 18.75 minutes	A1	3.2a	Obtain correct value for $t$ Allow 18.7, 18.8 or 19 mins



# Exponentials and Logarithms - equations

- Follow the log laws! You should know these by now...
  - Adding logs... multiply the input
  - Subtracting logs... divide the input
  - Factor of a log... becomes input's power
    - And vice versa!
- Note: check your answers, your input cannot be negative for a logarithm

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



2. Find the exact solutions, in their simplest form, to the equations

(a)  $e^{3x-9} = 8$

(3)

(b)  $\ln(2y + 5) = 2 + \ln(4 - y)$

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



2.(a)	$\begin{aligned} e^{3x-9} &= 8 \Rightarrow 3x - 9 = \ln 8 \\ \Rightarrow x &= \frac{\ln 8 + 9}{3}, = \ln 2 + 3 \end{aligned}$
(b)	$\begin{aligned} \ln(2y+5) &= 2 + \ln(4-y) \\ \ln\left(\frac{2y+5}{4-y}\right) &= 2 \\ \left(\frac{2y+5}{4-y}\right) &= e^2 \\ 2y+5 &= e^2(4-y) \Rightarrow 2y + e^2y = 4e^2 - 5 \Rightarrow y = \frac{4e^2 - 5}{2 + e^2} \end{aligned}$



(ii) Given that

$$2 \log_4(3x + 5) = \log_4(3x + 8) + 1, \quad x > -\frac{5}{3}$$

(a) show that

$$9x^2 + 18x - 7 = 0 \quad (4)$$

(b) Hence solve the equation

$$2 \log_4(3x + 5) = \log_4(3x + 8) + 1, \quad x > -\frac{5}{3} \quad (2)$$

DO NOT WRITE IN THIS AREA

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



	(ii) (a)	<p>Ignore labels (a) and (b) in part ii and mark work as seen</p> <p><math>\log_4(3x+5)^2 =</math> <span style="float: right;">Applies power law of logarithms</span></p> <p>Uses <math>\log_4 4 = 1</math> or <math>4^1 = 4</math></p> <p>Uses quotient or product rule so e.g. <math>\log(3x+5)^2 = \log 4(3x+8)</math> or <math>\log \frac{(3x+5)^2}{(3x+8)} = 1</math></p> <p>Obtains with no errors <math>9x^2 + 18x - 7 = 0 *</math></p>	M1 M1 M1 A1* cso (4)
	(b)	<p>Solves given or “their” quadratic equation by any of the standard methods</p> <p>Obtains <math>x = \frac{1}{3}</math> and <math>-\frac{7}{3}</math> and rejects <math>-\frac{7}{3}</math> to give just <math>\frac{1}{3}</math></p>	M1 A1 <b>(2)</b> <b>[8]</b>



7. (i)  $2 \log(x + a) = \log(16a^6)$ , where  $a$  is a positive constant

Find  $x$  in terms of  $a$ , giving your answer in its simplest form.

(3)

(ii)  $\log_3(9y + b) - \log_3(2y - b) = 2$ , where  $b$  is a positive constant

Find  $y$  in terms of  $b$ , giving your answer in its simplest form.

(4)

DO NOT WRITE IN THIS AREA

Handwriting practice lines for the word "BUCÉN MATH". There are 10 sets of horizontal lines for tracing.



	Use of power rule so $\log(x+a)^2 = \log 16a^6$ or $2\log(x+a) = 2\log 4a^3$ or $\log(x+a) = \log(16a^6)^{\frac{1}{2}}$ Removes logs and square roots, <b>or</b> halves then removes logs to give $(x+a) = 4a^3$ Or $x^2 + 2ax + a^2 - 16a^6 = 0$ followed by factorisation or formula to give $x = \sqrt{16a^6} - a$ $(x =) 4a^3 - a$ (depends on previous M's and must be this expression or equivalent)	M1 M1 M1 A1cao (3)
(ii) Way 1	$\log_3 \frac{(9y+b)}{(2y-b)} = 2$ $\frac{(9y+b)}{(2y-b)} = 3^2$ $(9y+b) = 9(2y-b) \Rightarrow y =$ $y = \frac{10}{9}b$	Applies quotient law of logarithms Uses $\log_3 3^2 = 2$ Multiplies across and makes y the subject A1cs (4)
Way 2	Or : $\log_3(9y+b) = \log_3 9 + \log_3(2y-b)$ $\log_3(9y+b) = \log_3 9(2y-b)$ $(9y+b) = 9(2y-b) \Rightarrow y = \frac{10}{9}b$	2 <sup>nd</sup> M mark 1 <sup>st</sup> M mark Multiplies across and makes y the subject A1cs (4) [7]