



# OCR A Level Physics



Your notes

## Ideal Gases

### Contents

- \* The Avogadro Constant
- \* Ideal Gas Equation
- \* Gas Laws



Your notes

## The Avogadro Constant

### Avogadro's Constant

- The atomic mass unit (u) is approximately the mass of a proton or neutron =  $1.66 \times 10^{-27}$  kg
- This means that the mass of an atom or molecule can be calculated using the number of protons and neutrons it contains
  - For example, a carbon-12 atom has a mass of:

$$12 \text{ u} = 12 \times 1.66 \times 10^{-27} = 1.99 \times 10^{-26} \text{ kg}$$

### The Mole

- In thermodynamics, the amount of substance is measured in the SI unit 'mole'
  - This has the symbol **mol**
  - The mole is a unit of **substance**, not a unit of mass

- The mole is defined as:

**The SI base unit of an 'amount of substance'. It is the amount containing as many particles (e.g., atoms or molecules) as there are atoms in 12 g of carbon-12**

- The mole is an important unit in thermodynamics
- If we consider the number of moles of two different gases under the same conditions, their physical properties are the same
- One mole of a substance is defined as the number of molecules in exactly 12 g of carbon:

$$1 \text{ mole} = \frac{0.012}{1.99 \times 10^{-26}} = 6.02 \times 10^{23} \text{ molecules}$$

### Avogadro's Constant

Avogadro's constant ( $N_A$ ) is defined as:

**The number of atoms of carbon-12 in 12 g of carbon-12; equal to  $6.02 \times 10^{23} \text{ mol}^{-1}$**

- For example, 1 mole of sodium (Na) contains  $6.02 \times 10^{23}$  **atoms** of sodium
- The number of atoms can be determined if the number of moles is known by multiplying by  $N_A$ .

- For example: 2.0 mol of nitrogen contains:  $2.0 \times N_A = 2.0 \times 6.02 \times 10^{23} = 1.20 \times 10^{24}$  atoms

## Molar Mass

- The molar mass of a substance is the mass, in grams, in one mole
  - Its unit is **g mol<sup>-1</sup>**
- The number of moles from this can be calculated using the equation:

$$\text{Number of moles} = \frac{\text{mass (g)}}{\text{molar mass (g mol}^{-1}\text{)}}$$



Your notes



Your notes

## Ideal Gas Equation

### Ideal Gas Equation

- An ideal gas is a specific type of gas which:
  - Has molecules with negligible volume
  - Collisions which are elastic
  - Cannot be liquified
  - Has no interactions between the molecules (except during collisions)
  - Obeys the (ideal) gas laws (Boyles law, Charles' law and Pressure law)
- All of these can occur at any temperature or pressure
- The ideal gas equation for number of moles can be expressed as:




Diagram illustrating the ideal gas equation  $pV = nRT$  with variables in boxes and arrows pointing to them:

- VOLUME ( $\text{m}^3$ ) points to  $V$
- PRESSURE (Pa) points to  $p$
- MOLAR GAS CONSTANT =  $8.31 \text{ J kg}^{-1} \text{ mol}^{-1}$  points to  $R$
- TEMPERATURE (K) points to  $T$
- NUMBER OF MOLES (mol) points to  $n$

Copyright © Save My Exams. All Rights Reserved

- The ideal gas equation for number of molecules can also be written in the form:




Diagram illustrating the ideal gas equation  $pV = NkT$  with variables in boxes and arrows pointing to them:

- VOLUME ( $\text{m}^3$ ) points to  $V$
- PRESSURE (Pa) points to  $p$
- BOLTZMANN CONSTANT =  $1.38 \times 10^{-23} \text{ J K}^{-1}$  points to  $k$
- TEMPERATURE (K) points to  $T$
- NUMBER OF MOLECULES points to  $N$

Copyright © Save My Exams. All Rights Reserved

- An ideal gas is therefore defined as:

**A gas which obeys the equation of state  $pV = nRT$  at all pressures, volumes and temperatures**



Your notes



### Worked Example

A storage cylinder of an ideal gas has a volume of  $8.3 \times 10^3 \text{ cm}^3$ . The gas is at a temperature of  $15^\circ\text{C}$  and a pressure of  $4.5 \times 10^7 \text{ Pa}$ .

Calculate the amount of gas in the cylinder, in moles.

**Answer:**

#### Step 1: State the known quantities

- Pressure,  $P = 4.5 \times 10^7 \text{ Pa}$
- Volume,  $V = 8.3 \times 10^3 \text{ cm}^3 = 8.3 \times 10^3 \times 10^{-6} = 8.3 \times 10^{-3} \text{ m}^3$
- Temperature,  $T = 15^\circ\text{C} = 15 + 273 = 288 \text{ K}$

#### Step 2: Write down the ideal gas equation

$$pV = nRT$$

#### Step 3: Rearrange for the number of moles $n$

$$n = \frac{pV}{RT}$$

#### Step 4: Substitute in values and calculate number of moles of gas

$$n = \frac{4.5 \times 10^7 \times 8.3 \times 10^{-3}}{8.31 \times 288} = 160 \text{ mol}$$



### Examiner Tips and Tricks

Don't worry about remembering the values of  $R$  and  $k$ , they will both be given in the equation sheet in your exam.



Your notes

## Gas Laws


# Investigating Gas Laws

## Boyle's Law

- If the temperature  $T$  of an ideal gas is constant, then **Boyle's Law** is given by:

$$P \propto \frac{1}{V}$$

- This means the pressure is **inversely proportional** to the volume of a gas



Copyright © Save My Exams. All Rights Reserved



**Pressure increases when a gas is compressed**

- The relationship between the pressure and volume for a fixed mass of gas at constant temperature can also be written as:

$$P_1V_1 = P_2V_2$$

- Where:
  - $P_1$  = initial pressure (Pa)
  - $P_2$  = final pressure (Pa)
  - $V_1$  = initial volume ( $\text{m}^3$ )

- $V_2$  = final volume ( $\text{m}^3$ )



### **Boyle's Law graph representing pressure inversely proportional to volume**

- If the temperature increases, the graph is further from the origin and vice versa

## **Pressure Law**

- If the volume  $V$  of an ideal gas is constant, the **Pressure law** is given by:



$$P \propto T$$

- This means the pressure is **proportional** to the temperature
- The relationship between the pressure and thermodynamic temperature for a fixed mass of gas at constant volume can also be written as:

$$\frac{P_1}{T_1} = \frac{P_2}{T_2}$$

- Where:
  - $P_1$  = initial pressure (Pa)
  - $P_2$  = final pressure (Pa)
  - $T_1$  = initial temperature (K)

- $T_2$  = final temperature (K)



**Pressure Law graph representing temperature (in °C) directly proportional to the volume**



### Worked Example

The pressure inside a bicycle tyre is  $5.10 \times 10^5$  Pa when the temperature is 279 K. After the bicycle has been ridden, the temperature of the air in the tyre is 299 K.

Calculate the new pressure in the tyre, assuming the volume is unchanged.

**Answer:**





Your notes

**Step 1: Choose which ideal gas law to use**

Since the volume is constant, the pressure law must be used

$$\frac{P_1}{T_1} = \frac{P_2}{T_2}$$

**Step 2: Write down the known quantities**

$$P_1 = 5.10 \times 10^5 \text{ Pa}$$

$$P_2 = ?$$

$$T_1 = 279 \text{ K}$$

$$T_2 = 299 \text{ K}$$

**Step 3: Substitute values into pressure law equation**

$$P_2 = \frac{P_1 T_2}{T_1} = \frac{(5.10 \times 10^5) \times 299}{279} = 5.47 \times 10^5 \text{ Pa}$$

**Examiner Tips and Tricks**

Remember when using any ideal gas law, including the ideal gas equation, the temperature  $T$  must always be in **kelvin** (K)

## Estimating Absolute Zero

- The value of **absolute zero** can be estimated by using the gas law **Charles' Law**
- If the pressure  $P$  of an ideal gas is constant, then **Charles's law** is given by:

$$V \propto T$$

- This means the volume is **proportional** to the temperature of a gas
- The relationship between the volume and thermodynamic temperature for a fixed mass of gas at constant pressure can also be written as:


$$\frac{V_1}{T_1} = \frac{V_2}{T_2}$$



Your notes

## ■ Where:

- $V_1$  = initial volume ( $\text{m}^3$ )
- $V_2$  = final volume ( $\text{m}^3$ )
- $T_1$  = initial temperature (K)
- $T_2$  = final temperature (K)

**Charles's Law graph representing temperature (in °C) directly proportional to the volume**

- The Charles's Law graph for temperature in **Kelvin** against volume is identical except that it is a straight line through the **origin**
  - Extrapolating backwards leads to a temperature of **absolute zero** at which the gas will have a **volume** of zero  $\text{m}^3$