

# Introduction to Deep Learning for Computer Vision

## Assignment 3: Simple Linear Classifier II

Paul Molina-Plant

January 30, 2019

### **Abstract**

In this assignment, I finished implementing a linear classifier. I derived the gradient for multinomial logistic regression and implemented it. I also finished implementing the training routine and cross validation.

# 1 Derivation of the gradient for cross entropy

## 1.1 The cross entropy function

$$L_i = -\log \frac{\sum_j t_{ij} e^{s_{ij}}}{\sum_k e^{s_{ik}}} \quad (1)$$

I assume  $\log$  base  $e$  for (1).

$$t_{ij} = \delta_{ij} = \begin{cases} 1 & j = y_i \\ 0 & j \neq y_i \end{cases} \quad (2)$$

Softmax  $P_{ij}$

$$P_{ij} = \frac{e^{s_{ij}}}{\sum_k e^{s_{ik}}} \quad (3)$$

Among the  $k$  classes,  $j = y_i$  for exactly one of them. Therefore, we can ignore the other  $k - 1$  terms in the sum of (1).

$$L_i = -\ln \frac{e^{s_{ij}}}{\sum_k e^{s_{ik}}} = -\ln P_{ij} \quad (4)$$

## 1.2 Gradient of $W$

Computing the gradient  $\frac{\partial L_i}{\partial W}$  via chain rule.

$$\frac{\partial L_i}{\partial W} = \frac{\partial L_i}{\partial P_{ij}} \frac{\partial P_{ij}}{\partial s_{ij}} \frac{\partial s_{ij}}{\partial W}$$

Linear classifier  $f(x_i, W) = s_{ij}$ .

$$s_{ij} = Wx_i + b_i \quad (5)$$

The partial derivative of  $s_{ij}$  wrt  $W$  is nonzero when  $j = y_i$  and zero everywhere else. This condition is expressed with  $\delta_{ij}$  (2).

$$\frac{\partial s_{ij}}{\partial W} = \delta_{ij} x_i$$

The partial derivative of  $P_{ij}$  (3) wrt  $W$  by applying the chain rule.

$$\begin{aligned} \frac{\partial P_{ij}}{\partial W} &= \frac{\partial}{\partial W} \frac{e^{s_{ij}}}{\sum_k e^{s_{ik}}} = \frac{\delta_{ij} x_i e^{s_{ij}} - e^{s_{ij}} x_i e^{s_{ij}}}{(\sum_k e^{s_{ik}})^2} \\ &= \left( \frac{e^{s_{ij}}}{\sum_k e^{s_{ik}}} \right) \left( \frac{\delta_{ij} x_i \sum_k e^{s_{ik}} - x_i e^{s_{ij}}}{\sum_k e^{s_{ik}}} \right) \\ &= P_{ij} (\delta_{ij} x_i - P_{ij} x_i) \\ &= P_{ij} x_i (\delta_{ij} - P_{ij}) \end{aligned}$$

Finally the partial derivative of  $L_i$  (4) wrt  $W$ .

$$\begin{aligned} \frac{\partial L_i}{\partial W} &= \frac{\partial}{\partial W} (-\ln P_{ij}) \\ &= -\frac{1}{P_{ij}} P_{ij} x_i (\delta_{ij} - P_{ij}) \\ &= (P_{ij} - \delta_{ij}) x_i \end{aligned} \quad (6)$$

### 1.3 Gradient of $b_i$

Computing the gradient  $\frac{\partial L_i}{\partial b_i}$  via chain rule.

$$\frac{\partial L_i}{\partial b_i} = \frac{\partial L_i}{\partial P_{ij}} \frac{\partial P_{ij}}{\partial s_{ij}} \frac{\partial s_{ij}}{\partial b_i}$$

The partial derivative of  $s_{ij}$  (5) wrt  $b_i$  is nonzero when  $j = y_i$  and zero everywhere else. This condition is expressed with  $\delta_{ij}$  (2).

$$\frac{\partial s_{ij}}{\partial W} = \delta_{ij}$$

The partial derivative of  $P_{ij}$  (3) wrt  $b_i$  by applying the chain rule.

$$\frac{\partial P_{ij}}{\partial b_i} = \frac{\partial}{\partial W} \frac{e^{s_{ij}}}{\sum_k e^{s_{ik}}} = \frac{\delta_{ij} e^{s_{ij}} \sum_k e^{s_{ik}} - e^{s_{ij}} e^{s_{ij}}}{(\sum_k e^{s_{ik}})^2} = \delta_{ij} P_{ij} - P_{ij}^2$$

Finally the partial derivative of  $L_i$  (4) wrt  $b_i$ .

$$\begin{aligned} \frac{\partial L_i}{\partial b_i} &= \frac{\partial}{\partial b_i} (-\ln P_{ij}) \\ &= -\frac{1}{P_{ij}} (\delta_{ij} P_{ij} - P_{ij}^2) \\ &= P_{ij} - \delta_{ij} \end{aligned} \tag{7}$$

### 1.4 Summary of derivations

The weights gradient was,

$$\frac{\partial L_i}{\partial W} = (P_{ij} - \delta_{ij}) x_i$$

The bias gradient was,

$$\frac{\partial L_i}{\partial b_i} = P_{ij} - \delta_{ij}$$

## 2 Training curves

The following curves were obtained by running the HOG model on cifar-10 dataset with default parameters.

### 2.1 Linear SVM

Best validation accuracy: 46.58%

Test accuracy: 46.47%




Figure 1: Validation Accuracy.




Figure 2: Validation Loss.

## 2.2 Logistic Regression

Best validation accuracy: 42.60%

Test accuracy: 42.45%




Figure 3: Validation Accuracy.




Figure 4: Validation Loss.



Figure 5: Example caption.

Example Section.

## 2.3 Example Subsection

Example Sub Section.

Example paragraph.

## 2.4 Example Equation

This is how a equation looks like.

$$y = ax^2 + bx + c \quad , \quad (8)$$

where an inline equation looks like  $a = b$ .

## 2.5 Example Figure

To put a figure, you can do as shown in Fig. 5.