(a)


Fan DKD bulk RNA-seq
P2RY6 vs PDGFRB
Normal (n=9), Early DKD (n=6), Adv. DKD (n=21)
rho = 0.62 [0.37, 0.79]
BH-adjusted p < 0.001
PDGFRB (log2)
P2RY6 (log2)
Normal
Early DKD
Adv. DKD

(b)


KPMP scRNA
Stroma P2RY6 vs PDGFRB (adjudicated)
n = 92 stroma samples
rho = -0.20 [-0.39, 0.00]
BH-adjusted p = 0.068
Stroma PDGFRB (pseudobulk)
Stroma P2RY6 (pseudobulk)
Adjudicated
Diabetic Kidney Disease
Hypertensive Kidney Disease
Cannot be determined
Other
Healthy Reference Tissue
Acute Kidney Injury

(c)


KPMP scRNA (DKD-focused)
Stroma P2RY6 vs PDGFRB
n = 66 stroma samples
rho = -0.21 [-0.43, 0.04]
BH-adjusted p = 0.093
Stroma PDGFRB (pseudobulk)
Stroma P2RY6 (pseudobulk)
Healthy Reference Tissue
Acute Kidney Injury
Chronic Kidney Disease

(d)


KPMP snRNA
FB P2RY6 vs PDGFRB
n = 130 fibroblast patients
rho = 0.23 [0.06, 0.39]
BH-adjusted p = 0.014
FB PDGFRB (pseudobulk)
FB P2RY6 (pseudobulk)
Healthy Reference
AKI
CKD

(e)


KPMP snRNA (DKD-focused)
FB P2RY6 vs PDGFRB
n = 94 fibroblast patients
rho = 0.34 [0.14, 0.51]
BH-adjusted p = 0.002
FB PDGFRB (pseudobulk)
FB P2RY6 (pseudobulk)
Healthy Reference
AKI
CKD