



Stochastic Processes

DTMCs: Some Important Results and Their Proofs, Examples
Problems

Karthik P. N.


Assistant Professor, Department of AI

Email: pnkarthik@ai.iith.ac.in

04/08 April 2025

Example

- Consider a DTMC with the following transition graph.



1. Is this Markov chain irreducible?
2. Does there exist a unique stationary distribution?

Some Important Results – 1

Lemma (Regarding a Transient State)

Consider a time-homogeneous DTMC on a discrete state space \mathcal{X} with TPM P . Let $x \in \mathcal{X}$ be **transient**. Then,

$$\lim_{n \rightarrow \infty} P_{x,x}^n = 0.$$

Furthermore,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n P_{x,x}^k = 0.$$

Some Important Results – 1

Lemma (Regarding a Transient State)

Consider a time-homogeneous DTMC on a discrete state space \mathcal{X} with TPM P . Let $x \in \mathcal{X}$ be **transient**. Then,

$$\lim_{n \rightarrow \infty} P_{x,x}^n = 0.$$

Furthermore,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n P_{x,x}^k = 0.$$

Some Important Results – 2

Lemma (Regarding a Recurrent State)

Consider a time-homogeneous DTMC on a discrete state space \mathcal{X} and TPM P . Let $x \in \mathcal{X}$ be **recurrent**. Then,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n P_{x,x}^k = \frac{1}{\mu_{xx}}.$$



Proof of Lemma – Overview

- The proof of consists of two parts

Proof of Lemma – Overview

- The proof of consists of two parts
- In the first part, we show that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n P_{x,x}^k \geq \frac{1}{\mu_{xx}}.$$

Proof of Lemma – Overview

- The proof consists of two parts
- In the first part, we show that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n P_{x,x}^k \geq \frac{1}{\mu_{xx}}.$$

- In the second part, we show that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n P_{x,x}^k \leq \frac{1}{\mu_{xx}}.$$



Proof of Lemma – Lower Bound

- Assume that $X_0 = x$

Proof of Lemma – Lower Bound

- Assume that $X_0 = x$
- Let $N_x(n)$ denote the number of visits to state x up to time n , i.e.,

$$N_x(n) := \sum_{k=1}^n \mathbf{1}_{\{X_k=x\}}.$$

Proof of Lemma – Lower Bound

- Assume that $X_0 = x$
- Let $N_x(n)$ denote the number of visits to state x up to time n , i.e.,

$$N_x(n) := \sum_{k=1}^n \mathbf{1}_{\{X_k=x\}}.$$

- Let $\tau_x^{(0)} := 0$, and let

$$\tau_x^{(k)} = \inf\{n > \tau_x^{(k-1)} : X_n = x\}, \quad k \in \mathbb{N},$$

denote the time instant of k th visit to x

Proof of Lemma – Lower Bound

- Assume that $X_0 = x$
- Let $N_x(n)$ denote the number of visits to state x up to time n , i.e.,

$$N_x(n) := \sum_{k=1}^n \mathbf{1}_{\{X_k=x\}}.$$


- Let $\tau_x^{(0)} := 0$, and let

$$\tau_x^{(k)} = \inf\{n > \tau_x^{(k-1)} : X_n = x\}, \quad k \in \mathbb{N},$$

denote the time instant of k th visit to x

- Because x is recurrent, $\mathbb{P}(\tau_x^{(k)} < +\infty) = 1$ for all $k \in \mathbb{N}$

Proof of Lemma – Lower Bound




- Consider $H_k = \tau_x^{(k)} - \tau_x^{(k-1)}$, $k \in \mathbb{N}$
- From IID block structure, we know that $\{H_k\}_{k \in \mathbb{N}}$ is an IID process

- **Claim:**

For each $n \in \mathbb{N}$, the random variable $N_x(n) + 1$ is a stopping time w.r.t. $\{H_k\}_{k \in \mathbb{N}}$

Proof of Lemma – Lower Bound




- Consider $H_k = \tau_x^{(k)} - \tau_x^{(k-1)}$, $k \in \mathbb{N}$
- From IID block structure, we know that $\{H_k\}_{k \in \mathbb{N}}$ is an IID process
- **Claim:**

For each $n \in \mathbb{N}$, the random variable $N_x(n) + 1$ is a stopping time w.r.t. $\{H_k\}_{k \in \mathbb{N}}$

$$\{N_x(n) + 1 = m\} = \{N_x(n) = m - 1\} = \left\{ \sum_{k=1}^{m-1} H_k \leq n < \sum_{k=1}^m H_k \right\}$$

Proof of Lemma – Lower Bound



- Consider $H_k = \tau_x^{(k)} - \tau_x^{(k-1)}$, $k \in \mathbb{N}$
- From IID block structure, we know that $\{H_k\}_{k \in \mathbb{N}}$ is an IID process
- **Claim:**

For each $n \in \mathbb{N}$, the random variable $N_x(n) + 1$ is a stopping time w.r.t. $\{H_k\}_{k \in \mathbb{N}}$

$$\{N_x(n) + 1 = m\} = \{N_x(n) = m - 1\} = \left\{ \sum_{k=1}^{m-1} H_k \leq n < \sum_{k=1}^m H_k \right\}$$

$$N_x(n) \leq n \implies \mathbb{P}(N_x(n) < +\infty) = 1, \quad \mathbb{E}[N_x(n)] \leq n < +\infty$$

Proof of Lemma – Lower Bound

- Observe that

$$\sum_{k=1}^{N_x(n)+1} H_k > n$$

Proof of Lemma – Lower Bound

- Observe that

$$\sum_{k=1}^{N_x(n)+1} H_k > n$$

- Applying $\mathbb{E}[\cdot]$ on both sides, and using Wald's lemma, we get

$$\mathbb{E}[N_x(n) + 1] \cdot \mathbb{E}[H_1] > n$$

Proof of Lemma – Lower Bound

- Observe that

$$\sum_{k=1}^{N_x(n)+1} H_k > n$$

- Applying $\mathbb{E}[\cdot]$ on both sides, and using Wald's lemma, we get

$$\mathbb{E}[N_x(n) + 1] \cdot \mathbb{E}[H_1] > n$$

- Observe that $\mathbb{E}[H_1] = \mu_{xx}$.

Proof of Lemma – Lower Bound

- Observe that

$$\sum_{k=1}^{N_x(n)+1} H_k > n$$

- Applying $\mathbb{E}[\cdot]$ on both sides, and using Wald's lemma, we get

$$\mathbb{E}[N_x(n) + 1] \cdot \mathbb{E}[H_1] > n$$

- Observe that $\mathbb{E}[H_1] = \mu_{xx}$.
- Therefore,

$$\frac{\mathbb{E}[N_x(n)]}{n} > \frac{1}{\mu_{xx}} - \frac{1}{n}$$

Proof of Lemma – Lower Bound

- Observe that

$$\sum_{k=1}^{N_x(n)+1} H_k > n$$

- Applying $\mathbb{E}[\cdot]$ on both sides, and using Wald's lemma, we get

$$\mathbb{E}[N_x(n) + 1] \cdot \mathbb{E}[H_1] > n$$

- Observe that $\mathbb{E}[H_1] = \mu_{xx}$.
- Therefore,

$$\frac{\mathbb{E}[N_x(n)]}{n} > \frac{1}{\mu_{xx}} - \frac{1}{n}$$

- Taking limits as $n \rightarrow \infty$ on either sides, we get the desired lower bound

Proof of Lemma – Upper Bound

- Let $X_0 = x$. Fix a positive integer $M \in \mathbb{N}$.

Proof of Lemma – Upper Bound

- Let $X_0 = x$. Fix a positive integer $M \in \mathbb{N}$.
- For each $k \in \mathbb{N}$, let

$$\bar{H}_k := \min\{H_k, M\}$$

Proof of Lemma – Upper Bound

- Let $X_0 = x$. Fix a positive integer $M \in \mathbb{N}$.
- For each $k \in \mathbb{N}$, let

$$\bar{H}_k := \min\{H_k, M\}$$

- $\{\bar{H}_k\}_{k \in \mathbb{N}}$ is an IID sequence, $\bar{H}_k \leq H_k$, $\bar{H}_k \leq M$ for all k

Proof of Lemma – Upper Bound

- Let $X_0 = x$. Fix a positive integer $M \in \mathbb{N}$.
- For each $k \in \mathbb{N}$, let

$$\bar{H}_k := \min\{H_k, M\}$$

- $\{\bar{H}_k\}_{k \in \mathbb{N}}$ is an IID sequence, $\bar{H}_k \leq H_k$, $\bar{H}_k \leq M$ for all k
- Let

$$\bar{\mu}_{xx} := \mathbb{E}[\bar{H}_1].$$

Proof of Lemma – Upper Bound

- Let $X_0 = x$. Fix a positive integer $M \in \mathbb{N}$.
- For each $k \in \mathbb{N}$, let

$$\bar{H}_k := \min\{H_k, M\}$$

- $\{\bar{H}_k\}_{k \in \mathbb{N}}$ is an IID sequence, $\bar{H}_k \leq H_k$, $\bar{H}_k \leq M$ for all k
- Let

$$\bar{\mu}_{xx} := \mathbb{E}[\bar{H}_1].$$

- Then, $\bar{\mu}_{xx} \leq \mu_{xx}$

Proof of Lemma – Upper Bound

- Let $\bar{\tau}_x^{(0)} = 0$, and let

$$\bar{\tau}_x^{(k)} = \sum_{i=1}^k \bar{H}_i, \quad k \in \mathbb{N}.$$

Clearly, $\bar{\tau}_x^{(k)} \leq \tau_x^{(k)}$

Proof of Lemma – Upper Bound

- Let $\bar{\tau}_x^{(0)} = 0$, and let

$$\bar{\tau}_x^{(k)} = \sum_{i=1}^k \bar{H}_i, \quad k \in \mathbb{N}.$$

Clearly, $\bar{\tau}_x^{(k)} \leq \tau_x^{(k)}$

- Let

$$\bar{N}_x(n) = \sum_{k \in \mathbb{N}} \mathbf{1}_{\{\bar{\tau}_x^{(k)} \leq n\}}, \quad n \in \mathbb{N}$$

Proof of Lemma – Upper Bound

- Let $\bar{\tau}_x^{(0)} = 0$, and let

$$\bar{\tau}_x^{(k)} = \sum_{i=1}^k \bar{H}_i, \quad k \in \mathbb{N}.$$

Clearly, $\bar{\tau}_x^{(k)} \leq \tau_x^{(k)}$

- Let

$$\bar{N}_x(n) = \sum_{k \in \mathbb{N}} \mathbf{1}_{\{\bar{\tau}_x^{(k)} \leq n\}}, \quad n \in \mathbb{N}$$

- Observe that $\bar{N}_x(n) \geq N_x(n)$

Proof of Lemma – Upper Bound

- Let $\bar{\tau}_x^{(0)} = 0$, and let

$$\bar{\tau}_x^{(k)} = \sum_{i=1}^k \bar{H}_i, \quad k \in \mathbb{N}.$$

Clearly, $\bar{\tau}_x^{(k)} \leq \tau_x^{(k)}$

- Let

$$\bar{N}_x(n) = \sum_{k \in \mathbb{N}} \mathbf{1}_{\{\bar{\tau}_x^{(k)} \leq n\}}, \quad n \in \mathbb{N}$$

- Observe that $\bar{N}_x(n) \geq N_x(n)$

- **Claim:**

$\bar{N}_x(n) + 1$ is a stopping time w.r.t. $\{\bar{H}_k\}_{k \in \mathbb{N}}$

Proof of Lemma – Upper Bound

- We then have

$$\sum_{k=1}^{\bar{N}_x(n)+1} \bar{H}_k$$

Proof of Lemma – Upper Bound

- We then have

$$\sum_{k=1}^{\bar{N}_x(n)+1} \bar{H}_k$$

Proof of Lemma – Upper Bound

- We then have

$$\sum_{k=1}^{\bar{N}_x(n)+1} \bar{H}_k = \bar{\tau}_x^{(\bar{N}_x(n)+1)}$$

Proof of Lemma – Upper Bound

- We then have

$$\sum_{k=1}^{\bar{N}_x(n)+1} \bar{H}_k = \bar{\tau}_x^{(\bar{N}_x(n)+1)} = \bar{\tau}_x^{(\bar{N}_x(n))} + \bar{H}_{\bar{N}_x(n)+1}$$

Proof of Lemma – Upper Bound

- We then have

$$\begin{aligned}\sum_{k=1}^{\bar{N}_x(n)+1} \bar{H}_k &= \bar{\tau}_x^{(\bar{N}_x(n)+1)} = \bar{\tau}_x^{(\bar{N}_x(n))} + \bar{H}_{\bar{N}_x(n)+1} \\ &\leq n + M\end{aligned}$$

- Applying $\mathbb{E}[\cdot]$ and using Wald's lemma,

$$\mathbb{E}[\bar{N}_x(n) + 1] \cdot \bar{\mu}_{xx} \leq n + M.$$

Proof of Lemma – Upper Bound

- We then have

$$\begin{aligned}\sum_{k=1}^{\bar{N}_x(n)+1} \bar{H}_k &= \bar{\tau}_x^{(\bar{N}_x(n)+1)} = \bar{\tau}_x^{(\bar{N}_x(n))} + \bar{H}_{\bar{N}_x(n)+1} \\ &\leq n + M\end{aligned}$$

- Applying $\mathbb{E}[\cdot]$ and using Wald's lemma,

$$\mathbb{E}[\bar{N}_x(n) + 1] \cdot \bar{\mu}_{xx} \leq n + M.$$

- Noting that $N_x(n) \leq \bar{N}_x(n)$, we have

$$\mathbb{E}[N_x(n) + 1] \cdot \bar{\mu}_{xx} \leq \mathbb{E}[\bar{N}_x(n) + 1] \cdot \bar{\mu}_{xx} \leq n + M.$$

Proof of Lemma – Upper Bound

- Dividing both sides by n and taking limits as $n \rightarrow \infty$, we get

$$\lim_{n \rightarrow \infty} \frac{\mathbb{E}[N_x(n)]}{n} \leq \frac{1}{\bar{\mu}_{xx}} = \frac{1}{\mathbb{E}[\bar{H}_1]} = \frac{1}{\mathbb{E}[\min\{H_1, M\}]}$$

- Left-hand side does not depend on M , and above equation holds for all $M \in \mathbb{N}$
- Taking $M \rightarrow \infty$ on the right hand side, we get the desired upper bound

Recap – Invariant Distributions

Proposition (On Existence and Uniqueness of Invariant Distribution)

Let $\{X_n\}_{n=0}^{\infty}$ be an **irreducible**, time-homogeneous DTMC on a discrete state space \mathcal{X} with TPM P .

Then, **a unique stationary distribution π exists if and only if P is positive recurrent.**

In this case, $\pi_x = \frac{1}{\mu_{xx}} > 0$ for all $x \in \mathcal{X}$.

Example

- Consider a time-homogeneous DTMC with $\mathcal{X} = \{0, 1, 2, \dots\}$ whose transition probabilities are given by

$$P_{0,i} = \left(\frac{1}{2}\right)^i, \quad P_{i,i+1} = \frac{1}{2} = P_{i,0}$$

for all $i \in \mathbb{N}$.

1. Draw the transition graph.
2. Is the Markov chain irreducible?
3. Classify the states as transient, positive recurrent, or null recurrent.
4. Compute μ_{ii} for all $i \in \{0, 1, 2, \dots\}$.

Example

- A fair coin is tossed repeatedly and independently until the pattern “HTH” is observed for the first time.
How many tosses will be required on the average?

Example

- **[Goldmann Sachs Interview Question @IITSc] [2017-18 Placement Season]**

Imagine that you are playing a game that involves tossing a coin of bias 0.9 repeatedly and independently, with 3 lives at the start of the game. At each round, if the coin toss results in a tail, you lose one life. However, if the coin toss results in a head, you gain back your lost lives one at a time. If you have all 3 lives and the coin toss results in a head, nothing changes, and you continue tossing. You are allowed to play until you lose all your lives. What is the expected number of times you will play this game?