



Stochastic Processes

Sampling Techniques: Inverse Transform Technique, Rejection
Sampling

Karthik P. N.

Assistant Professor, Department of AI

Email: pnkarthik@ai.iith.ac.in

15 April 2025

Sampling from a Given Distribution

Objective

Given a cumulative distribution function (CDF) $F : \mathbb{R} \rightarrow [0, 1]$, generate a sample $X \sim F$.

Sampling from a Given Distribution

Objective

Given a cumulative distribution function (CDF) $F : \mathbb{R} \rightarrow [0, 1]$, generate a sample $X \sim F$.

Remarks:

- The CDF F may or may not be continuous

Sampling from a Given Distribution

Objective

Given a cumulative distribution function (CDF) $F : \mathbb{R} \rightarrow [0, 1]$, generate a sample $X \sim F$.

Remarks:

- The CDF F may or may not be continuous
- If F is continuous and differentiable, it admits a probability density function f

Sampling from a Given Distribution

Objective

Given a cumulative distribution function (CDF) $F : \mathbb{R} \rightarrow [0, 1]$, generate a sample $X \sim F$.

Remarks:

- The CDF F may or may not be continuous
- If F is continuous and differentiable, it admits a probability density function f
- Instead of the CDF F , we may be given a target PMF or PDF from which to sample

Sampling from a Given Distribution

Objective

Given a cumulative distribution function (CDF) $F : \mathbb{R} \rightarrow [0, 1]$, generate a sample $X \sim F$.

Remarks:

- The CDF F may or may not be continuous
- If F is continuous and differentiable, it admits a probability density function f
- Instead of the CDF F , we may be given a target PMF or PDF from which to sample

Sampling from a Given Distribution

Objective

Given a cumulative distribution function (CDF) $F : \mathbb{R} \rightarrow [0, 1]$, generate a sample $X \sim F$.

Remarks:

- The CDF F may or may not be continuous
- If F is continuous and differentiable, it admits a probability density function f
- Instead of the CDF F , we may be given a target PMF or PDF from which to sample

The basic ingredient: **Uniform $(0, 1)$** sample(s) via PRNGs

Inverse Transform Technique (ITT)

Inverse Transform Technique

Objective

Given a cumulative distribution function (CDF) $F : \mathbb{R} \rightarrow [0, 1]$, generate a sample $X \sim F$.

Inverse Transform Technique

Objective

Given a cumulative distribution function (CDF) $F : \mathbb{R} \rightarrow [0, 1]$, generate a sample $X \sim F$.

- Sample $U \sim \text{Unif}(0, 1)$

Inverse Transform Technique

Objective

Given a cumulative distribution function (CDF) $F : \mathbb{R} \rightarrow [0, 1]$, generate a sample $X \sim F$.

- Sample $U \sim \text{Unif}(0, 1)$
- Define the function F^{-1} as

$$F^{-1}(u) := \min\{x \in \mathbb{R} : F(x) \geq u\}$$

Inverse Transform Technique

Objective

Given a cumulative distribution function (CDF) $F : \mathbb{R} \rightarrow [0, 1]$, generate a sample $X \sim F$.

- Sample $U \sim \text{Unif}(0, 1)$
- Define the function F^{-1} as

$$F^{-1}(u) := \min\{x \in \mathbb{R} : F(x) \geq u\}$$

- F^{-1} is called the **quantile** function

Inverse Transform Technique

Objective

Given a cumulative distribution function (CDF) $F : \mathbb{R} \rightarrow [0, 1]$, generate a sample $X \sim F$.

- Sample $U \sim \text{Unif}(0, 1)$
- Define the function F^{-1} as

$$F^{-1}(u) := \min\{x \in \mathbb{R} : F(x) \geq u\}$$

- F^{-1} is called the **quantile** function
- Set $X = F^{-1}(U)$

Inverse Transform Technique

Objective

Given a cumulative distribution function (CDF) $F : \mathbb{R} \rightarrow [0, 1]$, generate a sample $X \sim F$.

- Sample $U \sim \text{Unif}(0, 1)$
- Define the function F^{-1} as

$$F^{-1}(u) := \min\{x \in \mathbb{R} : F(x) \geq u\}$$

- F^{-1} is called the **quantile** function
- Set $X = F^{-1}(U)$
- **Claim:** The CDF of X is exactly equal to F , i.e., $F_X = F$

Example

- Let X be a discrete random variable with the following PMF:

$$p_X(x) = \begin{cases} 0.1, & x = 10, \\ 0.2, & x = 20, \\ 0.3, & x = 30, \\ 0.4, & x = 40, \\ 0, & \text{otherwise.} \end{cases}$$

Use the inverse transform method to generate a sample from the above distribution.

Example

- **[Generating a Sample from Rayleigh Distribution]**

The PDF of the Rayleigh distribution is given by

$$f(r) = r e^{-r^2/2}, \quad r > 0.$$

Use the inverse transform method to generate a sample from the above distribution.

Gaussian Samples on Python via ITT

- Python's built-in module

```
numpy.random.normal(loc, scale, size)
```

generates n independent samples from $\mathcal{N}(\mu, \sigma^2)$, where

$$n = \text{size}, \quad \mu = \text{loc}, \quad \sigma = \text{scale}.$$

- In principle, the above module uses the inverse transform technique

Gaussian Samples on Python via ITT

1. Let $U_1, U_2 \stackrel{\text{i.i.d.}}{\sim} \text{Unif}(0, 1)$
2. Let R and Θ be two random variables defined via

$$R = F_1^{-1}(U_1), \quad \Theta = 2\pi U_2,$$

where F_1 is the CDF of the Rayleigh distribution

3. Let Y_1 and Y_2 be defined as

$$Y_1 = R \cos(\Theta), \quad Y_2 = R \sin(\Theta).$$

4. Then, $Y_1, Y_2 \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, 1)$
5. To get $X \sim \mathcal{N}(\mu, \sigma^2)$, simply **discard Y_2 , and**

$$X = \sigma Y_1 + \mu.$$

6. Repeat steps 1-5 a total of n times to get $X_1, X_2, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(\mu, \sigma^2)$

Rejection Sampling (Accept-Reject) Technique



Rejection Sampling (Accept-Reject) Technique

- Many a times, computing F^{-1} is tedious

Rejection Sampling (Accept-Reject) Technique

- Many a times, computing F^{-1} is tedious
- For instance, consider the PDF

$$f(x) = 6x(1 - x), \quad x \in [0, 1]$$

Rejection Sampling (Accept-Reject) Technique

- Many a times, computing F^{-1} is tedious
- For instance, consider the PDF

$$f(x) = 6x(1 - x), \quad x \in [0, 1]$$

- The CDF corresponding to the above PDF is given by

$$F(x) = \begin{cases} 0, & x < 0, \\ 3x^2 - 2x^3, & 0 \leq x < 1, \\ 1, & x \geq 1. \end{cases}$$

Rejection Sampling (Accept-Reject) Technique

- Many a times, computing F^{-1} is tedious
- For instance, consider the PDF

$$f(x) = 6x(1 - x), \quad x \in [0, 1]$$

- The CDF corresponding to the above PDF is given by

$$F(x) = \begin{cases} 0, & x < 0, \\ 3x^2 - 2x^3, & 0 \leq x < 1, \\ 1, & x \geq 1. \end{cases}$$

- Computation of F^{-1} is non-trivial; need to use numerical techniques

Rejection Sampling (Accept-Reject) Technique

- Many a times, computing F^{-1} is tedious
- For instance, consider the PDF

$$f(x) = 6x(1 - x), \quad x \in [0, 1]$$

- The CDF corresponding to the above PDF is given by

$$F(x) = \begin{cases} 0, & x < 0, \\ 3x^2 - 2x^3, & 0 \leq x < 1, \\ 1, & x \geq 1. \end{cases}$$

- Computation of F^{-1} is non-trivial; need to use numerical techniques
- **Alternative solution:** Rejection sampling!

Rejection Sampling (Accept-Reject) Technique

Objective

Given a PDF f with CDF F for which computing F^{-1} is tedious, generate a sample $X \sim f$.

Rejection Sampling (Accept-Reject) Technique

Objective

Given a PDF f with CDF F for which computing F^{-1} is tedious, generate a sample $X \sim f$.

- Let U, Z be a pair of continuous random variables satisfying the following properties:

Rejection Sampling (Accept-Reject) Technique

Objective

Given a PDF f with CDF F for which computing F^{-1} is tedious, generate a sample $X \sim f$.

- Let U, Z be a **pair of continuous random variables** satisfying the following properties:
 1. $U \sim \text{Unif}(0, 1)$.

Rejection Sampling (Accept-Reject) Technique

Objective

Given a PDF f with CDF F for which computing F^{-1} is tedious, generate a sample $X \sim f$.

- Let U, Z be a **pair of continuous random variables** satisfying the following properties:
 1. $U \sim \text{Unif}(0, 1)$.
 2. U is independent of Z .

Rejection Sampling (Accept-Reject) Technique

Objective

Given a PDF f with CDF F for which computing F^{-1} is tedious, generate a sample $X \sim f$.

- Let U, Z be a **pair of continuous random variables** satisfying the following properties:
 - $U \sim \text{Unif}(0, 1)$.
 - U is independent of Z .
 - There exists a real number $a \geq 1$ such that

$$f(x) \leq a f_Z(x) \quad \forall x \in \mathbb{R},$$

where f_Z is the PDF of Z

Rejection Sampling (Accept-Reject) Technique

Objective

Given a PDF f with CDF F for which computing F^{-1} is tedious, generate a sample $X \sim f$.

- Let U, Z be a **pair of continuous random variables** satisfying the following properties:
 - $U \sim \text{Unif}(0, 1)$.
 - U is independent of Z .
 - There exists a real number $a \geq 1$ such that

$$f(x) \leq a f_Z(x) \quad \forall x \in \mathbb{R},$$

where f_Z is the PDF of Z

- Generate (U, Z) satisfying 1-3 above

Rejection Sampling (Accept-Reject) Technique

Objective

Given a PDF f with CDF F for which computing F^{-1} is tedious, generate a sample $X \sim f$.

- Let U, Z be a **pair of continuous random variables** satisfying the following properties:
 - $U \sim \text{Unif}(0, 1)$.
 - U is independent of Z .
 - There exists a real number $a \geq 1$ such that

$$f(x) \leq a f_Z(x) \quad \forall x \in \mathbb{R},$$

where f_Z is the PDF of Z

- Generate (U, Z) satisfying 1-3 above
 - If $a U f_Z(U) \leq f(U)$, **accept** (U, Z) , and set $X = Z$.

Rejection Sampling (Accept-Reject) Technique

Objective

Given a PDF f with CDF F for which computing F^{-1} is tedious, generate a sample $X \sim f$.

- Let U, Z be a **pair of continuous random variables** satisfying the following properties:
 - $U \sim \text{Unif}(0, 1)$.
 - U is independent of Z .
 - There exists a real number $a \geq 1$ such that

$$f(x) \leq a f_Z(x) \quad \forall x \in \mathbb{R},$$

where f_Z is the PDF of Z

- Generate (U, Z) satisfying 1-3 above
 - If $a U f_Z(U) \leq f(U)$, **accept** (U, Z) , and set $X = Z$.
 - If not, **reject** current (U, Z) and generate new (U, Z) . Repeat till (i) holds.

Rejection Sampling (Accept-Reject) Technique

- Let U, Z be a **pair of continuous random variables** satisfying the following properties:
 - $U \sim \text{Unif}(0, 1)$.
 - U is independent of Z .
 - There exists a real number $a \geq 1$ such that

$$f(x) \leq a f_Z(x) \quad \forall x \in \mathbb{R},$$

where f_Z is the PDF of Z

- Generate (U, Z) satisfying 1-3 above
 - If $a U f_Z(Z) \leq f(Z)$, **accept** (U, Z) , and set $X = Z$.
 - If not, **reject** current (U, Z) and generate new (U, Z) . Repeat till (i) holds.

Rejection Sampling (Accept-Reject) Technique

- Let U, Z be a **pair of continuous random variables** satisfying the following properties:
 - $U \sim \text{Unif}(0, 1)$.
 - U is independent of Z .
 - There exists a real number $a \geq 1$ such that

$$f(x) \leq a f_Z(x) \quad \forall x \in \mathbb{R},$$

where f_Z is the PDF of Z

- Generate (U, Z) satisfying 1-3 above
 - If $a U f_Z(Z) \leq f(Z)$, **accept** (U, Z) , and set $X = Z$.
 - If not, **reject** current (U, Z) and generate new (U, Z) . Repeat till (i) holds.

Theorem (Rejection Sampling)

Let $E = \{a U f_Z(Z) \leq f(Z)\}$. Then,

$$F_X(x) = \mathbb{P}(X \leq x) = \mathbb{P}(Z \leq x \mid E).$$

Example

- Design an algorithm to generate a sample $X \sim f$, where

$$f(x) = 6x(1 - x), \quad x \in [0, 1].$$

Example

- For fixed constants $\lambda, t > 0$, the $\text{Gamma}(\lambda, t)$ PDF is given by

$$f(x) = e^{-\lambda x} \frac{\lambda^t x^{t-1}}{\Gamma(t)}, \quad x > 0.$$

1. When $t \in \mathbb{N}$, suggest a technique to generate a sample $X \sim f$ via ITT.
2. When $t \notin \mathbb{N}$, design an algorithm to sample $X \sim f$ via rejection sampling.

Hint: Take $Z \sim \text{Exponential}(1/t)$.