



# Stochastic Processes

Stopping Times, Wald's Lemma, Strong Independence Property,  
Properties of Stopping Times

**Karthik P. N.**

**Assistant Professor, Department of AI**

**Email: [pnkarthik@ai.iith.ac.in](mailto:pnkarthik@ai.iith.ac.in)**

28 February 2025



## Filtrations

Fix a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ .

Let  $\mathcal{T}$  be an **ordered** index set.

## Filtrations

Fix a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ .

Let  $\mathcal{T}$  be an **ordered** index set.

### Definition (Filtration)

Consider a collection of  $\sigma$ -algebras  $\mathcal{G}_\bullet = \{\mathcal{G}_t : t \in \mathcal{T}\}$  such that  $\mathcal{G}_t \subseteq \mathcal{F}$  for all  $t$ .  
The above collection is called a **filtration** if

$$\mathcal{G}_s \subseteq \mathcal{G}_t \quad \forall s \leq t.$$

## Filtrations

Fix a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ .

Let  $\mathcal{T}$  be an **ordered** index set.

### Definition (Filtration)

Consider a collection of  $\sigma$ -algebras  $\mathcal{G}_\bullet = \{\mathcal{G}_t : t \in \mathcal{T}\}$  such that  $\mathcal{G}_t \subseteq \mathcal{F}$  for all  $t$ .  
The above collection is called a **filtration** if

$$\mathcal{G}_s \subseteq \mathcal{G}_t \quad \forall s \leq t.$$

Example:

Let  $\{X_t : t \in \mathcal{T}\}$  be a stochastic process defined w.r.t.  $\mathcal{F}$ . Then,

$$\mathcal{G}_t = \sigma(X_s : s \leq t)$$

is called the **natural filtration** associated with the process  $\{X_t : t \in \mathcal{T}\}$ .

## Stopping Time

Fix a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ .

Let  $\mathcal{T}$  be an **ordered** index set.

Fix a filtration  $\mathcal{G}_\bullet = \{\mathcal{G}_t : t \in \mathcal{T}\}$ .

### Definition (Stopping Time)

A random variable  $\tau$  is called a **stopping time w.r.t. the filtration  $\mathcal{G}_\bullet$**  if:

- $\mathbb{P}(\tau < +\infty) = 1$ .
- For each  $t \in \mathcal{T}$ ,

$$\{\tau \leq t\} \in \mathcal{G}_t.$$

## Stopping Time

Fix a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ .

Let  $\mathcal{T}$  be an **ordered** index set.

Fix a filtration  $\mathcal{G}_\bullet = \{\mathcal{G}_t : t \in \mathcal{T}\}$ .

### Definition (Stopping Time)

A random variable  $\tau$  is called a **stopping time w.r.t. the filtration  $\mathcal{G}_\bullet$**  if:

- $\mathbb{P}(\tau < +\infty) = 1$ .
- For each  $t \in \mathcal{T}$ ,

$$\{\tau \leq t\} \in \mathcal{G}_t.$$

If  $\mathcal{G}_t = \sigma(X_s : s \leq t)$ , then the question “is  $\tau \leq t$ ?” can be answered by simply looking at the process up to time  $t$ .

## Stopping Time w.r.t. a Process

Fix a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ .

Let  $\mathcal{T}$  be an **ordered** index set.

Let  $\{X_t : t \in \mathcal{T}\}$  be a process w.r.t.  $\mathcal{F}$ .

### Definition (Stopping Time w.r.t. a Process)

A random variable  $\tau$  is called a **stopping time w.r.t. the process  $\{X_t : t \in \mathcal{T}\}$**  if:

- $\mathbb{P}(\tau < +\infty) = 1$ .
- For each  $t \in \mathcal{T}$ ,

$$\{\tau \leq t\} \in \sigma(X_s : s \leq t).$$

That is, the question “is  $\tau \leq t$ ?” can be answered by simply looking at the process up to time  $t$ .

## Examples

- Let  $\{X_n : n \in \mathbb{N}\}$  be a process.

Fix a set  $A \subseteq \mathbb{R}$ .

Let  $\tau_X^A$  be defined as

$$\tau_X^A := \inf\{n \in \mathbb{N} : X_n \in A\}.$$

Is  $\tau_X^A$  a stopping time w.r.t. the process  $\{X_n : n \in \mathbb{N}\}$ ?





## Discrete Stopping Times

Fix a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ .

Let  $\{\mathcal{G}_n : n \in \mathbb{N}\}$  be a filtration.

## Discrete Stopping Times

Fix a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ .

Let  $\{\mathcal{G}_n : n \in \mathbb{N}\}$  be a filtration.

### Lemma (Discrete Stopping Times)

A **discrete** random variable  $\tau$  is a stopping time w.r.t.  $\{\mathcal{G}_n\}_{n=1}^{\infty}$  if and only if

$$\mathbb{P}(\tau < +\infty) = 1, \quad \{\tau = n\} \in \mathcal{G}_n \quad \forall n \in \mathbb{N}.$$



$$\mathbb{P}(\tau < +\infty) = 1, \quad \{\tau \leq n\} \in \mathcal{G}_n \quad \forall n \in \mathbb{N}.$$

## Proof of Lemma

Assume  $\mathbb{P}(\tau < +\infty) = 1$ .

- Suppose that  $\{\tau = k\} \in \mathcal{G}_k$  for all  $k \in \mathbb{N}$

## Proof of Lemma

Assume  $\mathbb{P}(\tau < +\infty) = 1$ .

- Suppose that  $\{\tau = k\} \in \mathcal{G}_k$  for all  $k \in \mathbb{N}$ 
  - We need to prove that  $\{\tau \leq n\} \in \mathcal{G}_n$  for all  $n \in \mathbb{N}$

## Proof of Lemma

Assume  $\mathbb{P}(\tau < +\infty) = 1$ .

- Suppose that  $\{\tau = k\} \in \mathcal{G}_k$  for all  $k \in \mathbb{N}$ 
  - We need to prove that  $\{\tau \leq n\} \in \mathcal{G}_n$  for all  $n \in \mathbb{N}$
  - Observe that for any  $n \in \mathbb{N}$ ,

$$\{\tau \leq n\} = \bigcup_{k=1}^n \{\tau = k\}$$

## Proof of Lemma

Assume  $\mathbb{P}(\tau < +\infty) = 1$ .

- Suppose that  $\{\tau = k\} \in \mathcal{G}_k$  for all  $k \in \mathbb{N}$ 
  - We need to prove that  $\{\tau \leq n\} \in \mathcal{G}_n$  for all  $n \in \mathbb{N}$
  - Observe that for any  $n \in \mathbb{N}$ ,

$$\{\tau \leq n\} = \bigcup_{k=1}^n \{\tau = k\}$$

- $\{\tau = k\} \in \mathcal{G}_k \subset \mathcal{G}_n \implies \{\tau = k\} \in \mathcal{G}_n$  for all  $k \in \{1, \dots, n\}$

## Proof of Lemma

Assume  $\mathbb{P}(\tau < +\infty) = 1$ .

- Suppose that  $\{\tau = k\} \in \mathcal{G}_k$  for all  $k \in \mathbb{N}$ 
  - We need to prove that  $\{\tau \leq n\} \in \mathcal{G}_n$  for all  $n \in \mathbb{N}$
  - Observe that for any  $n \in \mathbb{N}$ ,

$$\{\tau \leq n\} = \bigcup_{k=1}^n \{\tau = k\}$$

- $\{\tau = k\} \in \mathcal{G}_k \subset \mathcal{G}_n \implies \{\tau = k\} \in \mathcal{G}_n$  for all  $k \in \{1, \dots, n\}$
- Suppose that  $\{\tau \leq k\} \in \mathcal{G}_k$  for all  $k \in \mathbb{N}$

## Proof of Lemma

Assume  $\mathbb{P}(\tau < +\infty) = 1$ .

- Suppose that  $\{\tau = k\} \in \mathcal{G}_k$  for all  $k \in \mathbb{N}$ 
  - We need to prove that  $\{\tau \leq n\} \in \mathcal{G}_n$  for all  $n \in \mathbb{N}$
  - Observe that for any  $n \in \mathbb{N}$ ,

$$\{\tau \leq n\} = \bigcup_{k=1}^n \{\tau = k\}$$

- $\{\tau = k\} \in \mathcal{G}_k \subset \mathcal{G}_n \implies \{\tau = k\} \in \mathcal{G}_n$  for all  $k \in \{1, \dots, n\}$
- Suppose that  $\{\tau \leq k\} \in \mathcal{G}_k$  for all  $k \in \mathbb{N}$ 
  - We need to prove that  $\{\tau = n\} \in \mathcal{G}_n$  for every  $n \in \mathbb{N}$



## Proof of Lemma

Assume  $\mathbb{P}(\tau < +\infty) = 1$ .

- Suppose that  $\{\tau = k\} \in \mathcal{G}_k$  for all  $k \in \mathbb{N}$ 
  - We need to prove that  $\{\tau \leq n\} \in \mathcal{G}_n$  for all  $n \in \mathbb{N}$
  - Observe that for any  $n \in \mathbb{N}$ ,

$$\{\tau \leq n\} = \bigcup_{k=1}^n \{\tau = k\}$$

- $\{\tau = k\} \in \mathcal{G}_k \subset \mathcal{G}_n \implies \{\tau = k\} \in \mathcal{G}_n$  for all  $k \in \{1, \dots, n\}$
- Suppose that  $\{\tau \leq k\} \in \mathcal{G}_k$  for all  $k \in \mathbb{N}$ 
  - We need to prove that  $\{\tau = n\} \in \mathcal{G}_n$  for every  $n \in \mathbb{N}$
  - Observe that

$$\{\tau = n\} = \{\tau \leq n\} \setminus \{\tau \leq n-1\}$$

## Proof of Lemma

Assume  $\mathbb{P}(\tau < +\infty) = 1$ .

- Suppose that  $\{\tau = k\} \in \mathcal{G}_k$  for all  $k \in \mathbb{N}$ 
  - We need to prove that  $\{\tau \leq n\} \in \mathcal{G}_n$  for all  $n \in \mathbb{N}$
  - Observe that for any  $n \in \mathbb{N}$ ,

$$\{\tau \leq n\} = \bigcup_{k=1}^n \{\tau = k\}$$

- $\{\tau = k\} \in \mathcal{G}_k \subset \mathcal{G}_n \implies \{\tau = k\} \in \mathcal{G}_n$  for all  $k \in \{1, \dots, n\}$
- Suppose that  $\{\tau \leq k\} \in \mathcal{G}_k$  for all  $k \in \mathbb{N}$ 
  - We need to prove that  $\{\tau = n\} \in \mathcal{G}_n$  for every  $n \in \mathbb{N}$
  - Observe that

$$\{\tau = n\} = \{\tau \leq n\} \setminus \{\tau \leq n-1\}$$

- $\{\tau = n-1\} \in \mathcal{G}_{n-1} \subset \mathcal{G}_n$

## Example

- Let  $\{X_n\}_{n=1}^\infty$  be an  $\mathbb{N}$ -valued process.  
Fix  $y \in \mathbb{N}$ .  
Let  $\tau_y^{(0)} := 0$ , and

$$\tau_y^{(k)} = \inf\{n > \tau_y^{(k-1)} : X_n = y\}, \quad k \in \mathbb{N}.$$

For each  $k \in \mathbb{N}$ , prove that  $\tau_y^{(k)}$  is a stopping time w.r.t. the process  $\{X_n\}_{n=1}^\infty$ .



## Wald's Lemma

Fix a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ .

## Wald's Lemma

Fix a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ .

### Lemma (Wald's Lemma [Wal44, Wal45])

Let  $\{X_n\}_{n=1}^{\infty}$  be an **IID** process w.r.t.  $\mathcal{F}$ , with  $\mathbb{E}|X_1| < +\infty$ .

For each  $n \in \mathbb{N}$ , let

$$S_n = \sum_{i=1}^n X_i.$$

If  $\tau$  is a **stopping time** w.r.t. the process  $\{X_n\}_{n=1}^{\infty}$ , with  $\mathbb{E}|\tau| < +\infty$ , then

$$\mathbb{E}[S_{\tau}] = \mathbb{E}\left[\sum_{i=1}^{\tau} X_i\right] = \mathbb{E}[\tau] \cdot \mathbb{E}[X_1].$$

## Example

- Suppose  $X_1, X_2, \dots \stackrel{\text{i.i.d.}}{\sim} \text{Geometric}(0.5)$ .  
For each  $n \in \mathbb{N}$ , let  $S_n = \sum_{i=1}^n X_i$ .

Let  $\tau$  be defined as

$$\tau := \inf \left\{ n \geq 1 : S_n = 33 \right\}.$$

Determine  $\mathbb{E}[\tau]$ .

## References



Abraham Wald.

On cumulative sums of random variables.

*The Annals of Mathematical Statistics*, 15(3):283–296, 1944.



Abraham Wald.

Some generalizations of the theory of cumulative sums of random variables.

*The Annals of Mathematical Statistics*, 16(3):287–293, 1945.