



Stochastic Processes

Markov Chain Monte Carlo Methods: Metropolis–Hastings
Algorithm, Examples

Karthik P. N.

Assistant Professor, Department of AI

Email: pnkarthik@ai.iith.ac.in

25 April 2025

Recap of MCMC Techniques

Objective

To evaluate a sum of the form

$$I = \sum_{x \in \mathcal{X}} f(x) \pi(x),$$

for given functions $f, \pi : \mathcal{X} \rightarrow \mathbb{R}$, where π is a probability distribution on \mathcal{X} .
Here, $\mathcal{X} \subseteq \mathbb{R}^d$ for some d .

Difficulties:

- Direct sampling from π (via ITT or rejection sampling methods) may be difficult
- High dimensionality: d may be large (e.g., 10000)
- f and/or π may have complex expressions, making it difficult to compute the sum

Metropolis–Hastings Algorithm

The Metropolis–Hastings MCMC Algorithm

- Originally proposed by Metropolis et al. [[MRR⁺53](#)] and Hastings [[Has70](#)]

The Metropolis–Hastings MCMC Algorithm

- Originally proposed by Metropolis et al. [[MRR⁺53](#)] and Hastings [[Has70](#)]
- The key idea: construct an ergodic Markov chain whose stationary distribution is π

The Metropolis–Hastings MCMC Algorithm

- Originally proposed by Metropolis et al. [[MRR⁺53](#)] and Hastings [[Has70](#)]
- The key idea: construct an ergodic Markov chain whose stationary distribution is π
- If $\{X_n\}_{n=0}^\infty$ is this Markov chain, it follows by the ergodic theorem that

$$\frac{1}{n} \sum_{k=1}^n f(X_k) \xrightarrow{\text{a.s.}} I.$$

The Metropolis–Hastings MCMC Algorithm

- Originally proposed by Metropolis et al. [[MRR⁺53](#)] and Hastings [[Has70](#)]
- The key idea: construct an ergodic Markov chain whose stationary distribution is π
- If $\{X_n\}_{n=0}^\infty$ is this Markov chain, it follows by the ergodic theorem that

$$\frac{1}{n} \sum_{k=1}^n f(X_k) \xrightarrow{\text{a.s.}} I.$$

- The method also applies when \mathcal{X} is countable or uncountable

Construction of Markov Chain

- Let Q be any row-stochastic matrix on $\mathcal{X} \times \mathcal{X}$, i.e.,

$$Q_{x,y} > 0, \quad \sum_y Q_{x,y} = 1 \quad \forall x, y \in \mathcal{X}.$$

Construction of Markov Chain

- Let Q be any row-stochastic matrix on $\mathcal{X} \times \mathcal{X}$, i.e.,

$$Q_{x,y} > 0, \quad \sum_y Q_{x,y} = 1 \quad \forall x, y \in \mathcal{X}.$$

- Let S be a non-negative, **symmetric** matrix having same size as Q

Construction of Markov Chain

- Let Q be any row-stochastic matrix on $\mathcal{X} \times \mathcal{X}$, i.e.,

$$Q_{x,y} > 0, \quad \sum_y Q_{x,y} = 1 \quad \forall x, y \in \mathcal{X}.$$

- Let S be a non-negative, **symmetric** matrix having same size as Q
- Let $A = [A_{x,y}]_{x,y \in \mathcal{X}}$ be defined via

$$A_{x,y} = \frac{S_{x,y}}{1 + \frac{\pi(x) Q_{x,y}}{\pi(y) Q_{y,x}}}.$$

Construction of Markov Chain

- Let Q be any row-stochastic matrix on $\mathcal{X} \times \mathcal{X}$, i.e.,

$$Q_{x,y} > 0, \quad \sum_y Q_{x,y} = 1 \quad \forall x, y \in \mathcal{X}.$$

- Let S be a non-negative, **symmetric** matrix having same size as Q
- Let $A = [A_{x,y}]_{x,y \in \mathcal{X}}$ be defined via

$$A_{x,y} = \frac{S_{x,y}}{1 + \frac{\pi(x) Q_{x,y}}{\pi(y) Q_{y,x}}}.$$

- The entries of S are chosen in such a way that $0 < A_{x,y} \leq 1$ for all $x, y \in \mathcal{X}$

Construction of Markov Chain

- Let Q be any row-stochastic matrix on $\mathcal{X} \times \mathcal{X}$, i.e.,

$$Q_{x,y} > 0, \quad \sum_y Q_{x,y} = 1 \quad \forall x, y \in \mathcal{X}.$$

- Let S be a non-negative, **symmetric** matrix having same size as Q
- Let $A = [A_{x,y}]_{x,y \in \mathcal{X}}$ be defined via

$$A_{x,y} = \frac{S_{x,y}}{1 + \frac{\pi(x) Q_{x,y}}{\pi(y) Q_{y,x}}}.$$

- The entries of S are chosen in such a way that $0 < A_{x,y} \leq 1$ for all $x, y \in \mathcal{X}$
- A is called the **acceptance matrix**

Construction of Markov Chain

- Choose an arbitrary starting state $X_0 = x$
- At each time step $n \in \mathbb{N}$, do the following two tasks:
 - Pick Y such that

$$\mathbb{P}(Y = y \mid X_{n-1} = x) = Q_{x,y}$$

- Define X_n as

$$X_n = \begin{cases} y, & \text{with probability } A_{x,y}, \\ X_{n-1}, & \text{with probability } 1 - A_{x,y}. \end{cases}$$

- **Claim:** The TPM of $\{X_n\}_{n=0}^{\infty}$ will have π as the unique stationary distribution!

Choice of S Matrix in Metropolis–Hastings Algorithm

- The matrix S used in [MRR⁺53] (call this $S^{(M)}$) is given by

$$S_{x,y}^{(M)} = \begin{cases} 1 + \frac{\pi_x Q_{x,y}}{\pi_y Q_{y,x}}, & \frac{\pi_y Q_{y,x}}{\pi_x Q_{x,y}} \geq 1, \\ 1 + \frac{\pi_y Q_{y,x}}{\pi_x Q_{x,y}}, & \frac{\pi_y Q_{y,x}}{\pi_x Q_{x,y}} < 1. \end{cases}$$

Choice of S Matrix in Metropolis–Hastings Algorithm

- The matrix S used in [MRR⁺53] (call this $S^{(M)}$) is given by

$$S_{x,y}^{(M)} = \begin{cases} 1 + \frac{\pi_x Q_{x,y}}{\pi_y Q_{y,x}}, & \frac{\pi_y Q_{y,x}}{\pi_x Q_{x,y}} \geq 1, \\ 1 + \frac{\pi_y Q_{y,x}}{\pi_x Q_{x,y}}, & \frac{\pi_y Q_{y,x}}{\pi_x Q_{x,y}} < 1. \end{cases}$$

Choice of S Matrix in Metropolis–Hastings Algorithm

- The matrix S used in [MRR⁺53] (call this $S^{(M)}$) is given by

$$S_{x,y}^{(M)} = \begin{cases} 1 + \frac{\pi_x Q_{x,y}}{\pi_y Q_{y,x}}, & \frac{\pi_y Q_{y,x}}{\pi_x Q_{x,y}} \geq 1, \\ 1 + \frac{\pi_y Q_{y,x}}{\pi_x Q_{x,y}}, & \frac{\pi_y Q_{y,x}}{\pi_x Q_{x,y}} < 1. \end{cases}$$

With $S = S^{(M)}$ as above, the acceptance matrix A becomes

$$A_{x,y} = \begin{cases} 1, & \frac{\pi_y Q_{y,x}}{\pi_x Q_{x,y}} \geq 1, \\ \frac{\pi_y Q_{y,x}}{\pi_x Q_{x,y}}, & \frac{\pi_y Q_{y,x}}{\pi_x Q_{x,y}} < 1 \end{cases} = \min \left\{ 1, \frac{\pi_y Q_{y,x}}{\pi_x Q_{x,y}} \right\}.$$

- Barker [Bar65] proposes using $S = S^{(B)}$, where $S_{x,y}^{(B)} = 1$ for all $x, y \in \mathcal{X}$

Choice of S Matrix in Metropolis–Hastings Algorithm

- If Q is symmetric (e.g., every row of Q is uniform), then

$$A_{x,y}^{(M)} = \min \left\{ 1, \frac{\pi_y}{\pi_x} \right\}, \quad A_{x,y}^{(B)} = \frac{\pi_y}{\pi_x + \pi_y}.$$




Choice of S Matrix in Metropolis–Hastings Algorithm

- If Q is symmetric (e.g., every row of Q is uniform), then

$$A_{x,y}^{(M)} = \min \left\{ 1, \frac{\pi_y}{\pi_x} \right\}, \quad A_{x,y}^{(B)} = \frac{\pi_y}{\pi_x + \pi_y}.$$

- Thus, if $\pi_x = \pi_y$, then $A_{x,y}^{(M)} = 1$, whereas $A_{x,y}^{(B)} = 0.5$
In this case, [MRR⁺53] suggests picking $X_{n+1} = y$ with probability 1
[Bar65] suggests taking $X_{n+1} = X_n = x$ with probability 0.5 and taking $X_{n+1} = y$ with probability 0.5

References

-  **Anthony Alfred Barker.**
Monte carlo calculations of the radial distribution functions for a proton? electron plasma.
Australian Journal of Physics, 18(2):119–134, 1965.
-  **W Keith Hastings.**
Monte carlo sampling methods using markov chains and their applications.
Biometrika, 1970.
-  **N Metropolis, A Rosenbluth, MN Rosenbluth, AH Teller, and E Teller.**
Perspective on “equation of state calculations by fast computing machines”.
J. Chem. Phys, 21:1087–1092, 1953.