



# Stochastic Processes

Communicating Classes, Class Properties, Irreducibility, Aperiodicity,  
Invariant Distribution

**Karthik P. N.**

**Assistant Professor, Department of AI**

**Email: [pnkarthik@ai.iith.ac.in](mailto:pnkarthik@ai.iith.ac.in)**

28 March 2025

## Definition (Reachability)

State  $y \in \mathcal{X}$  is said to be **reachable** from state  $x \in \mathcal{X}$  if there exists  $n \in \mathbb{N} \cup \{0\}$  such that the probability of reaching  $y$  in  $n$  steps starting from  $x$  is strictly positive.

Notation:  $x \longrightarrow y$ .

## Definition (Reachability)

State  $y \in \mathcal{X}$  is said to be **reachable** from state  $x \in \mathcal{X}$  if there exists  $n \in \mathbb{N} \cup \{0\}$  such that the probability of reaching  $y$  in  $n$  steps starting from  $x$  is strictly positive.

Notation:  $x \longrightarrow y$ .

Remark: For a time-homogeneous Markov chain with state space  $\mathcal{X}$  and TPM  $P$ ,

$$x \longrightarrow y \iff \exists n \in \mathbb{N} \cup \{0\} \text{ such that } P_{x,y}^n > 0.$$

# Communication

## Definition (Communication)

Two states  $x$  and  $y$  are said to **communicate** with each other if  $x \longrightarrow y$  and  $y \longrightarrow x$ .

Notation:  $x \longleftrightarrow y$ .

# Communication is an Equivalence Relation

## Proposition (Communication is an Equivalence Relation)

$\longleftrightarrow$  defines an **equivalence relation** on  $\mathcal{X} \times \mathcal{X}$ . Formally:

1. **(Reflexive)**:  $x \longleftrightarrow x$  for all  $x \in \mathcal{X}$ .
2. **(Symmetric)**: For all  $x, y \in \mathcal{X}$ ,

$$x \longleftrightarrow y \iff y \longleftrightarrow x.$$

3. **(Transitive)**: For all  $x, y, z \in \mathcal{X}$ ,

$$x \longleftrightarrow y, \quad y \longleftrightarrow z \implies x \longleftrightarrow z.$$

## Proof of Proposition

- Reflexive:

$$P^0 = I \iff P_{x,x}^0 = 1 > 0 \iff x \longleftrightarrow x.$$

## Proof of Proposition

- Reflexive:

$$P^0 = I \iff P_{x,x}^0 = 1 > 0 \iff x \longleftrightarrow x.$$

- Symmetric:

$$x \longleftrightarrow y \iff x \longrightarrow y, \quad y \longrightarrow x \iff y \longleftrightarrow x.$$

## Proof of Proposition

- Reflexive:

$$P^0 = I \iff P_{x,x}^0 = 1 > 0 \iff x \longleftrightarrow x.$$

- Symmetric:

$$x \longleftrightarrow y \iff x \longrightarrow y, \quad y \longrightarrow x \iff y \longleftrightarrow x.$$

- Transitive:

$$x \longleftrightarrow y, \quad y \longleftrightarrow z \implies \exists m, n \in \mathbb{N} \cup \{0\} \text{ such that } P_{x,y}^m > 0, \quad P_{y,z}^n > 0$$



## Proof of Proposition

- Reflexive:

$$P^0 = I \iff P_{x,x}^0 = 1 > 0 \iff x \longleftrightarrow x.$$

- Symmetric:

$$x \longleftrightarrow y \iff x \longrightarrow y, \quad y \longrightarrow x \iff y \longleftrightarrow x.$$

- Transitive:

$$x \longleftrightarrow y, \quad y \longleftrightarrow z \implies \exists m, n \in \mathbb{N} \cup \{0\} \text{ such that } P_{x,y}^m > 0, \quad P_{y,z}^n > 0$$

## Proof of Proposition

- Reflexive:

$$P^0 = I \iff P_{x,x}^0 = 1 > 0 \iff x \longleftrightarrow x.$$

- Symmetric:

$$x \longleftrightarrow y \iff x \longrightarrow y, \quad y \longrightarrow x \iff y \longleftrightarrow x.$$

- Transitive:

$$x \longleftrightarrow y, \quad y \longleftrightarrow z \implies \exists m, n \in \mathbb{N} \cup \{0\} \text{ such that } P_{x,y}^m > 0, \quad P_{y,z}^n > 0$$
$$P_{x,z}^{m+n} = \sum_{w \in \mathcal{X}} P_{x,w}^m P_{w,z}^n$$

## Proof of Proposition

- Reflexive:

$$P^0 = I \iff P_{x,x}^0 = 1 > 0 \iff x \longleftrightarrow x.$$

- Symmetric:

$$x \longleftrightarrow y \iff x \longrightarrow y, \quad y \longrightarrow x \iff y \longleftrightarrow x.$$

- Transitive:

$$x \longleftrightarrow y, \quad y \longleftrightarrow z \implies \exists m, n \in \mathbb{N} \cup \{0\} \text{ such that } P_{x,y}^m > 0, \quad P_{y,z}^n > 0$$

$$\begin{aligned} P_{x,z}^{m+n} &= \sum_{w \in \mathcal{X}} P_{x,w}^m P_{w,z}^n \\ &\geq P_{x,y}^m P_{y,z}^n > 0 \end{aligned}$$

## Proof of Proposition

- Reflexive:

$$P^0 = I \iff P_{x,x}^0 = 1 > 0 \iff x \longleftrightarrow x.$$

- Symmetric:

$$x \longleftrightarrow y \iff x \longrightarrow y, \quad y \longrightarrow x \iff y \longleftrightarrow x.$$

- Transitive:

$$x \longleftrightarrow y, \quad y \longleftrightarrow z \implies \exists m, n \in \mathbb{N} \cup \{0\} \text{ such that } P_{x,y}^m > 0, \quad P_{y,z}^n > 0$$

$$P_{x,z}^{m+n} = \sum_{w \in \mathcal{X}} P_{x,w}^m P_{w,z}^n$$

$$\geq P_{x,y}^m P_{y,z}^n > 0$$

$$\implies x \longleftrightarrow z.$$

## Communicating Class

### Definition (Communicating Class)

The communication relation  $\longleftrightarrow$  creates a **partition** of the state space  $\mathcal{X}$ .  
Each element of the partition is referred to as a **communicating class**.

## Open and Closed Communicating Classes

### Definition (Open and Closed Communicating Classes)

A communicating class  $\mathcal{C}$  is said to be **open** if there exists an edge that leaves the class, i.e., there exists  $x \in \mathcal{C}$  and  $y \in \mathcal{C}^c$  such that  $P_{x,y} > 0$ .

A communicating class  $\mathcal{C}$  is said to be **closed** if there is no edge leaving the class, i.e.,

$$P_{x,y} = 0 \quad \forall x \in \mathcal{C}, y \in \mathcal{C}^c.$$

# Irreducibility and Periodicity

# Irreducible Markov Chain

## Definition (Irreducible Markov Chain)

A time-homogeneous DTMC is said to be **irreducible** if its entire state space constitutes a single communicating class.

That is, for all  $x, y \in \mathcal{X}$ , there exists  $n \in \mathbb{N} \cup \{0\}$  such that  $P_{x,y}^n > 0$ .

Remark:

Some textbooks (particularly on RL) refer to irreducibility as **unichain** property.



## Period

- Consider a time-homogeneous DTMC on a discrete state space  $\mathcal{X}$  and TPM  $P$
- For any  $x \in \mathcal{X}$ , let

$$\mathcal{R}(x) := \left\{ n \in \mathbb{N} : P_{x,x}^n > 0 \right\}$$

denote the set of **return times** to  $x$

## Period

- Consider a time-homogeneous DTMC on a discrete state space  $\mathcal{X}$  and TPM  $P$
- For any  $x \in \mathcal{X}$ , let

$$\mathcal{R}(x) := \left\{ n \in \mathbb{N} : P_{x,x}^n > 0 \right\}$$

denote the set of **return times** to  $x$

### Definition (Period)

The **period** of a state  $x \in \mathcal{X}$ , denoted  $d(x)$ , is defined as the greatest common divisor of the set of return times, i.e.,

$$d(x) := \gcd\{n \in \mathbb{N} : P_{x,x}^n > 0\}.$$

## Period

- Consider a time-homogeneous DTMC on a discrete state space  $\mathcal{X}$  and TPM  $P$
- For any  $x \in \mathcal{X}$ , let

$$\mathcal{R}(x) := \left\{ n \in \mathbb{N} : P_{x,x}^n > 0 \right\}$$

denote the set of **return times** to  $x$

### Definition (Period)

The **period** of a state  $x \in \mathcal{X}$ , denoted  $d(x)$ , is defined as the greatest common divisor of the set of return times, i.e.,

$$d(x) := \gcd\{n \in \mathbb{N} : P_{x,x}^n > 0\}.$$

A state  $x$  is called **aperiodic** if  $d(x) = 1$ .

If  $d(x) > 1$ , the state  $x$  is said to be **periodic**.

## Aperiodic Markov Chain

### Definition (Aperiodic Markov Chain)

A time-homogeneous DTMC is said to be **aperiodic** if

1. The Markov chain is irreducible, and
2. The period of every state is 1.

# Class Properties

## Period is a Class Property

### Proposition (Period is a Class Property)

If  $x \longleftrightarrow y$ , then  $d(x) = d(y)$ .

Thus, all states within a communicating class possess the same period.

## Proof of Proposition – 1

- Observe that

$$x \longleftrightarrow y \quad \implies \quad \exists m, n \in \mathbb{N} \cup \{0\} \text{ such that } P_{x,y}^m > 0, \ P_{y,x}^n > 0.$$

## Proof of Proposition – 1

- Observe that

$$x \longleftrightarrow y \implies \exists m, n \in \mathbb{N} \cup \{0\} \text{ such that } P_{x,y}^m > 0, P_{y,x}^n > 0.$$

- Suppose  $r \in \mathcal{R}(x)$ , i.e.,  $P_{x,x}^r > 0$ . Then,



## Proof of Proposition – 1

- Observe that

$$x \longleftrightarrow y \implies \exists m, n \in \mathbb{N} \cup \{0\} \text{ such that } P_{x,y}^m > 0, P_{y,x}^n > 0.$$

- Suppose  $r \in \mathcal{R}(x)$ , i.e.,  $P_{x,x}^r > 0$ . Then,

## Proof of Proposition – 1

- Observe that

$$x \longleftrightarrow y \implies \exists m, n \in \mathbb{N} \cup \{0\} \text{ such that } P_{x,y}^m > 0, P_{y,x}^n > 0.$$

- Suppose  $r \in \mathcal{R}(x)$ , i.e.,  $P_{x,x}^r > 0$ . Then,

$$P_{y,y}^{n+m} \geq P_{y,x}^n \cdot P_{x,y}^m > 0, \quad P_{y,y}^{n+r+m} \geq P_{y,x}^n \cdot P_{x,x}^r \cdot P_{x,y}^m > 0,$$

## Proof of Proposition – 1

- Observe that

$$x \longleftrightarrow y \implies \exists m, n \in \mathbb{N} \cup \{0\} \text{ such that } P_{x,y}^m > 0, P_{y,x}^n > 0.$$

- Suppose  $r \in \mathcal{R}(x)$ , i.e.,  $P_{x,x}^r > 0$ . Then,

$$P_{y,y}^{n+m} \geq P_{y,x}^n \cdot P_{x,y}^m > 0, \quad P_{y,y}^{n+r+m} \geq P_{y,x}^n \cdot P_{x,x}^r \cdot P_{x,y}^m > 0,$$

thereby implying that

$$n + m \in \mathcal{R}(y), \quad n + m + r \in \mathcal{R}(y)$$

## Proof of Proposition – 1

- Observe that

$$x \longleftrightarrow y \implies \exists m, n \in \mathbb{N} \cup \{0\} \text{ such that } P_{x,y}^m > 0, P_{y,x}^n > 0.$$

- Suppose  $r \in \mathcal{R}(x)$ , i.e.,  $P_{x,x}^r > 0$ . Then,

$$P_{y,y}^{n+m} \geq P_{y,x}^n \cdot P_{x,y}^m > 0, \quad P_{y,y}^{n+r+m} \geq P_{y,x}^n \cdot P_{x,x}^r \cdot P_{x,y}^m > 0,$$

thereby implying that

$$\begin{aligned} n+m &\in \mathcal{R}(y), & n+m+r &\in \mathcal{R}(y) \\ \implies d(y) &\mid n+m, & d(y) &\mid n+m+r \end{aligned}$$

## Proof of Proposition – 1

- Observe that

$$x \longleftrightarrow y \implies \exists m, n \in \mathbb{N} \cup \{0\} \text{ such that } P_{x,y}^m > 0, P_{y,x}^n > 0.$$

- Suppose  $r \in \mathcal{R}(x)$ , i.e.,  $P_{x,x}^r > 0$ . Then,

$$P_{y,y}^{n+m} \geq P_{y,x}^n \cdot P_{x,y}^m > 0, \quad P_{y,y}^{n+r+m} \geq P_{y,x}^n \cdot P_{x,x}^r \cdot P_{x,y}^m > 0,$$

thereby implying that

$$\begin{aligned} n+m &\in \mathcal{R}(y), & n+m+r &\in \mathcal{R}(y) \\ \implies d(y) &\mid n+m, & d(y) &\mid n+m+r \\ \implies d(y) &\mid r \end{aligned}$$

## Proof of Proposition – 2

- Thus, if  $r \in \mathcal{R}(x)$ , then  $d(y) \mid r$ , implying that  $d(y)$  is a common divisor of  $\mathcal{R}(x)$

## Proof of Proposition – 2

- Thus, if  $r \in \mathcal{R}(x)$ , then  $d(y) \mid r$ , implying that  $d(y)$  is a common divisor of  $\mathcal{R}(x)$
- This implies  $d(y) \leq d(x)$ , as  $d(x)$  is the greatest common divisor of  $\mathcal{R}(x)$

## Proof of Proposition – 2

- Thus, if  $r \in \mathcal{R}(x)$ , then  $d(y) \mid r$ , implying that  $d(y)$  is a common divisor of  $\mathcal{R}(x)$
- This implies  $d(y) \leq d(x)$ , as  $d(x)$  is the greatest common divisor of  $\mathcal{R}(x)$
- Switching the roles of  $x$  and  $y$ , we can establish that  $d(x) \leq d(y)$



## Transience and Recurrence are Class Properties

### Proposition (Transience and Recurrence are Class Properties)

Transience and recurrence are class properties, i.e., the states within a communicating class are either **all transient** or **all recurrent**.



## Proof of Proposition – 1

- Suppose  $x$  is recurrent, and  $x \longleftrightarrow y$

## Proof of Proposition – 1

- Suppose  $x$  is recurrent, and  $x \longleftrightarrow y$
- By definition of communication, there exist  $m, n \in \mathbb{N} \cup \{0\}$  such that

$$P_{x,y}^m > 0, \quad P_{y,x}^n > 0.$$

## Proof of Proposition - 1

- Suppose  $x$  is recurrent, and  $x \longleftrightarrow y$
- By definition of communication, there exist  $m, n \in \mathbb{N} \cup \{0\}$  such that

$$P_{x,y}^m > 0, \quad P_{y,x}^n > 0.$$

- $x$  recurrent  $\iff \mathbb{E}[N_x \mid X_0 = x] = \sum_{s \in \mathbb{N}} P_{x,x}^s = +\infty$

## Proof of Proposition – 1

- Suppose  $x$  is recurrent, and  $x \longleftrightarrow y$
- By definition of communication, there exist  $m, n \in \mathbb{N} \cup \{0\}$  such that

$$P_{x,y}^m > 0, \quad P_{y,x}^n > 0.$$

- $x$  recurrent  $\iff \mathbb{E}[N_x \mid X_0 = x] = \sum_{s \in \mathbb{N}} P_{x,x}^s = +\infty$
- Then, it follows that

$$\mathbb{E}[N_y \mid X_0 = y] = \sum_{k \in \mathbb{N}} P_{y,y}^k$$

## Proof of Proposition – 1

- Suppose  $x$  is recurrent, and  $x \longleftrightarrow y$
- By definition of communication, there exist  $m, n \in \mathbb{N} \cup \{0\}$  such that

$$P_{x,y}^m > 0, \quad P_{y,x}^n > 0.$$

- $x$  recurrent  $\iff \mathbb{E}[N_x \mid X_0 = x] = \sum_{s \in \mathbb{N}} P_{x,x}^s = +\infty$
- Then, it follows that

$$\mathbb{E}[N_y \mid X_0 = y] = \sum_{k \in \mathbb{N}} P_{y,y}^k$$

## Proof of Proposition – 1

- Suppose  $x$  is recurrent, and  $x \longleftrightarrow y$
- By definition of communication, there exist  $m, n \in \mathbb{N} \cup \{0\}$  such that

$$P_{x,y}^m > 0, \quad P_{y,x}^n > 0.$$

- $x$  recurrent  $\iff \mathbb{E}[N_x \mid X_0 = x] = \sum_{s \in \mathbb{N}} P_{x,x}^s = +\infty$
- Then, it follows that

$$\begin{aligned} \mathbb{E}[N_y \mid X_0 = y] &= \sum_{k \in \mathbb{N}} P_{y,y}^k \\ &\geq P_{y,y}^{m+n+1} + P_{y,y}^{m+n+2} + \dots \end{aligned}$$

## Proof of Proposition – 1

- Suppose  $x$  is recurrent, and  $x \longleftrightarrow y$
- By definition of communication, there exist  $m, n \in \mathbb{N} \cup \{0\}$  such that

$$P_{x,y}^m > 0, \quad P_{y,x}^n > 0.$$

- $x$  recurrent  $\iff \mathbb{E}[N_x \mid X_0 = x] = \sum_{s \in \mathbb{N}} P_{x,x}^s = +\infty$
- Then, it follows that

$$\begin{aligned} \mathbb{E}[N_y \mid X_0 = y] &= \sum_{k \in \mathbb{N}} P_{y,y}^k \\ &\geq P_{y,y}^{m+n+1} + P_{y,y}^{m+n+2} + \dots \\ &= \sum_{s \in \mathbb{N}} P_{y,y}^{m+n+s} \end{aligned}$$



## Proof of Proposition - 1

- Suppose  $x$  is recurrent, and  $x \longleftrightarrow y$
- By definition of communication, there exist  $m, n \in \mathbb{N} \cup \{0\}$  such that

$$P_{x,y}^m > 0, \quad P_{y,x}^n > 0.$$

- $x$  recurrent  $\iff \mathbb{E}[N_x \mid X_0 = x] = \sum_{s \in \mathbb{N}} P_{x,x}^s = +\infty$
- Then, it follows that

$$\begin{aligned} \mathbb{E}[N_y \mid X_0 = y] &= \sum_{k \in \mathbb{N}} P_{y,y}^k \\ &\geq P_{y,y}^{m+n+1} + P_{y,y}^{m+n+2} + \dots \\ &= \sum_{s \in \mathbb{N}} P_{y,y}^{m+n+s} \\ &\geq \sum_{s \in \mathbb{N}} P_{y,x}^n P_{x,x}^s P_{x,y}^m \end{aligned}$$

## Proof of Proposition – 1

- Suppose  $x$  is recurrent, and  $x \longleftrightarrow y$
- By definition of communication, there exist  $m, n \in \mathbb{N} \cup \{0\}$  such that

$$P_{x,y}^m > 0, \quad P_{y,x}^n > 0.$$

- $x$  recurrent  $\iff \mathbb{E}[N_x \mid X_0 = x] = \sum_{s \in \mathbb{N}} P_{x,x}^s = +\infty$
- Then, it follows that

$$\begin{aligned} \mathbb{E}[N_y \mid X_0 = y] &= \sum_{k \in \mathbb{N}} P_{y,y}^k \\ &\geq P_{y,y}^{m+n+1} + P_{y,y}^{m+n+2} + \dots \\ &= \sum_{s \in \mathbb{N}} P_{y,y}^{m+n+s} \\ &\geq \sum_{s \in \mathbb{N}} P_{y,x}^n P_{x,x}^s P_{x,y}^m = P_{y,x}^n P_{x,y}^m \left( \sum_{s \in \mathbb{N}} P_{x,x}^s \right) \end{aligned}$$

## Proof of Proposition - 1

- Suppose  $x$  is recurrent, and  $x \longleftrightarrow y$
- By definition of communication, there exist  $m, n \in \mathbb{N} \cup \{0\}$  such that

$$P_{x,y}^m > 0, \quad P_{y,x}^n > 0.$$

- $x$  recurrent  $\iff \mathbb{E}[N_x \mid X_0 = x] = \sum_{s \in \mathbb{N}} P_{x,x}^s = +\infty$
- Then, it follows that

$$\begin{aligned} \mathbb{E}[N_y \mid X_0 = y] &= \sum_{k \in \mathbb{N}} P_{y,y}^k \\ &\geq P_{y,y}^{m+n+1} + P_{y,y}^{m+n+2} + \dots \\ &= \sum_{s \in \mathbb{N}} P_{y,y}^{m+n+s} \\ &\geq \sum_{s \in \mathbb{N}} P_{y,x}^n P_{x,x}^s P_{x,y}^m = P_{y,x}^n P_{x,y}^m \left( \sum_{s \in \mathbb{N}} P_{x,x}^s \right) = +\infty \end{aligned}$$

## Proof of Proposition – 2

- Suppose  $x$  is transient, and  $x \longleftrightarrow y$

## Proof of Proposition – 2

- Suppose  $x$  is transient, and  $x \longleftrightarrow y$
- By definition of communication, there exist  $m, n \in \mathbb{N} \cup \{0\}$  such that

$$P_{x,y}^m > 0, \quad P_{y,x}^n > 0.$$

## Proof of Proposition - 2

- Suppose  $x$  is transient, and  $x \longleftrightarrow y$
- By definition of communication, there exist  $m, n \in \mathbb{N} \cup \{0\}$  such that

$$P_{x,y}^m > 0, \quad P_{y,x}^n > 0.$$

- $x$  transient  $\iff \mathbb{E}[N_x \mid X_0 = x] = \sum_{s \in \mathbb{N}} P_{x,x}^s < +\infty$

## Proof of Proposition - 2

- Suppose  $x$  is transient, and  $x \longleftrightarrow y$
- By definition of communication, there exist  $m, n \in \mathbb{N} \cup \{0\}$  such that

$$P_{x,y}^m > 0, \quad P_{y,x}^n > 0.$$

- $x$  transient  $\iff \mathbb{E}[N_x \mid X_0 = x] = \sum_{s \in \mathbb{N}} P_{x,x}^s < +\infty$
- Then, it follows that

$$\mathbb{E}[N_y \mid X_0 = y] = \sum_{k \in \mathbb{N}} P_{y,y}^k$$

## Proof of Proposition - 2

- Suppose  $x$  is transient, and  $x \longleftrightarrow y$
- By definition of communication, there exist  $m, n \in \mathbb{N} \cup \{0\}$  such that

$$P_{x,y}^m > 0, \quad P_{y,x}^n > 0.$$

- $x$  transient  $\iff \mathbb{E}[N_x \mid X_0 = x] = \sum_{s \in \mathbb{N}} P_{x,x}^s < +\infty$
- Then, it follows that

$$\mathbb{E}[N_y \mid X_0 = y] = \sum_{k \in \mathbb{N}} P_{y,y}^k$$



## Proof of Proposition - 2

- Suppose  $x$  is transient, and  $x \longleftrightarrow y$
- By definition of communication, there exist  $m, n \in \mathbb{N} \cup \{0\}$  such that

$$P_{x,y}^m > 0, \quad P_{y,x}^n > 0.$$

- $x$  transient  $\iff \mathbb{E}[N_x \mid X_0 = x] = \sum_{s \in \mathbb{N}} P_{x,x}^s < +\infty$
- Then, it follows that

$$\begin{aligned} \mathbb{E}[N_y \mid X_0 = y] &= \sum_{k \in \mathbb{N}} P_{y,y}^k \\ &= \frac{1}{P_{x,y}^m P_{y,x}^n} \sum_{s \in \mathbb{N}} P_{x,y}^m P_{y,y}^s P_{y,x}^n \end{aligned}$$

## Proof of Proposition - 2

- Suppose  $x$  is transient, and  $x \longleftrightarrow y$
- By definition of communication, there exist  $m, n \in \mathbb{N} \cup \{0\}$  such that

$$P_{x,y}^m > 0, \quad P_{y,x}^n > 0.$$

- $x$  transient  $\iff \mathbb{E}[N_x \mid X_0 = x] = \sum_{s \in \mathbb{N}} P_{x,x}^s < +\infty$
- Then, it follows that

$$\begin{aligned} \mathbb{E}[N_y \mid X_0 = y] &= \sum_{k \in \mathbb{N}} P_{y,y}^k \\ &= \frac{1}{P_{x,y}^m P_{y,x}^n} \sum_{s \in \mathbb{N}} P_{x,y}^m P_{y,y}^s P_{y,x}^n \\ &\leq \frac{1}{P_{x,y}^m P_{y,x}^n} \sum_{s \in \mathbb{N}} P_{x,x}^{m+n+s} \end{aligned}$$

## Proof of Proposition - 2

- Suppose  $x$  is transient, and  $x \longleftrightarrow y$
- By definition of communication, there exist  $m, n \in \mathbb{N} \cup \{0\}$  such that

$$P_{x,y}^m > 0, \quad P_{y,x}^n > 0.$$

- $x$  transient  $\iff \mathbb{E}[N_x \mid X_0 = x] = \sum_{s \in \mathbb{N}} P_{x,x}^s < +\infty$
- Then, it follows that

$$\begin{aligned} \mathbb{E}[N_y \mid X_0 = y] &= \sum_{k \in \mathbb{N}} P_{y,y}^k \\ &= \frac{1}{P_{x,y}^m P_{y,x}^n} \sum_{s \in \mathbb{N}} P_{x,y}^m P_{y,y}^s P_{y,x}^n \\ &\leq \frac{1}{P_{x,y}^m P_{y,x}^n} \sum_{s \in \mathbb{N}} P_{x,x}^{m+n+s} \leq \frac{1}{P_{x,y}^m P_{y,x}^n} \sum_{k \in \mathbb{N}} P_{x,x}^k \end{aligned}$$

## Proof of Proposition - 2

- Suppose  $x$  is transient, and  $x \longleftrightarrow y$
- By definition of communication, there exist  $m, n \in \mathbb{N} \cup \{0\}$  such that

$$P_{x,y}^m > 0, \quad P_{y,x}^n > 0.$$

- $x$  transient  $\iff \mathbb{E}[N_x \mid X_0 = x] = \sum_{s \in \mathbb{N}} P_{x,x}^s < +\infty$
- Then, it follows that

$$\begin{aligned} \mathbb{E}[N_y \mid X_0 = y] &= \sum_{k \in \mathbb{N}} P_{y,y}^k \\ &= \frac{1}{P_{x,y}^m P_{y,x}^n} \sum_{s \in \mathbb{N}} P_{x,y}^m P_{y,y}^s P_{y,x}^n \\ &\leq \frac{1}{P_{x,y}^m P_{y,x}^n} \sum_{s \in \mathbb{N}} P_{x,x}^{m+n+s} \leq \frac{1}{P_{x,y}^m P_{y,x}^n} \sum_{k \in \mathbb{N}} P_{x,x}^k < +\infty \end{aligned}$$

## Positive/Null Recurrence are Class Properties

### Proposition (Positive/Null Recurrence are Class Properties)

Positive recurrence and null recurrence are class properties, i.e., the states within a communicating class are either **all positive recurrent** or **all null recurrent**.

A guided proof of this will be demonstrated in the homework.

## An Important Result About Open and Closed Communicating Classes

### Proposition (Result about Open/Closed Communicating Classes)

1. If  $\mathcal{C}$  is an **open** communicating class, then every state within  $\mathcal{C}$  is **transient**.
2. If  $\mathcal{C}$  is a **closed** communicating class, and  $|\mathcal{C}| < +\infty$ , then every state within  $\mathcal{C}$  is **positive recurrent**.

As a corollary, an irreducible DTMC with a finite state space is positive recurrent.