



Stochastic Processes

DTMCs: Recap of Important Results, Ergodicity, Convergence to Stationary Distribution

Karthik P. N.

Assistant Professor, Department of AI

Email: pnkarthik@ai.iith.ac.in

11 April 2025



Recap of Important Results

- Period, transience, positive recurrence, and null recurrence are class properties

Recap of Important Results

- Period, transience, positive recurrence, and null recurrence are class properties
- Open communicating classes are transient.
Closed communicating classes are recurrent.
Finite closed communicating classes are positive recurrent.

Recap of Important Results

- Period, transience, positive recurrence, and null recurrence are class properties
- Open communicating classes are transient.
Closed communicating classes are recurrent.
Finite closed communicating classes are positive recurrent.
- The following are equivalent.

Recap of Important Results

- Period, transience, positive recurrence, and null recurrence are class properties
- Open communicating classes are transient.
Closed communicating classes are recurrent.
Finite closed communicating classes are positive recurrent.
- The following are equivalent.
 - State x is transient.

Recap of Important Results

- Period, transience, positive recurrence, and null recurrence are class properties
- Open communicating classes are transient.
Closed communicating classes are recurrent.
Finite closed communicating classes are positive recurrent.
- The following are equivalent.
 - State x is transient.
 - $\mathbb{E}[N_x \mid X_0 = x] = \sum_{n \in \mathbb{N}} P_{x,x}^n < +\infty$

Recap of Important Results

- Period, transience, positive recurrence, and null recurrence are class properties
- Open communicating classes are transient.
Closed communicating classes are recurrent.
Finite closed communicating classes are positive recurrent.
- The following are equivalent.

- State x is transient.

- $\mathbb{E}[N_x \mid X_0 = x] = \sum_{n \in \mathbb{N}} P_{x,x}^n < +\infty$

- $f_{xx} = \mathbb{P}(\tau_x^{(1)} < +\infty \mid X_0 = x) = \sum_{n \in \mathbb{N}} f_{xx}^{(n)} < 1,$

$$f_{xx}^{(n)} = \mathbb{P}(\tau_x^{(1)} = n \mid X_0 = x).$$

Recap of Important Results

- Period, transience, positive recurrence, and null recurrence are class properties
- Open communicating classes are transient.
Closed communicating classes are recurrent.
Finite closed communicating classes are positive recurrent.

- The following are equivalent.

- State x is transient.

- $\mathbb{E}[N_x \mid X_0 = x] = \sum_{n \in \mathbb{N}} P_{x,x}^n < +\infty$

- $f_{xx} = \mathbb{P}(\tau_x^{(1)} < +\infty \mid X_0 = x) = \sum_{n \in \mathbb{N}} f_{xx}^{(n)} < 1,$

$$f_{xx}^{(n)} = \mathbb{P}(\tau_x^{(1)} = n \mid X_0 = x).$$

- The following are equivalent.

Recap of Important Results

- Period, transience, positive recurrence, and null recurrence are class properties
- Open communicating classes are transient.
Closed communicating classes are recurrent.
Finite closed communicating classes are positive recurrent.

- The following are equivalent.

- State x is transient.

- $\mathbb{E}[N_x \mid X_0 = x] = \sum_{n \in \mathbb{N}} P_{x,x}^n < +\infty$

- $f_{xx} = \mathbb{P}(\tau_x^{(1)} < +\infty \mid X_0 = x) = \sum_{n \in \mathbb{N}} f_{xx}^{(n)} < 1,$

$$f_{xx}^{(n)} = \mathbb{P}(\tau_x^{(1)} = n \mid X_0 = x).$$

- The following are equivalent.
 - State x is recurrent.

Recap of Important Results

- Period, transience, positive recurrence, and null recurrence are class properties
- Open communicating classes are transient.
Closed communicating classes are recurrent.
Finite closed communicating classes are positive recurrent.

- The following are equivalent.

- State x is transient.

- $\mathbb{E}[N_x \mid X_0 = x] = \sum_{n \in \mathbb{N}} P_{x,x}^n < +\infty$

- $f_{xx} = \mathbb{P}(\tau_x^{(1)} < +\infty \mid X_0 = x) = \sum_{n \in \mathbb{N}} f_{xx}^{(n)} < 1,$

$$f_{xx}^{(n)} = \mathbb{P}(\tau_x^{(1)} = n \mid X_0 = x).$$

- The following are equivalent.

- State x is recurrent.

- $\mathbb{E}[N_x \mid X_0 = x] = \sum_{n \in \mathbb{N}} P_{x,x}^n = +\infty$

Recap of Important Results

- Period, transience, positive recurrence, and null recurrence are class properties
- Open communicating classes are transient.
Closed communicating classes are recurrent.
Finite closed communicating classes are positive recurrent.

- The following are equivalent.

- State x is transient.

- $\mathbb{E}[N_x \mid X_0 = x] = \sum_{n \in \mathbb{N}} P_{x,x}^n < +\infty$

- $f_{xx} = \mathbb{P}(\tau_x^{(1)} < +\infty \mid X_0 = x) = \sum_{n \in \mathbb{N}} f_{xx}^{(n)} < 1,$

$$f_{xx}^{(n)} = \mathbb{P}(\tau_x^{(1)} = n \mid X_0 = x).$$

- The following are equivalent.

- State x is recurrent.

- $\mathbb{E}[N_x \mid X_0 = x] = \sum_{n \in \mathbb{N}} P_{x,x}^n = +\infty$

- $f_{xx} = \mathbb{P}(\tau_x^{(1)} < +\infty \mid X_0 = x) = \sum_{n \in \mathbb{N}} f_{xx}^{(n)} = 1,$

$$f_{xx}^{(n)} = \mathbb{P}(\tau_x^{(1)} = n \mid X_0 = x).$$

Recap of Important Results

- If x is transient, then

$$\lim_{n \rightarrow \infty} P_{x,x}^n = 0,$$

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n P_{x,x}^k = 0.$$

Recap of Important Results

- If x is transient, then

$$\lim_{n \rightarrow \infty} P_{x,x}^n = 0, \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n P_{x,x}^k = 0.$$

- If x is recurrent, then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n P_{x,x}^k = \frac{1}{\mu_{xx}}.$$

Recap of Important Results

- If x is transient, then

$$\lim_{n \rightarrow \infty} P_{x,x}^n = 0, \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n P_{x,x}^k = 0.$$

- If x is recurrent, then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n P_{x,x}^k = \frac{1}{\mu_{xx}}.$$

Recap of Important Results

- If x is transient, then

$$\lim_{n \rightarrow \infty} P_{x,x}^n = 0, \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n P_{x,x}^k = 0.$$

- If x is recurrent, then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n P_{x,x}^k = \frac{1}{\mu_{xx}}.$$

In addition, if x is aperiodic, then

$$\lim_{n \rightarrow \infty} P_{x,x}^n = \frac{1}{\mu_{xx}}.$$

- For an irreducible Markov chain,

Unique stationary distribution exists \iff Markov chain is positive recurrent.

Furthermore, in this case, $\pi_x = \frac{1}{\mu_{xx}}$ for all x .

Recap of Important Results

- If $x \longleftrightarrow y$, and y is recurrent, then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n P_{x,y}^k = \frac{1}{\mu_{yy}}.$$

Recap of Important Results

- If $x \longleftrightarrow y$, and y is recurrent, then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n P_{x,y}^k = \frac{1}{\mu_{yy}}.$$

In addition, if y is aperiodic, then

$$\lim_{n \rightarrow \infty} P_{x,y}^n = \frac{1}{\mu_{yy}}.$$

Ergodicity and Ergodic Theorem

Definition (Ergodic Markov Chain)

A time-homogeneous DTMC (on a finite or countably infinite state space) with TPM P is said to be **ergodic** if P is irreducible, aperiodic, and positive recurrent.

Ergodicity and Convergence to Stationary Distribution

Theorem (Ergodicity and Convergence to Stationary Distribution)

Consider a time-homogeneous DTMC $\{X_n\}_{n=0}^{\infty}$ on a discrete state space \mathcal{X} with TPM P . Assume that $X_0 = x$, and for each $n \in \mathbb{N}$, let π_n denote the PMF of X_n . If P is ergodic with associated stationary distribution π , then

$$\lim_{n \rightarrow \infty} d_{\text{TV}}(\pi_n, \pi) = \lim_{n \rightarrow \infty} \frac{1}{2} \|\pi_n - \pi\|_1 = 0,$$

where d_{TV} denotes the **total variation distance**.

Ergodicity and Convergence to Stationary Distribution

Theorem (Ergodicity and Convergence to Stationary Distribution)

Consider a time-homogeneous DTMC $\{X_n\}_{n=0}^{\infty}$ on a discrete state space \mathcal{X} with TPM P . Assume that $X_0 = x$, and for each $n \in \mathbb{N}$, let π_n denote the PMF of X_n . If P is ergodic with associated stationary distribution π , then

$$\lim_{n \rightarrow \infty} d_{\text{TV}}(\pi_n, \pi) = \lim_{n \rightarrow \infty} \frac{1}{2} \|\pi_n - \pi\|_1 = 0,$$

where d_{TV} denotes the **total variation distance**.

Remark: We shall present the proof only for \mathcal{X} finite.

We note that the result holds even when \mathcal{X} is countably infinite.

Proof of Theorem via Coupling

- Consider an **independent copy** of $\{X_n\}_{n=0}^{\infty}$, say $\{Y_n\}_{n=0}^{\infty}$

Proof of Theorem via Coupling

- Consider an **independent copy** of $\{X_n\}_{n=0}^\infty$, say $\{Y_n\}_{n=0}^\infty$
- Assume that $X_0 = x$ and $Y_0 \sim \pi$

Proof of Theorem via Coupling

- Consider an **independent copy** of $\{X_n\}_{n=0}^\infty$, say $\{Y_n\}_{n=0}^\infty$
- Assume that $X_0 = x$ and $Y_0 \sim \pi$
- Let τ be the first time at which both Markov chains meet, i.e.,

$$\tau := \inf\{n \geq 0 : X_n = Y_n\}.$$

Proof of Theorem via Coupling

- Consider an **independent copy** of $\{X_n\}_{n=0}^\infty$, say $\{Y_n\}_{n=0}^\infty$
- Assume that $X_0 = x$ and $Y_0 \sim \pi$
- Let τ be the first time at which both Markov chains meet, i.e.,

$$\tau := \inf\{n \geq 0 : X_n = Y_n\}.$$

- The random variable τ is called the **coupling time**

Proof of Theorem via Coupling

- Consider an **independent copy** of $\{X_n\}_{n=0}^\infty$, say $\{Y_n\}_{n=0}^\infty$
- Assume that $X_0 = x$ and $Y_0 \sim \pi$
- Let τ be the first time at which both Markov chains meet, i.e.,

$$\tau := \inf\{n \geq 0 : X_n = Y_n\}.$$

- The random variable τ is called the **coupling time**
- Let $Z_n = (X_n, Y_n)$; clearly, $\{Z_n\}_{n=0}^\infty$ is a DTMC on $\mathcal{X} \times \mathcal{X}$

Proof of Theorem via Coupling

- Consider an **independent copy** of $\{X_n\}_{n=0}^\infty$, say $\{Y_n\}_{n=0}^\infty$
- Assume that $X_0 = x$ and $Y_0 \sim \pi$
- Let τ be the first time at which both Markov chains meet, i.e.,

$$\tau := \inf\{n \geq 0 : X_n = Y_n\}.$$

- The random variable τ is called the **coupling time**
- Let $Z_n = (X_n, Y_n)$; clearly, $\{Z_n\}_{n=0}^\infty$ is a DTMC on $\mathcal{X} \times \mathcal{X}$
- Define the TPM Q of $\{Z_n\}_{n=0}^\infty$ as

$$Q_{(x,w),(y,z)} = P_{x,y} \cdot P_{w,z}.$$

Proof of Theorem via Coupling

- Consider an **independent copy** of $\{X_n\}_{n=0}^\infty$, say $\{Y_n\}_{n=0}^\infty$
- Assume that $X_0 = x$ and $Y_0 \sim \pi$
- Let τ be the first time at which both Markov chains meet, i.e.,

$$\tau := \inf\{n \geq 0 : X_n = Y_n\}.$$

- The random variable τ is called the **coupling time**
- Let $Z_n = (X_n, Y_n)$; clearly, $\{Z_n\}_{n=0}^\infty$ is a DTMC on $\mathcal{X} \times \mathcal{X}$
- Define the TPM Q of $\{Z_n\}_{n=0}^\infty$ as

$$Q_{(x,w),(y,z)} = P_{x,y} \cdot P_{w,z}.$$

- By Chapman-Kolmogorov,

$$Q_{(x,w),(y,z)}^n = P_{x,y}^n \cdot P_{w,z}^n \quad \forall n \in \mathbb{N}.$$

Proof of Theorem via Coupling

- Because $\{X_n\}_{n=0}^{\infty}$ and $\{Y_n\}_{n=0}^{\infty}$ are irreducible and aperiodic, there exists $N \in \mathbb{N}$ sufficiently large such that

$$P^n > 0 \quad \forall n \geq N.$$

Proof of Theorem via Coupling

- Because $\{X_n\}_{n=0}^{\infty}$ and $\{Y_n\}_{n=0}^{\infty}$ are irreducible and aperiodic, there exists $N \in \mathbb{N}$ sufficiently large such that

$$P^n > 0 \quad \forall n \geq N.$$

- As a consequence, it follows that $Q^n > 0$ for all $n \geq N$.
This in turn implies $\{Z_n\}_{n=0}^{\infty}$ is irreducible and aperiodic

Proof of Theorem via Coupling

- Because $\{X_n\}_{n=0}^{\infty}$ and $\{Y_n\}_{n=0}^{\infty}$ are irreducible and aperiodic, there exists $N \in \mathbb{N}$ sufficiently large such that

$$P^n > 0 \quad \forall n \geq N.$$

- As a consequence, it follows that $Q^n > 0$ for all $n \geq N$.
This in turn implies $\{Z_n\}_{n=0}^{\infty}$ is irreducible and aperiodic
- Important observation:**
 $\lambda = 1$ is a simple eigenvalue of Q^n for all $n \geq N$.
Therefore, there exists a unique probability vector θ such that $\theta = \theta Q$.
This means that $\{Z_n\}_{n=0}^{\infty}$ is positive recurrent, and $\mathbb{P}(\tau < +\infty) = 1$.

Proof of Theorem via Coupling

- Because $\{X_n\}_{n=0}^{\infty}$ and $\{Y_n\}_{n=0}^{\infty}$ are irreducible and aperiodic, there exists $N \in \mathbb{N}$ sufficiently large such that

$$P^n > 0 \quad \forall n \geq N.$$

- As a consequence, it follows that $Q^n > 0$ for all $n \geq N$.
This in turn implies $\{Z_n\}_{n=0}^{\infty}$ is irreducible and aperiodic
- Important observation:**
 $\lambda = 1$ is a simple eigenvalue of Q^n for all $n \geq N$.
Therefore, there exists a unique probability vector θ such that $\theta = \theta Q$.
This means that $\{Z_n\}_{n=0}^{\infty}$ is positive recurrent, and $\mathbb{P}(\tau < +\infty) = 1$.
- A simple observation shows that θ defined via

$$\theta(x, w) = \pi(x) \cdot \pi(w)$$

is a stationary distribution for Q , and the only such one

Proof of Theorem via Coupling

- From the coupling time onwards, the two processes $\{X_n\}_{n=0}^{\infty}$ and $\{Y_n\}_{n=0}^{\infty}$ will have identical statistics:

$$\mathbb{P}(X_n = \gamma, n \geq \tau \mid X_\tau = x) = \mathbb{P}(Y_n = \gamma, n \geq \tau \mid Y_\tau = x)$$

Proof of Theorem via Coupling

- From the coupling time onwards, the two processes $\{X_n\}_{n=0}^{\infty}$ and $\{Y_n\}_{n=0}^{\infty}$ will have identical statistics:

$$\mathbb{P}(X_n = \gamma, n \geq \tau \mid X_\tau = x) = \mathbb{P}(Y_n = \gamma, n \geq \tau \mid Y_\tau = x)$$

- Finally, we note that

$$|\pi_n(\gamma) - \pi(\gamma)| = \left| \mathbb{P}(X_n = \gamma) - \mathbb{P}(Y_n = \gamma) \right|$$

Proof of Theorem via Coupling

- From the coupling time onwards, the two processes $\{X_n\}_{n=0}^{\infty}$ and $\{Y_n\}_{n=0}^{\infty}$ will have identical statistics:

$$\mathbb{P}(X_n = \gamma, n \geq \tau \mid X_\tau = x) = \mathbb{P}(Y_n = \gamma, n \geq \tau \mid Y_\tau = x)$$

- Finally, we note that

$$|\pi_n(\gamma) - \pi(\gamma)| = \left| \mathbb{P}(X_n = \gamma) - \mathbb{P}(Y_n = \gamma) \right|$$

Proof of Theorem via Coupling

- From the coupling time onwards, the two processes $\{X_n\}_{n=0}^{\infty}$ and $\{Y_n\}_{n=0}^{\infty}$ will have identical statistics:

$$\mathbb{P}(X_n = \gamma, n \geq \tau \mid X_\tau = x) = \mathbb{P}(Y_n = \gamma, n \geq \tau \mid Y_\tau = x)$$

- Finally, we note that

$$\begin{aligned} |\pi_n(\gamma) - \pi(\gamma)| &= \left| \mathbb{P}(X_n = \gamma) - \mathbb{P}(Y_n = \gamma) \right| \\ &= \left| \mathbb{P}(X_n = \gamma, \tau > n) - \mathbb{P}(Y_n = \gamma, \tau > n) \right| \end{aligned}$$

Proof of Theorem via Coupling

- From the coupling time onwards, the two processes $\{X_n\}_{n=0}^{\infty}$ and $\{Y_n\}_{n=0}^{\infty}$ will have identical statistics:

$$\mathbb{P}(X_n = \gamma, n \geq \tau \mid X_\tau = x) = \mathbb{P}(Y_n = \gamma, n \geq \tau \mid Y_\tau = x)$$

- Finally, we note that

$$\begin{aligned} |\pi_n(\gamma) - \pi(\gamma)| &= \left| \mathbb{P}(X_n = \gamma) - \mathbb{P}(Y_n = \gamma) \right| \\ &= \left| \mathbb{P}(X_n = \gamma, \tau > n) - \mathbb{P}(Y_n = \gamma, \tau > n) \right| \\ &\leq 2 \mathbb{P}(\tau > n) \end{aligned}$$

- Taking limits as $n \rightarrow \infty$, and noting that $\mathbb{P}(\tau < +\infty) = 1$, we arrive at the result