



Stochastic Processes

Hitting Times and Recurrence, Transience, Positive/Null Recurrence

Karthik P. N.

Assistant Professor, Department of AI

Email: pnkarthik@ai.iith.ac.in

21 March 2025

Hitting Times

Definition (Hitting Times)

Let $\{X_n\}_{n=1}^{\infty}$ be DTMC on a discrete state space \mathcal{X} with TPM P .

Fix $y \in \mathcal{X}$.

Let $\tau_y^{(0)} := 0$, and

$$\tau_y^{(k)} = \inf\{n > \tau_y^{(k-1)} : X_n = y\}, \quad k \in \mathbb{N}.$$

For each $k \in \mathbb{N}$, the random variable $\tau_y^{(k)}$ is called the “ **k th hitting time of state y** ”.

Exercise:

For each $k \in \mathbb{N}$, verify that $\{\tau_y^k = n\} \in \sigma(X_1, \dots, X_n)$ for all n .


An Important Observation Regarding $\tau_y^{(k)}$

Lemma (An Important Observation Regarding $\tau_y^{(k)}$)

For each $k \in \mathbb{N}$, suppose that $\mathbb{P}(\tau_y^{(k)} < +\infty) = 1$.

Then, the history up to $\tau_y^{(k)}$ is independent of the future **unconditionally**, i.e.,

$$(X_1, \dots, X_{\tau_y^{(k)}-1}) \perp\!\!\!\perp (X_{\tau_y^{(k)}+1}, X_{\tau_y^{(k)}+2}, \dots).$$



An Important Observation Regarding $\tau_Y^{(k)}$

Lemma (An Important Observation Regarding $\tau_Y^{(k)}$)

For each $k \in \mathbb{N}$, suppose that $\mathbb{P}(\tau_Y^{(k)} < +\infty) = 1$.

Then, the history up to $\tau_Y^{(k)}$ is independent of the future **unconditionally**, i.e.,

$$(X_1, \dots, X_{\tau_Y^{(k)}-1}) \perp\!\!\!\perp (X_{\tau_Y^{(k)}+1}, X_{\tau_Y^{(k)}+2}, \dots).$$

Proof of Lemma:

- According to strong Markov property,

$$(X_1, \dots, X_{\tau_Y^{(k)}-1}) \perp\!\!\!\perp (X_{\tau_Y^{(k)}+1}, X_{\tau_Y^{(k)}+2}, \dots) \mid X_{\tau_Y^{(k)}}.$$

An Important Observation Regarding $\tau_Y^{(k)}$

Lemma (An Important Observation Regarding $\tau_Y^{(k)}$)

For each $k \in \mathbb{N}$, suppose that $\mathbb{P}(\tau_Y^{(k)} < +\infty) = 1$.

Then, the history up to $\tau_Y^{(k)}$ is independent of the future **unconditionally**, i.e.,

$$(X_1, \dots, X_{\tau_Y^{(k)}-1}) \perp\!\!\!\perp (X_{\tau_Y^{(k)}+1}, X_{\tau_Y^{(k)}+2}, \dots).$$

Proof of Lemma:

- According to strong Markov property,

$$(X_1, \dots, X_{\tau_Y^{(k)}-1}) \perp\!\!\!\perp (X_{\tau_Y^{(k)}+1}, X_{\tau_Y^{(k)}+2}, \dots) \mid X_{\tau_Y^{(k)}}.$$

- However, $X_{\tau_Y^{(k)}}$ is a constant random variable (taking the constant value y).
Thus, conditioning on $X_{\tau_Y^{(k)}}$ is as good as not conditioning at all.

IID Block Structure


Lemma (IID Block Structure)

For each $k \in \mathbb{N} \cup \{0\}$, define the k th block \mathcal{B}_k as

$$\mathcal{B}_k := (X_{\tau_Y^{(k-1)}+1}, X_{\tau_Y^{(k-1)}+2}, \dots, X_{\tau_Y^{(k)}-1}), \quad k \in \mathbb{N} \cup \{0\}.$$

Suppose that $\mathbb{P}(\tau_Y^{(k)} < +\infty) = 1$ for all k . Then, the following holds.

1. The collection $\{\mathcal{B}_1, \mathcal{B}_2, \mathcal{B}_3, \dots\}$ is independent.
2. The collection $\{\mathcal{B}_2, \mathcal{B}_3, \dots\}$ is identically distributed.



Proof of Lemma

- $(X_1, \dots, X_{\tau_Y^{(k)}-1}) \perp\!\!\!\perp (X_{\tau_Y^{(k)}+1}, X_{\tau_Y^{(k)}+1}, \dots) \implies B_k \perp\!\!\!\perp B_{k+1}$

Proof of Lemma

- $(X_1, \dots, X_{\tau_Y^{(k)}-1}) \perp\!\!\!\perp (X_{\tau_Y^{(k)}+1}, X_{\tau_Y^{(k)}+1}, \dots) \implies B_k \perp\!\!\!\perp B_{k+1}$
- To show identical distribution of $\{B_2, B_3, \dots\}$, it suffices to show that

$$X_{\tau_Y^{(1)}+1} \stackrel{d.}{=} X_{\tau_Y^{(2)}+1} \stackrel{d.}{=} X_{\tau_Y^{(3)}+1} \stackrel{d.}{=} X_{\tau_Y^{(4)}+1} \cdots$$

Proof of Lemma

- $(X_1, \dots, X_{\tau_Y^{(k)}-1}) \perp\!\!\!\perp (X_{\tau_Y^{(k)}+1}, X_{\tau_Y^{(k)}+1}, \dots) \implies B_k \perp\!\!\!\perp B_{k+1}$
- To show identical distribution of $\{B_2, B_3, \dots\}$, it suffices to show that

$$X_{\tau_Y^{(1)}+1} \stackrel{d.}{=} X_{\tau_Y^{(2)}+1} \stackrel{d.}{=} X_{\tau_Y^{(3)}+1} \stackrel{d.}{=} X_{\tau_Y^{(4)}+1} \cdots$$

- For any $k \in \mathbb{N}$, we have

$$\mathbb{P}(X_{\tau_Y^{(k)}+1} = j) = \mathbb{P}(X_{\tau_Y^{(k)}+1} = j \mid X_{\tau_Y^{(k)}} = y)$$

Proof of Lemma

- $(X_1, \dots, X_{\tau_Y^{(k)}-1}) \perp\!\!\!\perp (X_{\tau_Y^{(k)}+1}, X_{\tau_Y^{(k)}+1}, \dots) \implies B_k \perp\!\!\!\perp B_{k+1}$
- To show identical distribution of $\{B_2, B_3, \dots\}$, it suffices to show that

$$X_{\tau_Y^{(1)}+1} \stackrel{d.}{=} X_{\tau_Y^{(2)}+1} \stackrel{d.}{=} X_{\tau_Y^{(3)}+1} \stackrel{d.}{=} X_{\tau_Y^{(4)}+1} \cdots$$

- For any $k \in \mathbb{N}$, we have

$$\mathbb{P}(X_{\tau_Y^{(k)}+1} = j) = \mathbb{P}(X_{\tau_Y^{(k)}+1} = j \mid X_{\tau_Y^{(k)}} = y)$$

Proof of Lemma

- $(X_1, \dots, X_{\tau_Y^{(k)}-1}) \perp\!\!\!\perp (X_{\tau_Y^{(k)}+1}, X_{\tau_Y^{(k)}+1}, \dots) \implies B_k \perp\!\!\!\perp B_{k+1}$
- To show identical distribution of $\{B_2, B_3, \dots\}$, it suffices to show that

$$X_{\tau_Y^{(1)}+1} \stackrel{d.}{=} X_{\tau_Y^{(2)}+1} \stackrel{d.}{=} X_{\tau_Y^{(3)}+1} \stackrel{d.}{=} X_{\tau_Y^{(4)}+1} \cdots$$

- For any $k \in \mathbb{N}$, we have

$$\begin{aligned} \mathbb{P}(X_{\tau_Y^{(k)}+1} = j) &= \mathbb{P}(X_{\tau_Y^{(k)}+1} = j \mid X_{\tau_Y^{(k)}} = y) \\ &= \mathbb{P}(X_2 = j \mid X_1 = y) \end{aligned}$$

Proof of Lemma

- $(X_1, \dots, X_{\tau_Y^{(k)}-1}) \perp\!\!\!\perp (X_{\tau_Y^{(k)}+1}, X_{\tau_Y^{(k)}+1}, \dots) \implies B_k \perp\!\!\!\perp B_{k+1}$
- To show identical distribution of $\{B_2, B_3, \dots\}$, it suffices to show that

$$X_{\tau_Y^{(1)}+1} \stackrel{d.}{=} X_{\tau_Y^{(2)}+1} \stackrel{d.}{=} X_{\tau_Y^{(3)}+1} \stackrel{d.}{=} X_{\tau_Y^{(4)}+1} \cdots$$

- For any $k \in \mathbb{N}$, we have

$$\begin{aligned} \mathbb{P}(X_{\tau_Y^{(k)}+1} = j) &= \mathbb{P}(X_{\tau_Y^{(k)}+1} = j \mid X_{\tau_Y^{(k)}} = y) \\ &= \mathbb{P}(X_2 = j \mid X_1 = y) \\ &= P_{y,j}. \end{aligned}$$

Recurrence Times

Definition (Recurrence Time)

Let $\{X_n\}_{n=1}^{\infty}$ be DTMC on a discrete state space \mathcal{X} with TPM P .

Fix $y \in \mathcal{X}$.

Let $\tau_y^{(0)} := 0$, and

$$\tau_y^{(k)} = \inf\{n > \tau_y^{(k-1)} : X_n = y\}, \quad k \in \mathbb{N}.$$

The k th recurrence time associated with state y is defined as

$$H_y^{(k)} := \tau_y^{(k)} - \tau_y^{(k-1)}, \quad k \in \mathbb{N}.$$

Recurrence Times

Definition (Recurrence Time)

Let $\{X_n\}_{n=1}^\infty$ be DTMC on a discrete state space \mathcal{X} with TPM P .

Fix $y \in \mathcal{X}$.

Let $\tau_y^{(0)} := 0$, and

$$\tau_y^{(k)} = \inf\{n > \tau_y^{(k-1)} : X_n = y\}, \quad k \in \mathbb{N}.$$

The k th recurrence time associated with state y is defined as

$$H_y^{(k)} := \tau_y^{(k)} - \tau_y^{(k-1)}, \quad k \in \mathbb{N}.$$

Note: $H_y^{(k)}$ is the (random) length of block \mathcal{B}_k .

I.I.D. Property of Recurrence Times

Lemma (I.I.D. Property of Recurrence Times)

Suppose that $\mathbb{P}(\tau_Y^{(k)} < +\infty) = 1$ for all $k \in \mathbb{N}$.

Then, $\{H_Y^{(k)}\}_{k=1}^\infty$ is a sequence of **i.i.d. random variables**.

I.I.D. Property of Recurrence Times

Lemma (I.I.D. Property of Recurrence Times)

Suppose that $\mathbb{P}(\tau_Y^{(k)} < +\infty) = 1$ for all $k \in \mathbb{N}$.

Then, $\{H_Y^{(k)}\}_{k=1}^\infty$ is a sequence of **i.i.d. random variables**.

Proof of Lemma:

- We note that

$$\{H_Y^{(k)} = n\} \in \sigma(X_{\tau_Y^{(k-1)}+1}, \dots, X_{\tau_Y^{(k-1)}+n})$$

I.I.D. Property of Recurrence Times

Lemma (I.I.D. Property of Recurrence Times)

Suppose that $\mathbb{P}(\tau_Y^{(k)} < +\infty) = 1$ for all $k \in \mathbb{N}$.

Then, $\{H_Y^{(k)}\}_{k=1}^\infty$ is a sequence of **i.i.d. random variables**.

Proof of Lemma:

- We note that

$$\{H_Y^{(k)} = n\} \in \sigma(X_{\tau_Y^{(k-1)}+1}, \dots, X_{\tau_Y^{(k-1)}+n})$$

- The result follows from the IID nature of the blocks $\{B_2, B_3, \dots\}$.

Some Tidbits

- For $x, y \in \mathcal{X}$, let

$$f_{xy}^{(n)} := \mathbb{P}(\tau_y^{(1)} = n \mid X_1 = x).$$

$f_{xy}^{(n)}$: **probability of first visit to state y at time n , starting from state x .**

Some Tidbits

- For $x, y \in \mathcal{X}$, let

$$f_{xy}^{(n)} := \mathbb{P}(\tau_y^{(1)} = n \mid X_1 = x).$$

$f_{xy}^{(n)}$: **probability of first visit to state y at time n , starting from state x .**

- Let f_{xy} be defined as

$$f_{xy} = \mathbb{P}(\tau_y^{(1)} < +\infty \mid X_1 = x) =$$

Some Tidbits

- For $x, y \in \mathcal{X}$, let

$$f_{xy}^{(n)} := \mathbb{P}(\tau_y^{(1)} = n \mid X_1 = x).$$

$f_{xy}^{(n)}$: **probability of first visit to state y at time n , starting from state x .**

- Let f_{xy} be defined as

$$f_{xy} = \mathbb{P}(\tau_y^{(1)} < +\infty \mid X_1 = x) =$$

Some Tidbits

- For $x, y \in \mathcal{X}$, let

$$f_{xy}^{(n)} := \mathbb{P}(\tau_y^{(1)} = n \mid X_1 = x).$$

$f_{xy}^{(n)}$: **probability of first visit to state y at time n , starting from state x .**

- Let f_{xy} be defined as

$$f_{xy} = \mathbb{P}(\tau_y^{(1)} < +\infty \mid X_1 = x) = \sum_{n \in \mathbb{N}} f_{xy}^{(n)}.$$

f_{xy} : **probability of eventually visiting state y , starting from state x .**

- If $f_{xy} = 1$ for all $x \in \mathcal{X}$, then $\mathbb{P}(\tau_y^{(1)} < +\infty) = 1$, and hence $\tau_y^{(1)}$ is a stopping time.

Some Tidbits

- For $x, y \in \mathcal{X}$, let

$$f_{xy}^{(n)} := \mathbb{P}(\tau_y^{(1)} = n \mid X_1 = x).$$

$f_{xy}^{(n)}$: **probability of first visit to state y at time n , starting from state x .**

- Let f_{xy} be defined as

$$f_{xy} = \mathbb{P}(\tau_y^{(1)} < +\infty \mid X_1 = x) = \sum_{n \in \mathbb{N}} f_{xy}^{(n)}.$$

f_{xy} : **probability of eventually visiting state y , starting from state x .**

- If $f_{xy} = 1$ for all $x \in \mathcal{X}$, then $\mathbb{P}(\tau_y^{(1)} < +\infty) = 1$, and hence $\tau_y^{(1)}$ is a stopping time.
- $1 - f_{xy}$: probability that starting from x , the state y is **never** visited

Recurrent and Transient States

Definition (Recurrent and Transient States)

A state $x \in \mathcal{X}$ is called **recurrent** if $f_{xx} = 1$.

If $f_{xx} < 1$, then x is called a **transient** state.

Remarks:

- The collection

$$\{f_{xx}^{(1)}, f_{xx}^{(2)}, \dots, 1 - f_{xx}\}$$

defines a valid PMF on $\mathbb{N} \cup \{+\infty\}$.

Recurrent and Transient States

Definition (Recurrent and Transient States)

A state $x \in \mathcal{X}$ is called **recurrent** if $f_{xx} = 1$.

If $f_{xx} < 1$, then x is called a **transient** state.

Remarks:

- The collection

$$\{f_{xx}^{(1)}, f_{xx}^{(2)}, \dots, 1 - f_{xx}\}$$

defines a valid PMF on $\mathbb{N} \cup \{+\infty\}$.

- The above PMF is called **first recurrence time distribution**.

Mean Recurrence Time

Definition (Mean Recurrence Time)

The **mean recurrence time** of a state $x \in \mathcal{X}$ is denoted by μ_{xx} and is defined by

$$\mu_{xx} := \mathbb{E}[\tau_x^{(1)} \mid X_1 = x].$$

Remarks:

- If x is transient, then $\mu_{xx} = +\infty$.

Mean Recurrence Time

Definition (Mean Recurrence Time)

The **mean recurrence time** of a state $x \in \mathcal{X}$ is denoted by μ_{xx} and is defined by

$$\mu_{xx} := \mathbb{E}[\tau_x^{(1)} \mid X_1 = x].$$

Remarks:

- If x is transient, then $\mu_{xx} = +\infty$.
- If x is recurrent, then

$$\mu_{xx} = \sum_{n \in \mathbb{N}} n f_{xx}^{(n)}.$$

Positive Recurrent and Null Recurrent States

Definition (Positive / Null Recurrent States)

A recurrent state $x \in \mathcal{X}$ is called **positive recurrent** if $\mu_{xx} < +\infty$.

Else, if $\mu_{xx} = +\infty$, then x is called **null recurrent**.

Proposition

Proposition

Fix $x, y \in \mathcal{X}$. Then,

$$\mathbb{P}(\text{state } y \text{ is visited exactly } k \text{ times} \mid X_1 = x) = \begin{cases} , & k = 0, \\ , & k \in \mathbb{N}. \end{cases}$$

Proposition

Proposition

Fix $x, y \in \mathcal{X}$. Then,

$$\mathbb{P}(\text{state } y \text{ is visited exactly } k \text{ times} \mid X_1 = x) = \begin{cases} 1 - f_{xy}, & k = 0, \\ f_{xy} f_{yy}^{k-1} (1 - f_{yy}), & k \in \mathbb{N}. \end{cases}$$

Proof of Proposition ($k \in \mathbb{N}$)

$\mathbb{P}(\text{state } y \text{ is visited exactly } k \text{ times} \mid X_1 = x)$

Proof of Proposition ($k \in \mathbb{N}$)

$$\begin{aligned} & \mathbb{P}(\text{state } y \text{ is visited exactly } k \text{ times} \mid X_1 = x) \\ = & \mathbb{P}(\tau_y^{(1)} < +\infty, \tau_y^{(2)} < +\infty, \dots, \tau_y^{(k)} < +\infty, \tau_y^{(k+1)} = +\infty \mid X_1 = x) \end{aligned}$$

Proof of Proposition ($k \in \mathbb{N}$)

$\mathbb{P}(\text{state } y \text{ is visited exactly } k \text{ times} \mid X_1 = x)$

$$= \mathbb{P}(\tau_y^{(1)} < +\infty, \tau_y^{(2)} < +\infty, \dots, \tau_y^{(k)} < +\infty, \tau_y^{(k+1)} = +\infty \mid X_1 = x)$$

$$= \mathbb{P}(\tau_y^{(1)} < +\infty, \tau_y^{(2)} - \tau_y^{(1)} < +\infty, \dots, \tau_y^{(k)} - \tau_1^{(k-1)} < +\infty, \tau_y^{(k+1)} - \tau_1^{(k)} = +\infty \mid X_1 = x)$$

Proof of Proposition ($k \in \mathbb{N}$)

$$\begin{aligned}
 & \mathbb{P}(\text{state } y \text{ is visited exactly } k \text{ times} \mid X_1 = x) \\
 = & \mathbb{P}(\tau_y^{(1)} < +\infty, \tau_y^{(2)} < +\infty, \dots, \tau_y^{(k)} < +\infty, \tau_y^{(k+1)} = +\infty \mid X_1 = x) \\
 = & \mathbb{P}(\tau_y^{(1)} < +\infty, \tau_y^{(2)} - \tau_y^{(1)} < +\infty, \dots, \tau_y^{(k)} - \tau_1^{(k-1)} < +\infty, \tau_y^{(k+1)} - \tau_1^{(k)} = +\infty \mid X_1 = x) \\
 = & \mathbb{P}(H_y^{(1)} < +\infty, H_y^{(2)} < +\infty, \dots, H_y^{(k)} < +\infty, H_y^{(k+1)} = +\infty \mid X_1 = x)
 \end{aligned}$$

Proof of Proposition ($k \in \mathbb{N}$)

$$\begin{aligned}
 & \mathbb{P}(\text{state } y \text{ is visited exactly } k \text{ times} \mid X_1 = x) \\
 = & \mathbb{P}(\tau_y^{(1)} < +\infty, \tau_y^{(2)} < +\infty, \dots, \tau_y^{(k)} < +\infty, \tau_y^{(k+1)} = +\infty \mid X_1 = x) \\
 = & \mathbb{P}(\tau_y^{(1)} < +\infty, \tau_y^{(2)} - \tau_y^{(1)} < +\infty, \dots, \tau_y^{(k)} - \tau_1^{(k-1)} < +\infty, \tau_y^{(k+1)} - \tau_1^{(k)} = +\infty \mid X_1 = x) \\
 = & \mathbb{P}(H_y^{(1)} < +\infty, H_y^{(2)} < +\infty, \dots, H_y^{(k)} < +\infty, H_y^{(k+1)} = +\infty \mid X_1 = x) \\
 = & \mathbb{P}(H_y^{(1)} < +\infty \mid X_1 = x) \times \left(\prod_{\ell=2}^k \mathbb{P}(H_y^{(\ell)} < +\infty) \right) \times \mathbb{P}(H_y^{(k+1)} = +\infty)
 \end{aligned}$$

Proof of Proposition ($k \in \mathbb{N}$)

$$\begin{aligned}
 & \mathbb{P}(\text{state } y \text{ is visited exactly } k \text{ times} \mid X_1 = x) \\
 = & \mathbb{P}(\tau_y^{(1)} < +\infty, \tau_y^{(2)} < +\infty, \dots, \tau_y^{(k)} < +\infty, \tau_y^{(k+1)} = +\infty \mid X_1 = x) \\
 = & \mathbb{P}(\tau_y^{(1)} < +\infty, \tau_y^{(2)} - \tau_y^{(1)} < +\infty, \dots, \tau_y^{(k)} - \tau_1^{(k-1)} < +\infty, \tau_y^{(k+1)} - \tau_1^{(k)} = +\infty \mid X_1 = x) \\
 = & \mathbb{P}(H_y^{(1)} < +\infty, H_y^{(2)} < +\infty, \dots, H_y^{(k)} < +\infty, H_y^{(k+1)} = +\infty \mid X_1 = x) \\
 = & \mathbb{P}(H_y^{(1)} < +\infty \mid X_1 = x) \times \left(\prod_{\ell=2}^k \mathbb{P}(H_y^{(\ell)} < +\infty) \right) \times \mathbb{P}(H_y^{(k+1)} = +\infty) \\
 = & f_{xy} \times \left(\prod_{\ell=2}^k f_{yy} \right) \times (1 - f_{yy}).
 \end{aligned}$$

Corollary to Proposition

Proposition

Fix $x, y \in \mathcal{X}$. Then,

$$\mathbb{P}(\text{state } y \text{ is visited exactly } k \text{ times} \mid X_1 = x) = \begin{cases} 1 - f_{xy}, & k = 0, \\ f_{xy} f_{yy}^{k-1} (1 - f_{yy}), & k \in \mathbb{N}. \end{cases}$$

Corollary

For any $x, y \in \mathcal{X}$,

$$\mathbb{P}(\text{state } y \text{ is visited finitely many times} \mid X_1 = x) =$$

Corollary to Proposition

Proposition

Fix $x, y \in \mathcal{X}$. Then,

$$\mathbb{P}(\text{state } y \text{ is visited exactly } k \text{ times} \mid X_1 = x) = \begin{cases} 1 - f_{xy}, & k = 0, \\ f_{xy} f_{yy}^{k-1} (1 - f_{yy}), & k \in \mathbb{N}. \end{cases}$$

Corollary

For any $x, y \in \mathcal{X}$,

$$\mathbb{P}(\text{state } y \text{ is visited finitely many times} \mid X_1 = x) = \begin{cases} 1, & f_{yy} < 1, \\ 1 - f_{xy}, & f_{yy} = 1. \end{cases}$$