

# Behind the Scenes of $Ax = b$ : Axioms and an Open Question



P. N. Karthik

Research Fellow,  
Institute of Data Science,  
National University of Singapore

Based on joint work with Rajesh Sundaresan

23 March 2022

# Linear Inverse Problem

# Linear Inverse Problem

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{k1} & a_{k2} & \cdots & a_{kn} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_k \end{bmatrix}$$

# Linear Inverse Problem

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{k1} & a_{k2} & \cdots & a_{kn} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_k \end{bmatrix}$$

$A$        $x$        $b$

# Linear Inverse Problem

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{k1} & a_{k2} & \cdots & a_{kn} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_k \end{bmatrix}$$

  $A$        $x$        $b$  

# Linear Inverse Problem

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{k1} & a_{k2} & \cdots & a_{kn} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_k \end{bmatrix}$$

  $A$         $x$        $b$  

# Linear Inverse Problem

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{k1} & a_{k2} & \cdots & a_{kn} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_k \end{bmatrix}$$

  $A$         $x$        $b$  

- Example 1: image reconstruction

# Linear Inverse Problem

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{k1} & a_{k2} & \cdots & a_{kn} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_k \end{bmatrix}$$

  $A$         $x$        $b$  

- Example 1: image reconstruction
  - The domain of an image is its pixels (say  $n$  pixels)

# Linear Inverse Problem

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{k1} & a_{k2} & \cdots & a_{kn} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_k \end{bmatrix}$$

  $A$         $x$        $b$  

- Example 1: image reconstruction

- The domain of an image is its pixels (say  $n$  pixels)
- $f_j$ : indicator of the  $j$ th pixel;

$$f = \sum_{j=1}^n x_j f_j$$

# Linear Inverse Problem

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{k1} & a_{k2} & \cdots & a_{kn} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_k \end{bmatrix}$$

  $A$         $x$        $b$  

- Example 1: image reconstruction

- The domain of an image is its pixels (say  $n$  pixels)

- $f_j$ : indicator of the  $j$ th pixel;

$$f = \sum_{j=1}^n x_j f_j$$

- Linear measurements

$$R_i f_j = a_{ij}, \quad R_i f = b_i, \quad i = 1, \dots, k$$

# Linear Inverse Problem

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{k1} & a_{k2} & \cdots & a_{kn} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_k \end{bmatrix}$$

  $A$         $x$        $b$  

- Example 1: image reconstruction

- The domain of an image is its pixels (say  $n$  pixels)

- $f_j$ : indicator of the  $j$ th pixel;

$$f = \sum_{j=1}^n x_j f_j$$

- Linear measurements

$$R_i f_j = a_{ij}, \quad R_i f = b_i, \quad i = 1, \dots, k$$

- Goal: to recover  $x_1, \dots, x_n$  from the measurements

# Linear Inverse Problem

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{k1} & a_{k2} & \cdots & a_{kn} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_k \end{bmatrix}$$

✓  $A$       🔍  $x$        $b$  ✓

- ◆ Example 1: image reconstruction

- ◆ The domain of an image is its pixels (say  $n$  pixels)
- ◆  $f_j$ : indicator of the  $j$ th pixel;

$$f = \sum_{j=1}^n x_j f_j$$

- ◆ Linear measurements  
 $R_i f_j = a_{ij}, R_i f = b_i, i = 1, \dots, k$
- ◆ Goal: to recover  $x_1, \dots, x_n$  from the measurements

- ◆ Example 2: learning the frequency of alphabets from a corpus

# Linear Inverse Problem

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{k1} & a_{k2} & \cdots & a_{kn} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_k \end{bmatrix}$$

✓  $A$       🔍  $x$        $b$  ✓

- Example 1: image reconstruction

- The domain of an image is its pixels (say  $n$  pixels)
- $f_j$ : indicator of the  $j$ th pixel;

$$f = \sum_{j=1}^n x_j f_j$$

- Linear measurements

$$R_i f_j = a_{ij}, \quad R_i f = b_i, \quad i = 1, \dots, k$$

- Goal: to recover  $x_1, \dots, x_n$  from the measurements

- Example 2: learning the frequency of alphabets from a corpus

- $n$  alphabets  $a_1, \dots, a_n$  occurring with **unknown probabilities**  $x_1, \dots, x_n$

# Linear Inverse Problem

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{k1} & a_{k2} & \cdots & a_{kn} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_k \end{bmatrix}$$

✓  $A$       🔍  $x$        $b$  ✓

- Example 1: image reconstruction

- The domain of an image is its pixels (say  $n$  pixels)

- $f_j$ : indicator of the  $j$ th pixel;

$$f = \sum_{j=1}^n x_j f_j$$

- Linear measurements

$$R_i f_j = a_{ij}, \quad R_i f = b_i, \quad i = 1, \dots, k$$

- Goal: to recover  $x_1, \dots, x_n$  from the measurements

- Example 2: learning the frequency of alphabets from a corpus

- $n$  alphabets  $a_1, \dots, a_n$  occurring with **unknown probabilities**  $x_1, \dots, x_n$

- Corpus: body of text containing strings of alphabets

# Linear Inverse Problem

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{k1} & a_{k2} & \cdots & a_{kn} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_k \end{bmatrix}$$

✓  $A$       🔍  $x$        $b$  ✓

- Example 1: image reconstruction

- The domain of an image is its pixels (say  $n$  pixels)
- $f_j$ : indicator of the  $j$ th pixel;

$$f = \sum_{j=1}^n x_j f_j$$

- Linear measurements  
 $R_i f_j = a_{ij}, R_i f = b_i, i = 1, \dots, k$
- Goal: to recover  $x_1, \dots, x_n$  from the measurements

- Example 2: learning the frequency of alphabets from a corpus

- $n$  alphabets  $a_1, \dots, a_n$  occurring with **unknown probabilities**  $x_1, \dots, x_n$
- Corpus: body of text containing strings of alphabets
- Linear measurements on the corpus:  
 $\mathbb{E}[T_i(Z)] = b_i, i = 1, \dots, k$

# Linear Inverse Problem

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{k1} & a_{k2} & \cdots & a_{kn} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_k \end{bmatrix}$$

✓  $A$       🔍  $x$        $b$  ✓

- Example 1: image reconstruction

- The domain of an image is its pixels (say  $n$  pixels)
- $f_j$ : indicator of the  $j$ th pixel;

$$f = \sum_{j=1}^n x_j f_j$$

- Linear measurements  
 $R_i f_j = a_{ij}, R_i f = b_i, i = 1, \dots, k$
- Goal: to recover  $x_1, \dots, x_n$  from the measurements

- Example 2: learning the frequency of alphabets from a corpus

- $n$  alphabets  $a_1, \dots, a_n$  occurring with **unknown probabilities**  $x_1, \dots, x_n$
- Corpus: body of text containing strings of alphabets
- Linear measurements on the corpus:  
 $\mathbb{E}[T_i(Z)] = b_i, i = 1, \dots, k$
- Goal: to recover  $x_1, \dots, x_n$  satisfying the moment constraints

# Linear Inverse Problem

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{k1} & a_{k2} & \cdots & a_{kn} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_k \end{bmatrix}$$

✓  $A$       🔍  $x$        $b$  ✓

- Example 1: image reconstruction

- The domain of an image is its pixels (say  $n$  pixels)
- $f_j$ : indicator of the  $j$ th pixel;  

$$f = \sum_{j=1}^n x_j f_j$$
- Linear measurements  
 $R_i f_j = a_{ij}, R_i f = b_i, i = 1, \dots, k$
- Goal: to recover  $x_1, \dots, x_n$  from the measurements

Is there a principled  
approach to solve the  
problem?

- Example 2: learning the frequency of alphabets from a corpus

- $n$  alphabets  $a_1, \dots, a_n$  occurring with **unknown probabilities**  $x_1, \dots, x_n$
- Corpus: body of text containing strings of alphabets
- Linear measurements on the corpus:  
 $\mathbb{E}[T_i(Z)] = b_i, i = 1, \dots, k$
- Goal: to recover  $x_1, \dots, x_n$  satisfying the moment constraints

# Approach 1: Function-Minimisation

# Approach 1: Function-Minimisation


- Define a “cost” function  $F$

# Approach 1: Function-Minimisation

- Define a “cost” function  $F$
- Recover  $x$  as a minimiser of  $F$


# Approach 1: Function-Minimisation

- Define a “cost” function  $F$
- Recover  $x$  as a minimiser of  $F$




$$x = \arg \min_{w \in L} F(w)$$

# Solution using Optimisation Techniques




$$\min_{w \in L} F(w)$$




$$\min_{w \in L} F(w)$$

# Solution using Optimisation Techniques




$$\min_{w \in L} F(w)$$

$$L = \{w \in \mathbb{R}^n : a^T w = b\}$$



$$\min_{w \in L} F(w)$$


# Solution using Optimisation Techniques



$$\min_{w \in L} F(w)$$


$$L = \{w \in \mathbb{R}^n : a^T w = b\}$$

$$G(w, \lambda) = F(w) - \lambda(a^T w - b)$$



$$\min_{w \in L} F(w)$$

# Solution using Optimisation Techniques




$$\min_{w \in L} F(w)$$

$$L = \{w \in \mathbb{R}^n : a^T w = b\}$$


$$G(w, \lambda) = F(w) - \lambda(a^T w - b)$$

$$\nabla_w G(w, \lambda) = 0 \iff \nabla F(w) = \lambda a$$



$$\min_{w \in L} F(w)$$

# Solution using Optimisation Techniques




$$\min_{w \in L} F(w)$$

$$L = \{w \in \mathbb{R}^n : a^T w = b\}$$


$$G(w, \lambda) = F(w) - \lambda(a^T w - b)$$

$$\nabla_w G(w, \lambda) = 0 \iff \boxed{\nabla F(w) = \lambda a}$$



$$\min_{w \in L} F(w)$$

# Solution using Optimisation Techniques




$$\min_{w \in L} F(w)$$

$$L = \{w \in \mathbb{R}^n : a^T w = b\}$$

$$G(w, \lambda) = F(w) - \lambda(a^T w - b)$$


$$\nabla_w G(w, \lambda) = 0 \iff \boxed{\nabla F(w) = \lambda a}$$



$$\min_{w \in L} F(w)$$

$$L = \{w \in \mathbb{R}^n : Aw = b\}$$

# Solution using Optimisation Techniques




$$\min_{w \in L} F(w)$$

$$L = \{w \in \mathbb{R}^n : a^T w = b\}$$

$$G(w, \lambda) = F(w) - \lambda(a^T w - b)$$

$$\nabla_w G(w, \lambda) = 0 \iff \boxed{\nabla F(w) = \lambda a}$$



$$\min_{w \in L} F(w)$$

$$L = \{w \in \mathbb{R}^n : Aw = b\}$$

$$G(w, \lambda) = F(w) - \lambda^T(Aw - b)$$

$$\nabla_w G(w, \lambda) = 0 \iff \nabla F(w) = A^T \lambda$$

# Commonly Used $F$

# Commonly Used $F$

- Least squares:  $F(w) = \|w\|^2$

# Commonly Used $F$

- Least squares:  $F(w) = \|w\|^2$

$$x = \arg \min_{w \in L} \|w\|^2 = \arg \min_{w \in L} \sum_{i=1}^n w_i^2$$

# Commonly Used $F$

- Least squares:  $F(w) = \|w\|^2$

$$x = \arg \min_{w \in L} \|w\|^2 = \arg \min_{w \in L} \sum_{i=1}^n w_i^2$$

- Maximum entropy:  $F(w) = \sum_i w_i \log w_i$

# Commonly Used $F$

- Least squares:  $F(w) = \|w\|^2$

$$x = \arg \min_{w \in L} \|w\|^2 = \arg \min_{w \in L} \sum_{i=1}^n w_i^2$$

- Maximum entropy:  $F(w) = \sum_i w_i \log w_i$

$$x = \arg \min_{w \in L \cap \mathbb{R}_+^n} \sum_{i=1}^n w_i \log w_i$$

## Commonly Used $F$ with Prior Guess

- Prior guess:  $x^*$

## Commonly Used $F$ with Prior Guess

- Prior guess:  $x^\star$
- Least squares:  $F(w | x^\star) = \|w - x^\star\|^2$

## Commonly Used $F$ with Prior Guess

- Prior guess:  $x^\star$
- Least squares:  $F(w | x^\star) = \|w - x^\star\|^2$

$$x = \arg \min_{w \in L} \|w - x^\star\|^2$$

## Commonly Used $F$ with Prior Guess

- Prior guess:  $x^\star$
- Least squares:  $F(w|x^\star) = \|w - x^\star\|^2$

$$x = \arg \min_{w \in L} \|w - x^\star\|^2$$

- Maximum entropy:  $F(w|x^\star) = \sum_i w_i \log \frac{w_i}{x_i^\star}$

## Commonly Used $F$ with Prior Guess

- Prior guess:  $x^\star$

- Least squares:  $F(w | x^\star) = \|w - x^\star\|^2$





$$x = \arg \min_{w \in L} \|w - x^\star\|^2$$

- Maximum entropy:  $F(w | x^\star) = \sum_i w_i \log \frac{w_i}{x_i^\star}$



$$x = \arg \min_{w \in L \cap \mathbb{R}_+^n} \sum_i w_i \log \frac{w_i}{x_i^\star}$$

# Projection Rules



# Projection Rules





# Projection Rules




Function-minimisation  
=  
projection rule




# Projection Rules



Function-minimisation  
=  
projection rule



$$\Pi : (L, x^\star) \mapsto \Pi(L | x^\star) \in L$$



# Projection Rules and Function Minimisation

# Projection Rules and Function Minimisation

$$\begin{aligned} F : \mathbb{R}^n \times \mathbb{R}^n &\rightarrow \mathbb{R} \\ (x^\star, w) &\mapsto F(w \mid x^\star) \end{aligned}$$

# Projection Rules and Function Minimisation

$$F : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$$

$$(x^\star, w) \mapsto F(w | x^\star)$$

$\Pi : \mathcal{L} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  projection rule

$$\Pi : (L, x^\star) \mapsto \Pi(L | x^\star) \in L$$

# Projection Rules and Function Minimisation

$$F : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$$

$$(x^\star, w) \mapsto F(w | x^\star)$$

$\Pi : \mathcal{L} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  projection rule

$$\Pi : (L, x^\star) \mapsto \Pi(L | x^\star) \in L$$

Definition:  $\Pi$  is generated by  $F$ , if

$$\Pi(L | x^\star) = \arg \min_{w \in L} F(w | x^\star)$$

# Projection Rules and Function Minimisation

$$F : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$$

$$(x^\star, w) \mapsto F(w | x^\star)$$


$\Pi : \mathcal{L} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  projection rule

$$\Pi : (L, x^\star) \mapsto \Pi(L | x^\star) \in L$$


Definition:

$\Pi$  is generated by  $F$ , if

$$\Pi(L | x^\star) = \arg \min_{w \in L} F(w | x^\star)$$



$$F(w | x^\star) = \|w - x^\star\|^2$$



$$F(w | x^\star) = \sum_i w_i \log \frac{w_i}{x_i^\star}$$


Is every projection rule generated by some function?

Is every projection rule generated by some function?

- ◆ Answers to this question were provided by Csiszár in 1991

# Is every projection rule generated by some function?

- ◆ Answers to this question were provided by Csiszár in 1991



*The Annals of Statistics*  
1991, Vol. 19, No. 4, 2032–2066

## WHY LEAST SQUARES AND MAXIMUM ENTROPY? AN AXIOMATIC APPROACH TO INFERENCE FOR LINEAR INVERSE PROBLEMS<sup>1</sup>

BY IMRE CSISZÁR

*Mathematical Institute of the Hungarian Academy of Sciences*

An attempt is made to determine the logically consistent rules for selecting a vector from any feasible set defined by linear constraints, when either all  $n$ -vectors or those with positive components or the probability vectors are permissible. Some basic postulates are satisfied if and only if the selection rule is to minimize a certain function which, if a “prior guess” is available, is a measure of distance from the prior guess. Two further natural postulates restrict the permissible distances to the author’s  $f$ -divergences and Bregman’s divergences, respectively. As corollaries, axiomatic characterizations of the methods of least squares and minimum discrimination information are arrived at. Alternatively, the latter are also characterized by a postulate of composition consistency. As a special case, a derivation of the method of maximum entropy from a small set of natural axioms is obtained.

The seminal work of Csiszár

# Approach 2: Axioms

# Approach 2: Axioms

- ◆ Csiszár demonstrated that if projection rules satisfy some naturally appealing axioms, they must be generated by “nice” functions

# Approach 2: Axioms

- ◆ Csiszár demonstrated that if projection rules satisfy some naturally appealing axioms, they must be generated by “nice” functions

Projection rules satisfies the axioms of

**Regularity** + Locality

**Regularity** + Locality + Subspace Transitivity

**Regularity** + Locality + Subspace Transitivity + Statistical

**Regularity** + Locality + Subspace Transitivity + Location Invariance + Scale Invariance

Projection rule is generated by

$$F(w|x^\star) = \sum_{i=1}^n f_i(w_i|x_i^\star), f_i \text{ continuously differentiable and strictly convex}$$

$F(w|x^\star)$  = Bregman's divergence

$$F(w|x^\star) = \sum_{i=1}^n w_i \log \frac{w_i}{x_i^\star}$$

$$F(w|x^\star) = \sum_{i=1}^n (w_i - x_i^\star)^2$$

# Why are these axioms of interest to us?

- ◆ Kumar and Sundaresan ('15) discovered a family of projection rules satisfying

Regularity 

Subspace transitivity 

Locality 

- ◆ These projection rules are generated by a parametric divergence known as relative  $\alpha$ -entropy  
(related to the well-known **Rényi divergence**)

M. A. Kumar and R. Sundaresan. (2015). Minimization problems based on relative  $\alpha$ -entropy I: Forward projection.  
IEEE Transactions on Information Theory, 61(9), 5063-5080.

M. A. Kumar and R. Sundaresan. (2015). Minimization problems based on relative  $\alpha$ -entropy II: Reverse projection.  
IEEE Transactions on Information Theory, 61(9), 5081-5095.

“Are the projection rules generated by **relative  $\alpha$ -entropy** the only rules which are **regular, subspace transitive and nonlocal?**”

Regularity



Subspace transitivity




Locality




# Our Quest

- ◆ Csiszár's results provide a necessary and sufficient axiomatic characterisation of many projection rules




# Our Quest

- ◆ Csiszár's results provide a necessary and sufficient axiomatic characterisation of many projection rules



# Our Quest

- ◆ Csiszár's results provide a necessary and sufficient axiomatic characterisation of many projection rules



# The axiom of regularity

# The axiom of regularity

$\Pi : \mathcal{L} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  projection rule

# The axiom of regularity

$\Pi : \mathcal{L} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  projection rule

$$\mathcal{M} = \left\{ \{w \in \mathbb{R}^n : a^T w = b\}, a \in \mathbb{R}^n, a \neq 0, b \in \mathbb{R} \right\}$$

# The axiom of regularity

$\Pi : \mathcal{L} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  projection rule

$$\mathcal{M} = \left\{ \{w \in \mathbb{R}^n : a^T w = b\}, a \in \mathbb{R}^n, a \neq 0, b \in \mathbb{R} \right\}$$

$\Pi$  satisfies regularity if, for all  $x^\star \in \mathbb{R}^n$ ,

(Consistency)

$$L' \subset L, \Pi(L | x^\star) \in L' \implies \Pi(L' | x^\star) = \Pi(L | x^\star)$$

(Distinctness)



$$L, \tilde{L} \in \mathcal{M}, L \neq \tilde{L}, x^\star \notin L \cap \tilde{L} \implies \Pi(L | x^\star) \neq \Pi(\tilde{L} | x^\star)$$

(Continuity)

$\Pi(\cdot | x^\star)$  restricted to any fixed dimension is continuous



Regularity

# The axiom of consistency



$$L' \subset L, \Pi(L | x^*) \in L' \implies \Pi(L' | x^*) = \Pi(L | x^*)$$


# The axiom of consistency



$$L' \subset L, \Pi(L | x^*) \in L' \implies \Pi(L' | x^*) = \Pi(L | x^*)$$

**Additional observations do not give a reason to change the original selection**

# The axiom of distinctness




$$\mathcal{M} = \left\{ \{w \in \mathbb{R}^n : a^T w = b\}, a \in \mathbb{R}^n, a \neq 0, b \in \mathbb{R} \right\}$$

$$L, \tilde{L} \in \mathcal{M}, L \neq \tilde{L}, x^* \notin L \cap \tilde{L} \implies \Pi(L | x^*) \neq \Pi(\tilde{L} | x^*)$$


# The axiom of continuity




# The axiom of continuity



# The axiom of continuity



# Axiom of subspace transitivity



$$L' \subset L \implies \Pi(L' | x^*) = \Pi(\Pi(L | x^*) | x^*)$$

# A Key Implication of Regularity

$$\mathcal{L}^0(x^\star) = \left\{ L \in \mathcal{L} : \Pi(L|x^\star) = x^\star \right\}, \quad x^\star \in \mathbb{R}^n$$

$$\mathcal{M} = \left\{ \{w \in \mathbb{R}^n : a^T w = b\}, a \in \mathbb{R}^n, a \neq 0, b \in \mathbb{R} \right\}$$

# A Key Implication of Regularity

$$\mathcal{L}^0(x^\star) = \left\{ L \in \mathcal{L} : \Pi(L|x^\star) = x^\star \right\}, \quad x^\star \in \mathbb{R}^n$$

$$\mathcal{M} = \left\{ \{w \in \mathbb{R}^n : a^T w = b\}, a \in \mathbb{R}^n, a \neq 0, b \in \mathbb{R} \right\}$$


$\Pi(\cdot|x^\star) : \mathcal{M} \setminus \mathcal{L}^0(x^\star) \rightarrow \mathbb{R}^n \setminus \{x^\star\}$  is a homeomorphism

# A Key Implication of Regularity

$$\mathcal{L}^0(x^\star) = \left\{ L \in \mathcal{L} : \Pi(L|x^\star) = x^\star \right\}, \quad x^\star \in \mathbb{R}^n$$

$$\mathcal{M} = \left\{ \{w \in \mathbb{R}^n : a^T w = b\}, a \in \mathbb{R}^n, a \neq 0, b \in \mathbb{R} \right\}$$


$\Pi(\cdot | x^\star) : \mathcal{M} \setminus \mathcal{L}^0(x^\star) \rightarrow \mathbb{R}^n \setminus \{x^\star\}$  is a homeomorphism



# A Key Implication of Regularity

$\Pi(\cdot | x^*) : \mathcal{M} \setminus \mathcal{L}^0(x^*) \rightarrow \mathbb{R}^n \setminus \{x^*\}$  is a homeomorphism



This means that for every  $w \neq x^*$ ,  
there exists a unique set  $L$  of dimension  $(n - 1)$   
such that  $\Pi(L | x^*) = w$ . Denote this  $L$  as  $L(w | x^*)$



# A Key Implication of Regularity

$\Pi(\cdot | x^*) : \mathcal{M} \setminus \mathcal{L}^0(x^*) \rightarrow \mathbb{R}^n \setminus \{x^*\}$  is a homeomorphism

This means that for every  $w \neq x^*$ ,  
there exists a unique set  $L$  of dimension  $(n - 1)$   
such that  $\Pi(L | x^*) = w$ . Denote this  $L$  as  $L(w | x^*)$





$$\begin{aligned}
 L(w | x^*) &= \left\{ y \in \mathbb{R}^n : \underbrace{\mathbf{a}^\top}_{\parallel \mathbf{a} \parallel} y = b \right\} \\
 &= \left\{ y \in \mathbb{R}^n : \mathbf{a}^\top y = \mathbf{a}^\top w \right\} \\
 &= \left\{ y \in \mathbb{R}^n : \underbrace{\frac{\mathbf{a}^\top}{\parallel \mathbf{a} \parallel}}_{\tilde{\mathbf{a}}(w | x^*)} (y - w) = 0 \right\} \\
 &= \left\{ y \in \mathbb{R}^n : \tilde{\mathbf{a}}(w | x^*)^\top (y - w) = 0 \right\}
 \end{aligned}$$

# A Key Implication of Regularity

$\Pi(\cdot | x^*) : \mathcal{M} \setminus \mathcal{L}^0(x^*) \rightarrow \mathbb{R}^n \setminus \{x^*\}$  is a homeomorphism

This means that for every  $w \neq x^*$ , there exists a unique set  $L$  of dimension  $(n - 1)$  such that  $\Pi(L | x^*) = w$ . Denote this  $L$  as  $L(w | x^*)$



$$\begin{aligned} L(w | x^*) &= \left\{ y \in \mathbb{R}^n : \underbrace{\underline{a}^\top}_{\parallel a \parallel} y = b \right\} \\ &= \left\{ y \in \mathbb{R}^n : \underline{a}^\top y = \underline{a}^\top w \right\} \\ &= \left\{ y \in \mathbb{R}^n : \underbrace{\frac{\underline{a}^\top}{\parallel a \parallel}}_{\parallel a \parallel} (y - w) = 0 \right\} \\ &= \left\{ y \in \mathbb{R}^n : \underline{\tilde{a}(w | x^*)}^\top (y - w) = 0 \right\} \end{aligned}$$

$w \mapsto \tilde{a}(w | x^*)$  is continuous

# Regularity and vector fields

- ◆ Mere regularity helps us extract a **vector field**

# Regularity and vector fields

- ◆ Mere regularity helps us extract a **vector field**

$$\tilde{a}(\cdot | x^\star) : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

$\tilde{a}(\cdot | x^\star)$  continuous

$$\tilde{a}(x^\star | x^\star) = 0,$$

$$\tilde{a}(w | x^\star) \neq 0 \text{ for all } w \neq x^\star$$


# Existence of a Non-Zero Scaling Function

$$\tilde{a}(\cdot | x^*) : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

$\tilde{a}(\cdot | x^*)$  continuous

$$\tilde{a}(x^* | x^*) = 0,$$

$$\tilde{a}(w | x^*) \neq 0 \text{ for all } w \neq x^*$$



$$\min_{w \in L} F(w)$$

$$L = \{w \in \mathbb{R}^n : a^T w = b\}$$

$$G(w, \lambda) = F(w) - \lambda(a^T w - b)$$

$$\nabla_w G(w, \lambda) = 0 \iff \boxed{\nabla F(w) = \lambda a}$$


# Existence of a Non-Zero Scaling Function

$$\tilde{a}(\cdot | x^\star) : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

$\tilde{a}(\cdot | x^\star)$  continuous

$$\tilde{a}(x^\star | x^\star) = 0,$$

$$\tilde{a}(w | x^\star) \neq 0 \text{ for all } w \neq x^\star$$



$$\min_{w \in L} F(w)$$

$$L = \{w \in \mathbb{R}^n : a^T w = b\}$$

$$G(w, \lambda) = F(w) - \lambda(a^T w - b)$$

$$\nabla_w G(w, \lambda) = 0 \iff \boxed{\nabla F(w) = \lambda a}$$

Is there  $\lambda : \mathbb{R}^n \rightarrow \mathbb{R}$  such that  
 $\lambda(w) \cdot \tilde{a}(w | x^\star) = \nabla F(w | x^\star)$ ?

# Conservative Vector Fields and Regularity

# Conservative Vector Fields and Regularity

$\Pi(\cdot | x^\star)$  regular

# Conservative Vector Fields and Regularity

$\Pi(\cdot | x^\star)$  regular 


# Conservative Vector Fields and Regularity

$\Pi(\cdot | x^\star)$  regular




$\tilde{a}(\cdot | x^\star)$  continuous vector field


# Conservative Vector Fields and Regularity




# Conservative Vector Fields and Regularity




# Conservative Vector Fields and Regularity




# Conservative Vector Fields and Regularity



# Conservative Vector Fields and Regularity



# Conservative Vector Fields and Regularity



# An Open Question

# An Open Question

Given a continuous vector field, does there exist a scaling function such that the **scaled vector field is conservative**?

# An Open Question

Given a continuous vector field, does there exist a scaling function such that the **scaled vector field is conservative**?

$$a : \mathbb{R}^n \rightarrow \mathbb{R}^n \text{ continuous}$$

$$\lambda : \mathbb{R}^n \rightarrow \mathbb{R}$$

$$\lambda \cdot a = \nabla F?$$

## Instances Where the Answers are Known / Not Unknown

Given a continuous vector field, does there exist a scaling function such that the **scaled vector field is conservative**?

$$a : \mathbb{R}^n \rightarrow \mathbb{R}^n \text{ continuous}$$

$$\lambda : \mathbb{R}^n \rightarrow \mathbb{R}$$

$$\lambda \cdot a = \nabla F?$$

## Instances Where the Answers are Known / Not Unknown

$$a(w) = \frac{w}{\|w\|} \quad \lambda(w) = \|w\|$$

Given a continuous vector field, does there exist a scaling function such that the **scaled vector field is conservative**?

$$a : \mathbb{R}^n \rightarrow \mathbb{R}^n \text{ continuous}$$

$$\lambda : \mathbb{R}^n \rightarrow \mathbb{R}$$

$$\lambda \cdot a = \nabla F?$$

## Instances Where the Answers are Known / Not Unknown

$$a(w) = \frac{w}{\|w\|} \quad \lambda(w) = \|w\|$$

Given a continuous vector field, does there exist a scaling function such that the **scaled vector field is conservative**?

$$\lambda(w) \ a(w) = \nabla F(w) \quad F(w) = \frac{1}{2} \|w\|^2$$

$$a : \mathbb{R}^n \rightarrow \mathbb{R}^n \text{ continuous}$$

$$\lambda : \mathbb{R}^n \rightarrow \mathbb{R}$$

$$\lambda \cdot a = \nabla F?$$

## Instances Where the Answers are Known / Not Unknown

$$a(w) = \frac{w}{\|w\|} \quad \lambda(w) = \|w\|$$

Given a continuous vector field, does there exist a scaling function such that the **scaled vector field is conservative**?

$$\lambda(w) \ a(w) = \nabla F(w) \quad F(w) = \frac{1}{2}\|w\|^2$$

$$a : \mathbb{R}^n \rightarrow \mathbb{R}^n \text{ continuous}$$

$$\lambda : \mathbb{R}^n \rightarrow \mathbb{R}$$

$$\lambda \cdot a = \nabla F?$$

$$a(x, y, z) = \begin{pmatrix} y \\ 0 \\ 1 \end{pmatrix} \implies \text{there is no } \lambda \text{ such that } \lambda \cdot a = \nabla F$$

## Instances Where the Answers are Known / Not Unknown

$$a(w) = \frac{w}{\|w\|} \quad \lambda(w) = \|w\|$$

Given a continuous vector field, does there exist a scaling function such that the **scaled vector field is conservative**?

$$\lambda(w) \ a(w) = \nabla F(w) \quad F(w) = \frac{1}{2}\|w\|^2$$

$$a : \mathbb{R}^n \rightarrow \mathbb{R}^n \text{ continuous}$$

$$\lambda : \mathbb{R}^n \rightarrow \mathbb{R}$$

$$\lambda \cdot a = \nabla F?$$

$$a(x, y, z) = \begin{pmatrix} y \\ 0 \\ 1 \end{pmatrix} \Rightarrow \text{there is no } \lambda \text{ such that } \lambda \cdot a = \nabla F$$

Can this arise from a regular projection rule?

# Summary

- ◆ Two approaches for solving systems of linear equations:

- ◆ Function-minimisation approach

Projection rules satisfies the axioms of

Regularity + Locality

$$F(w|x^\star) = \sum_{i=1}^n f_i(w_i|x_i^\star), f_i \text{ continuously differentiable and strictly convex}$$

Regularity + Locality + Subspace Transitivity

$$F(w|x^\star) = \text{Bregman's divergence}$$

Regularity + Locality + Subspace Transitivity + Statistical

$$F(w|x^\star) = \sum_{i=1}^n w_i \log \frac{w_i}{x_i^\star}$$

Regularity + Locality + Subspace Transitivity + Location Invariance + Scale Invariance

$$F(w|x^\star) = \sum_{i=1}^n (w_i - x_i^\star)^2$$

- ◆ Axiomatic approach

- ◆ Regularity: a fundamental axiom

- ◆ Regularity has connections with conservative vector fields

- ◆ Given a continuous vector field that is not necessarily conservative, is there a continuous scaling that can make the product vector field conservative? Open!

# Thank you!

[karthik@nus.edu.sg](mailto:karthik@nus.edu.sg)