



Stochastic Processes

Application of Metropolis–Hastings to Bayesian Logistic Regression,
Test for Irreducibility, Sample Paths of a Random Process in
Simulations

Karthik P. N.

Assistant Professor, Department of AI

Email: pnkarthik@ai.iith.ac.in

29 April 2025



Bayesian Logistic Regression using Challenger O-Ring Dataset

- Between 1981 and 1985, NASA launched 23 space-shuttle missions before the ill-fated Challenger flight on 28 Jan 1986



Bayesian Logistic Regression using Challenger O-Ring Dataset

- Between 1981 and 1985, NASA launched 23 space-shuttle missions before the ill-fated Challenger flight on 28 Jan 1986
- For every flight $i \in \{1, \dots, 23\}$, engineers recorded

Bayesian Logistic Regression using Challenger O-Ring Dataset

- Between 1981 and 1985, NASA launched 23 space-shuttle missions before the ill-fated Challenger flight on 28 Jan 1986
- For every flight $i \in \{1, \dots, 23\}$, engineers recorded
 - the ambient air-temperature at lift-off x_i (in °F), and

Bayesian Logistic Regression using Challenger O-Ring Dataset


- Between 1981 and 1985, NASA launched 23 space-shuttle missions before the ill-fated Challenger flight on 28 Jan 1986
- For every flight $i \in \{1, \dots, 23\}$, engineers recorded
 - the ambient air-temperature at lift-off x_i (in °F), and
 - $y_i = \mathbf{1}\{\text{whether at least one solid-rocket-booster O-ring was damaged during ascent}\}$

Bayesian Logistic Regression using Challenger O-Ring Dataset

- Between 1981 and 1985, NASA launched 23 space-shuttle missions before the ill-fated Challenger flight on 28 Jan 1986
- For every flight $i \in \{1, \dots, 23\}$, engineers recorded
 - the ambient air-temperature at lift-off x_i (in °F), and
 - $y_i = \mathbf{1}\{\text{whether at least one solid-rocket-booster O-ring was damaged during ascent}\}$
- The lowest temperature available in the dataset was 53°F, whereas Challenger launched at 31°F despite warnings from engineers

Observed Empirical Relationship


- Empirical relationship between probability of ring damage and temperature was observed to be as shown below.



Regression Model

- For data point i , let

$$\eta_i = \beta_0 + \beta_1 x_i.$$




Regression Model

- For data point i , let


$$\eta_i = \beta_0 + \beta_1 x_i.$$

- Taking $\beta = (\beta_0, \beta_1)$, let

$$\pi_i = \text{Logit}(\eta_i) = \frac{\exp \eta_i}{1 + \exp \eta_i}.$$



Regression Model



- For data point i , let

$$\eta_i = \beta_0 + \beta_1 x_i.$$


- Taking $\beta = (\beta_0, \beta_1)$, let

$$\pi_i = \text{Logit}(\eta_i) = \frac{\exp \eta_i}{1 + \exp \eta_i}.$$

- Conditional distribution of one output data point:

$$\mathbb{P}(y_i = 1 \mid x_i, \beta) = \pi_i = 1 - \mathbb{P}(y_i = 0 \mid x_i, \beta).$$

Regression Model



- For data point i , let

$$\eta_i = \beta_0 + \beta_1 x_i.$$

- Taking $\beta = (\beta_0, \beta_1)$, let

$$\pi_i = \text{Logit}(\eta_i) = \frac{\exp \eta_i}{1 + \exp \eta_i}.$$

- Conditional distribution of one output data point:

$$\mathbb{P}(y_i = 1 \mid x_i, \beta) = \pi_i = 1 - \mathbb{P}(y_i = 0 \mid x_i, \beta).$$

- Data points $\{(x_i, y_i)\}_{i=1}^{23}$ are IID

Prior, Likelihood, and Posterior

- $\beta = (\beta_0, \beta_1)$ unknown. A reasonable prior is

$$\beta_0, \beta_1 \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, \sigma^2), \quad g(\beta) = \frac{1}{2\pi\sigma^2} \exp\left(-\frac{\beta_0^2 + \beta_1^2}{2\sigma^2}\right).$$

Prior, Likelihood, and Posterior

- $\beta = (\beta_0, \beta_1)$ unknown. A reasonable prior is

$$\beta_0, \beta_1 \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, \sigma^2), \quad g(\beta) = \frac{1}{2\pi\sigma^2} \exp\left(-\frac{\beta_0^2 + \beta_1^2}{2\sigma^2}\right).$$

- Likelihood function is given by

$$p(\mathbf{y} \mid \mathbf{x}, \beta) = \prod_{i=1}^{23} p(y_i \mid x_i, \beta) = \prod_{i=1}^{23} \pi_i^{y_i} (1 - \pi_i)^{1-y_i}.$$

Prior, Likelihood, and Posterior

- $\beta = (\beta_0, \beta_1)$ unknown. A reasonable prior is

$$\beta_0, \beta_1 \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, \sigma^2), \quad g(\beta) = \frac{1}{2\pi\sigma^2} \exp\left(-\frac{\beta_0^2 + \beta_1^2}{2\sigma^2}\right).$$

- Likelihood function is given by

$$p(\mathbf{y} \mid \mathbf{x}, \beta) = \prod_{i=1}^{23} p(y_i \mid x_i, \beta) = \prod_{i=1}^{23} \pi_i^{y_i} (1 - \pi_i)^{1-y_i}.$$

- Posterior function is given by

$$f(\beta \mid \mathbf{x}, \mathbf{y}) = \frac{p(\mathbf{y} \mid \mathbf{x}, \beta) g(\beta)}{c}, \quad c = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} p(\mathbf{y} \mid \mathbf{x}, \beta') g(\beta') d\beta'_0 d\beta'_1.$$

Prior, Likelihood, and Posterior

- $\beta = (\beta_0, \beta_1)$ unknown. A reasonable prior is

$$\beta_0, \beta_1 \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, \sigma^2), \quad g(\beta) = \frac{1}{2\pi\sigma^2} \exp\left(-\frac{\beta_0^2 + \beta_1^2}{2\sigma^2}\right).$$

- Likelihood function is given by

$$p(\mathbf{y} \mid \mathbf{x}, \beta) = \prod_{i=1}^{23} p(y_i \mid x_i, \beta) = \prod_{i=1}^{23} \pi_i^{y_i} (1 - \pi_i)^{1-y_i}.$$

- Posterior function is given by

$$f(\beta \mid \mathbf{x}, \mathbf{y}) = \frac{p(\mathbf{y} \mid \mathbf{x}, \beta) g(\beta)}{\mathcal{C}}, \quad \mathcal{C} = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} p(\mathbf{y} \mid \mathbf{x}, \beta') g(\beta') d\beta'_0 d\beta'_1.$$

- **No closed-form expression for \mathcal{C}**

Setting up Metropolis-Hastings

- Set Q matrix (kernel) as

$$Q(\beta, \cdot) = \mathcal{N}(\beta, \Sigma),$$

for some fixed covariance matrix Σ

Setting up Metropolis-Hastings

- Set Q matrix (kernel) as

$$Q(\beta, \cdot) = \mathcal{N}(\beta, \Sigma),$$

for some fixed covariance matrix Σ

- Q is symmetric

Setting up Metropolis-Hastings

- Set Q matrix (kernel) as

$$Q(\beta, \cdot) = \mathcal{N}(\beta, \Sigma),$$

for some fixed covariance matrix Σ

- Q is symmetric
- The acceptance probabilities are then given by

$$A_{\beta, \beta'} = \min \left\{ 1, \frac{f(\beta' | \mathbf{x}, \mathbf{y}) \cdot Q(\beta', \beta)}{f(\beta | \mathbf{x}, \mathbf{y}) \cdot Q(\beta, \beta')} \right\} = \min \left\{ 1, \frac{p(\mathbf{y} | \mathbf{x}, \beta') \cdot g(\beta')}{p(\mathbf{y} | \mathbf{x}, \beta) \cdot g(\beta)} \right\}.$$

Setting up Metropolis-Hastings

- Set Q matrix (kernel) as

$$Q(\beta, \cdot) = \mathcal{N}(\beta, \Sigma),$$

for some fixed covariance matrix Σ

- Q is symmetric
- The acceptance probabilities are then given by

$$A_{\beta, \beta'} = \min \left\{ 1, \frac{f(\beta' | \mathbf{x}, \mathbf{y}) \cdot Q(\beta', \beta)}{f(\beta | \mathbf{x}, \mathbf{y}) \cdot Q(\beta, \beta')} \right\} = \min \left\{ 1, \frac{p(\mathbf{y} | \mathbf{x}, \beta') \cdot g(\beta')}{p(\mathbf{y} | \mathbf{x}, \beta) \cdot g(\beta)} \right\}.$$

- For any bounded, continuous function h , we have by the ergodic theorem that

$$\frac{1}{n} \sum_{k=1}^n h(X_k) \xrightarrow{\text{a.s.}} \int h(\beta) f(\beta | \mathbf{x}, \mathbf{y}) \, d\beta,$$

where $\{X_k\}_{k=0}^{\infty}$ is a DTMC with stationary kernel $f(\cdot | \mathbf{x}, \mathbf{y})$

Other Tidbits

Test for Irreducibility

Theorem (Test for Irreducibility)

A row stochastic matrix P of size $d \times d$ is irreducible if and only if

$$\sum_{k=0}^{d-1} P^k > \mathbf{0},$$

where $P^0 = I$, $\mathbf{0}$ denotes the all-zeros matrix of size $d \times d$, and the inequality is entry-wise.

A simple algorithm to check for irreducibility:

- Start with $M_0 = I$.

Test for Irreducibility

Theorem (Test for Irreducibility)

A row stochastic matrix P of size $d \times d$ is irreducible if and only if

$$\sum_{k=0}^{d-1} P^k > \mathbf{0},$$

where $P^0 = I$, $\mathbf{0}$ denotes the all-zeros matrix of size $d \times d$, and the inequality is entry-wise.

A simple algorithm to check for irreducibility:

- Start with $M_0 = I$.
- Recursively define

$$M_{\ell+1} = I + PM_{\ell}.$$

Test for Irreducibility

Theorem (Test for Irreducibility)

A row stochastic matrix P of size $d \times d$ is irreducible if and only if

$$\sum_{k=0}^{d-1} P^k > \mathbf{0},$$

where $P^0 = I$, $\mathbf{0}$ denotes the all-zeros matrix of size $d \times d$, and the inequality is entry-wise.

A simple algorithm to check for irreducibility:

- Start with $M_0 = I$.
- Recursively define

$$M_{\ell+1} = I + PM_{\ell}.$$

- If $M_{\ell} > \mathbf{0}$ for any $1 \leq \ell \leq d - 1$, then P is irreducible.

Sample Path of a Random Process in Simulations

```

1 import numpy as np
2 import matplotlib.pyplot as plt
3 from matplotlib.animation import FuncAnimation
4 from scipy.stats import bernoulli, norm, gaussian_kde
5 from IPython.display import HTML
6
7 # Parameters of Bernoulli distribution
8 p = 0.5
9 mu = p
10 sigma = np.sqrt(p * (1 - p))
11
12 # Sample Size Progression
13 x_range = np.linspace(-4, 4, 500)
14 x_val = np.pi / np.sqrt(3)
15 sample_sizes = np.linspace(10, 1000, 50, dtype=int)
16
17 # True standard normal PDF
18 true_pdf = norm.pdf(x_range)
19
20 # Initialize figure
21 fig, ax = plt.subplots(figsize=(10, 6))
22 line_kde = ax.plot([], [], label='Estimated KDE PDF', color='blue')
23 line_true = ax.plot(x_range, true_pdf, '--', label='Standard Normal', color='black')
24 vline = ax.axvline(x_val, color="gray", linestyle=":")
25 title = ax.set_title('')
26 ax.set_xlim(-4, 4)
27 ax.set_ylim(0, 0.5)
28 ax.legend()
29
30
31 # Update function for animation
32 def update(n):
33     samples = bernoulli.rvs(p, size=(10000, n))
34     means = np.mean(samples, axis=1)
35     Z = (means - mu) / (sigma / np.sqrt(n))
36     kde = gaussian_kde(Z)
37     kde_vals = kde(x_range)
38     line_kde.set_data(x_range, kde_vals)
39     est_pdf_at_x = kde(x_val).item()
40     true_pdf_at_x = norm.pdf(x_val)
41     title.set_text(f"n={n}, PDF@x={x_val:.2f} = {est_pdf_at_x:.4f}, N(0,1) PDF@x = {true_pdf_at_x:.4f}")
42     return line_kde, line_true, vline, title
43
44 # Create animation
45 ani = FuncAnimation(fig, update, frames=sample_sizes, blit=True, interval=200)
46 HTML(ani.to_jshtml())
47

```


Sample Path of a Random Process in Simulations

```
1 # Generate an array of 10 numbers according to a unit Gaussian distribution
2 rng.standard_normal(10)
```

```
array([ 0.52788774, -0.89551364,  0.78844006,  0.19032279,  0.92833631,
        0.00333179, -1.08593089, -0.52056583,  1.06936692, -0.51129284])
```

Question


Can we assume that the generated samples correspond to a given $\omega \in \Omega$?

That is, can we assume that

$$X_1(\omega) = 0.52788774, \quad X_2(\omega) = -0.89551364, \dots$$

Back to Basics


$$(\Omega, \mathcal{F}, \mathbb{P}) + X \longrightarrow F_X$$



$$F_X(x) = \mathbb{P}_X((-\infty, x]) = \mathbb{P}(X^{-1}((-\infty, x])), \quad x \in \mathbb{R}$$

Back to Basics

$$(\Omega, \mathcal{F}, \mathbb{P}) + X \longrightarrow F_X$$



$$F_X(x) = \mathbb{P}_X((-\infty, x]) = \mathbb{P}(X^{-1}((-\infty, x])), \quad x \in \mathbb{R}$$

Proposition (From CDF to Probability Space and Random Variable)

Given a function $F : \mathbb{R} \rightarrow [0, 1]$ satisfying the properties of a CDF, there exists $(\Omega', \mathcal{F}', \mathbb{P}')$ and a random variable $X' : \Omega' \rightarrow \mathbb{R}$ such that $F_{X'} = F$.

Significance of Proposition

- In most practical scenarios, we are not given $(\Omega, \mathcal{F}, \mathbb{P})$ and X to begin with

Significance of Proposition

- In most practical scenarios, we are not given $(\Omega, \mathcal{F}, \mathbb{P})$ and X to begin with
- Despite this restriction, we generate samples from a custom CDF on computer, and treat these samples as output of a random variable (process)

Significance of Proposition

- In most practical scenarios, we are not given $(\Omega, \mathcal{F}, \mathbb{P})$ and X to begin with
- Despite this restriction, we generate samples from a custom CDF on computer, and treat these samples as output of a random variable (process)
- The natural question is: is this treatment justified?

Significance of Proposition

- In most practical scenarios, we are not given $(\Omega, \mathcal{F}, \mathbb{P})$ and X to begin with
- Despite this restriction, we generate samples from a custom CDF on computer, and treat these samples as output of a random variable (process)
- The natural question is: is this treatment justified?
- The preceding theorem says: **YES!**

From Single Random Variable to a Random Process

- Suppose that we generate n samples x_1, x_2, \dots, x_n , all from a custom CDF F

From Single Random Variable to a Random Process

- Suppose that we generate n samples x_1, x_2, \dots, x_n , all from a custom CDF F
- We would like to associate each sample to one random variable (say x_i to a RV X_i), and treat the samples as the outputs of the random variables

From Single Random Variable to a Random Process

- Suppose that we generate n samples x_1, x_2, \dots, x_n , all from a custom CDF F
- We would like to associate each sample to one random variable (say x_i to a RV X_i), and treat the samples as the outputs of the random variables
- However, it would not be desirable to have

$$x_1 = X_1(\omega_1), \quad x_2 = X_2(\omega_2), \dots,$$

because there exists no theory for analysing the behaviour of such samples

From Single Random Variable to a Random Process

- Suppose that we generate n samples x_1, x_2, \dots, x_n , all from a custom CDF F
- We would like to associate each sample to one random variable (say x_i to a RV X_i), and treat the samples as the outputs of the random variables
- However, it would not be desirable to have

$$x_1 = X_1(\omega_1), \quad x_2 = X_2(\omega_2), \dots,$$

because there exists no theory for analysing the behaviour of such samples

- What we desire is

$$x_1 = X_1(\omega), \quad x_2 = X_2(\omega), \dots,$$

a scenario wherein all samples correspond to a given ω

From Single Random Variable to a Random Process

- Suppose that we generate n samples x_1, x_2, \dots, x_n , all from a custom CDF F
- We would like to associate each sample to one random variable (say x_i to a RV X_i), and treat the samples as the outputs of the random variables
- However, it would not be desirable to have

$$x_1 = X_1(\omega_1), \quad x_2 = X_2(\omega_2), \dots,$$

because there exists no theory for analysing the behaviour of such samples

- What we desire is

$$x_1 = X_1(\omega), \quad x_2 = X_2(\omega), \dots,$$

a scenario wherein all samples correspond to a given ω

- Is there a feasible fit for the latter viewpoint?

From Single Random Variable to a Random Process

- Suppose that we generate n samples x_1, x_2, \dots, x_n , all from a custom CDF F
- We would like to associate each sample to one random variable (say x_i to a RV X_i), and treat the samples as the outputs of the random variables
- However, it would not be desirable to have

$$x_1 = X_1(\omega_1), \quad x_2 = X_2(\omega_2), \dots,$$

because there exists no theory for analysing the behaviour of such samples

- What we desire is

$$x_1 = X_1(\omega), \quad x_2 = X_2(\omega), \dots,$$

a scenario wherein all samples correspond to a given ω

- Is there a feasible fit for the latter viewpoint?

From Single Random Variable to a Random Process

- Suppose that we generate n samples x_1, x_2, \dots, x_n , all from a custom CDF F
- We would like to associate each sample to one random variable (say x_i to a RV X_i), and treat the samples as the outputs of the random variables
- However, it would not be desirable to have

$$x_1 = X_1(\omega_1), \quad x_2 = X_2(\omega_2), \dots,$$

because there exists no theory for analysing the behaviour of such samples

- What we desire is

$$x_1 = X_1(\omega), \quad x_2 = X_2(\omega), \dots,$$

a scenario wherein all samples correspond to a given ω

- Is there a feasible fit for the latter viewpoint? **YES!**

Proposition (From CDF to Probability Space and Random Variable)

Given a function $F : \mathbb{R} \times \mathbb{R} \times \cdots \rightarrow [0, 1]$ satisfying the properties of a joint CDF, there exists $(\Omega', \mathcal{F}', \mathbb{P}')$ and a random process $X' : \Omega' \rightarrow \mathbb{R} \times \mathbb{R} \times \cdots$ such that $F_{X'} = F$.