

/ Move down content because we have a fixed navbar that is 4.5rem tall / body { padding-top: 4.5rem; } hr { margin-top: 1rem; margin-bottom: 1rem; border: 1px solid #ccc; border-top: 1px solid #ccc; }

Eddy summary

/Users/kevin/eddy-squeeze/eddy_squeeze/eddy_squeeze_lib/
eddy_summary

Number of Eddy data included in the summary: 2

[Back to study level summary »](#)

Data summary for each Eddy runs

Basic information

subject	eddy_dir	number of volumes	max b value	min b value	unique b values	number of b's
subject	/Users/kevin/eddy-squeeze/tests/prac_study_dir/subject		.	.	[. . . .]	.
subject	/Users/kevin/eddy-squeeze/tests/prac_study_dir/subject		.	.	[. . . .]	.

Outlier information


subject	number of outlier slices	Sum of standard deviations in outlier slices	Mean of standard deviations in outlier slices	Standard deviation of standard deviations in outlier slices
subject		.	.	.
subject		.	.	.


Motion information

subject	absolute restricted movement	relative restricted movement
subject	.	.
subject	.	.

Summary figures


/Users/kevin/eddy-squeeze/eddy_squeeze/eddy_squeeze_lib/eddy_summary/plot_post_eddy_shell_PE_Deduced PE-translations (mm) relative b0-shell.png
These between shell PE-translations were calculated using MI between shell means
Deduced PE-translations (mm) relative b0-shell







/Users/kevin/eddy-squeeze/eddy_squeeze/eddy_squeeze_lib/eddy_summary/plot_post_eddy_shell_These between shell parameters were calculated using MI between shell means.png

Relative movement parameters between the shells
These between shell parameters were calculated using MI between shell means







/Users/kevin/eddy-squeeze/eddy_squeeze/eddy_squeeze_lib/eddy_summary/plot_post_eddy_shell_These between shell
parameters were calculated using interspersed b -volumes.png
Movement parameters relative b0-shell
These between shell parameters were calculated using interspersed b0-volumes




relative b0-shell from direct registration to mean b0-shell .png
These between shell PE-translations were calculated using MI between shell means
PE-translations (mm) relative b0-shell from direct registration to mean b0






/Users/kevin/eddy-squeeze/eddy_squeeze/eddy_squeeze_lib/eddy_summary/plot_post_eddy_shell_PE_Relative PE-translations
(mm) between the shells.png
These between shell PE-translations were calculated using MI between shell means
Relative PE-translations (mm) between the shells





/Users/kevin/eddy-squeeze/eddy_squeeze/eddy_squeeze_lib/eddy_summary/plot_post_eddy_shell_These are the movement parameters that have been applied to the data.png


Movement parameters relative b0-shell
These are the movement parameters that have been applied to the data



Standard deviation of standard deviations in outlier slices









BRIGHAM AND
WOMEN'S HOSPITAL



HARVARD
MEDICAL SCHOOL

© Kang Ik Kevin Cho & Ofer Pasternak, Department of Psychiatry, Psychiatry Neuroimaging Laboratory, Brigham and Women's Hospital, Harvard Medical School, Boston, MA

Next Generation Diffusion MRI Biomarkers for Prodromal Schizophrenia (R MH - A)