NCCP 00446 · VERSION 7B

Your treatment, explained

Metastatic colorectal cancer · NCCP 00446 v7b · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Modified FOLFOX-6 + bevacizumab

14-day cycle

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

ACT ON SYMPTOMS, NOT THE CALENDAR

When to get help

Emergency: call 112 / 999

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

Contact your oncology service immediately

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. Contact promptly for new chest symptoms, severe or persistent diarrhoea, reduced urine, inability to drink, severe abdominal pain or a line/pump problem. New neurological symptoms, uncontrolled bleeding or a swollen painful limb also need urgent assessment.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.

Open your CUH contact / treatment card

Sources: nccp · hse

Choose a topic to read

YOUR COMPLETE REFERENCE

Read what you need, when you need it.

Start with the overview, explore the cycle, or go straight to a question. Urgent advice is always available.

- How each drug works
- Your treatment plan
- Your 14-day timeline
- Baseline and cycle-specific checks
- Whole treatment, scans and checks
- Your port and treatment pump
- Tablets, visits and practical planning
- Your treatment aim and review plan
- Bowel symptoms: act early
- Tests and preparation
- Eating, activity and daily life
- Medicines and supportive care
- Prepare for the next conversation
- Bevacizumab: vessels, wounds and checks
- The nerve-check schedule
- Blood pressure, urine and anticoagulants

Supportive medicines
Understanding the evidence
Questions to ask
Sources

UNDERSTAND YOUR TREATMENT

Your medicines and what they do

Modified FOLFOX-6 + bevacizumab is used for cancer of the colon or rectum that has spread. Treatment aims to control the cancer; the likely benefit and review plan depend on your individual situation. Surgery may be reconsidered in selected cases after multidisciplinary review.

ANTI-VEGF TARGETED ANTIBODY

Bevacizumab

Avastin or a prescribed biosimilar



Blocks a blood-vessel growth signal

Bevacizumab binds VEGF, a signal that helps blood vessels grow. Blocking it can restrict the blood supply supporting a tumour. It also affects normal blood vessels and healing, which is why blood pressure, urine and planned procedures matter.

VEGF signal → Bevacizumab binds VEGF → New-vessel signalling is reduced

What to watch for

Report bleeding, a wound that is not healing, severe abdominal pain, new swelling, chest symptoms or a severe headache. Tell the team before surgery, dental procedures or port insertion.

Sources: nccp · patient

PLATINUM CHEMOTHERAPY

Oxaliplatin

Eloxatin



Links DNA strands

Oxaliplatin makes links within DNA. These interfere with copying and repair, which can stop a cancer cell dividing. Healthy tissues can also be affected.

Platinum enters cell → DNA links form → Copying is disrupted

What to watch for

Cold can trigger tingling or throat discomfort. Tell the team about persistent numbness, difficulty with buttons or walking, or infusion symptoms. New breathing difficulty always needs assessment.

Sources: nccp · patient

CHEMOTHERAPY MODULATOR

Folinic acid

Calcium leucovorin



Strengthens fluorouracil action

Folinic acid helps fluorouracil remain bound to an enzyme needed to make DNA. In this regimen it increases fluorouracil activity and can also increase its unwanted effects. It is not a vitamin supplement to start yourself.

Folinic acid → Stronger enzyme binding → Fluorouracil effect increases

---

What to watch for


The combined treatment can increase mouth soreness and diarrhoea. Report these promptly. Rare allergic reactions can occur.

Sources: nccp · patient

CHEMOTHERAPY

Fluorouracil

5-FU



Disrupts cell building blocks

Fluorouracil interferes with the production and use of DNA and RNA, the instructions cells need to grow. The infusion extends exposure over approximately two days.

Drug enters cell → DNA / RNA disrupted → Growth slows

What to watch for

Diarrhoea, mouth sores and skin soreness may become serious. Chest pain during the pump infusion needs emergency assessment. Unexpected severe early toxicity can occur even after DPD testing.

Sources: nccp · patient

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

Explore your existing anatomy and side-effect map

THE BIGGER TREATMENT PLAN

Your treatment plan

DAY WARD → HOME PUMP → REVIEW

14-day treatment cycle

D1 D2 D3 D14

Day 1: bevacizumab, oxaliplatin, folinic acid and fluorouracil, then a 46-hour fluorouracil pump. A cycle lasts 14 days; your team confirms the next treatment.

Treatment is reviewed against benefit, side effects and your priorities. Reductions, breaks or a maintenance plan are individual decisions, not an automatic change after a set number of cycles.

Sources: nccp

BASELINE → TREATMENT → FOLLOW-UP

Checks across the whole treatment course

The columns show the order of care, not equal lengths of time. Confirm who books each check, who reviews it and whom to call if it is missing.

Check	Before starting	Each treatment cycle	Between doses	Decision / response review	Completion / change
Blood count / kidneys / liver	Baseline bloods; DPD before fluoropyrimidine	Repeat before each cycle	Earlier tests if unwell, dehydrated or toxicity develops	Results and recovery guide any delay or dose change	Follow unresolved abnormalities
Symptoms and function	Document existing symptoms and usual activity	Review worst symptoms since the last treatment	Use urgent contacts; do not wait for the next visit	Balance benefit, side effects and priorities	Agree recovery care and ongoing support
Blood pressure / urine / wounds	BP, urine protein and wound/procedure review	Blood pressure before each cycle AND after treatment; urine protein each cycle	Report bleeding, severe headache or abdominal pain urgently	Additional urine tests and specialist review if abnormal	Confirm wound-healing and procedure plan
Neuropathy / cold sensitivity	Record existing nerve symptoms	Formal neuropathy evaluation every two cycles; review symptoms sooner if they occur	Report worsening numbness or cold-triggered throat symptoms	Consultant reviews the oxaliplatin component separately	Nerve symptoms can continue after treatment; arrange review
Heart checks	Review cardiac history; ECG when cardiac function is compromised or clinical findings warrant it. No automatic serial echocardiogram schedule is specified for this regimen.	Ask about new chest pain or breathlessness	Chest pain needs urgent assessment	Repeat heart tests if clinically indicated	Follow any heart problem; no fixed echo series
Response / scans	Confirm cancer setting, disease assessment and intended duration	Review clinical benefit and symptoms	New symptoms may bring review forward	Consultant-directed scans and response assessment	Agree follow-up, completion or another treatment
INR only if taking warfarin	Tell the team about warfarin and arrange a baseline INR	INR weekly until the warfarin dose is stable, then before each cycle	Report bleeding or unusual bruising	Only change anticoagulants as instructed	Continue the individually agreed anticoagulation follow-up

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00446 v7b · official regimen

THE WHOLE TREATMENT JOURNEY

Modified FOLFOX-6 + bevacizumab: treatment and checks

Metastatic colorectal cancer

Day 1: bevacizumab, oxaliplatin, folinic acid and fluorouracil, then a 46-hour fluorouracil pump. A cycle lasts 14 days; your team confirms the next treatment.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

On a small screen, swipe the map sideways. The checks below give the same timing in words.

Treatment, tests and reviews across the course

Care / check	Before starting	Each treatment cycle	Between doses	Decision / response review	Completion / change
Blood count / kidneys / liver NCCP 00446	Baseline bloods; DPD before fluoropyrimidine	Repeat before each cycle	Earlier tests if unwell, dehydrated or toxicity develops	Results and recovery guide any delay or dose change	Follow unresolved abnormalities
Symptoms and function NCCP 00446	Document existing symptoms and usual activity	Review worst symptoms since the last treatment	Use urgent contacts; do not wait for the next visit	Balance benefit, side effects and priorities	Agree recovery care and ongoing support
Blood pressure / urine / wounds NCCP 00446	BP, urine protein and wound/procedure review	Blood pressure before each cycle AND after treatment; urine protein each cycle	Report bleeding, severe headache or abdominal pain urgently	Additional urine tests and specialist review if abnormal	Confirm wound-healing and procedure plan
Neuropathy / cold sensitivity NCCP 00446	Record existing nerve symptoms	Formal neuropathy evaluation every two cycles; review symptoms sooner if they occur	Report worsening numbness or cold-triggered throat symptoms	Consultant reviews the oxaliplatin component separately	Nerve symptoms can continue after treatment; arrange review
Heart checks NCCP 00446	Review cardiac history; ECG when cardiac function is compromised or clinical findings warrant it. No automatic serial echocardiogram schedule is specified for this regimen.	Ask about new chest pain or breathlessness	Chest pain needs urgent assessment	Repeat heart tests if clinically indicated	Follow any heart problem; no fixed echo series
Response / scans NCCP 00446	Confirm cancer setting, disease assessment and intended duration	Review clinical benefit and symptoms	New symptoms may bring review forward	Consultant-directed scans and response assessment	Agree follow-up, completion or another treatment
INR only if taking warfarin NCCP 00446	Tell the team about warfarin and arrange a baseline INR	INR weekly until the warfarin dose is stable, then before each cycle	Report bleeding or unusual bruising	Only change anticoagulants as instructed	Continue the individually agreed anticoagulation follow-up

What each check is for

1

14-day treatment cycle

Day ward → home pump → review

Modified FOLFOX-6 + bevacizumab is used for cancer of the colon or rectum that has spread. Treatment aims to control the cancer; the likely benefit and review plan depend on your individual situation. Surgery may be reconsidered in selected cases after multidisciplinary review.

Ask / act: Confirm the phase, prescription, tests and supportive medicines before starting.

Sources: NCCP 00446

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?


This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-04 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.33

Day 1 → Day 14: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

Day 1: bevacizumab, oxaliplatin, folinic acid and fluorouracil, then a 46-hour fluorouracil pump. A cycle lasts 14 days; your team confirms the next treatment.



Mouth / bowel / energy	Record symptoms throughout
Bevacizumab effects	BP / urine checks; report new symptoms anytime

Day ward and home pump: Day 1 infusions lead into a 46-hour pump. The team confirms the disconnection appointment.

No safe window: Fever or sudden illness needs prompt action on any day, including after pump removal.

Cold sensitivity: Tingling or throat discomfort can occur early and may persist; follow the precautions your team gives you.

Ongoing checks: Blood pressure and urine checks may detect bevacizumab effects before you feel them. Report new symptoms throughout treatment.

> **Why these bands stay broad:** Bars show planned treatment and possible symptom patterns, not measured risk. Side effects can overlap and persist. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00446 v7b · official regimen ↗ · eviQ: patient explanation (NCCP defines your regimen schedule) ↗

> **There are no guaranteed “safe days”.** Act on fever, sudden illness or a new serious symptom whenever it occurs. Do not wait for the next appointment.

Read the whole cycle as text

Day 1: Infusion and pump connection

The team reviews symptoms, blood results and your treatment plan, including blood pressure and urine protein. Day-ward infusions are followed by connection of the fluorouracil pump. Before leaving, confirm the pump-care instructions, disconnection appointment and contact number.

Day 2–3: The pump is delivering treatment

The pump runs for approximately 46 hours from connection. Keep it and the line secure and dry as taught. The team confirms the actual disconnection appointment. Call for a leaking, disconnected or unexpectedly stopped pump. Do not reconnect, alter settings or disconnect it yourself unless specifically trained and instructed.

Day 4–7: After disconnection

Nausea, diarrhoea, mouth symptoms, fatigue or cold sensitivity may continue. Use prescribed support. New symptoms deserve advice even if the pump has finished. Report poor intake, persistent diarrhoea or mouth pain early. Chest pain or major breathing difficulty requires emergency help.

Day 8–14: Between visits and the next review

Blood counts can fall between visits. Tell the team about bowel symptoms, mouth pain and how you are managing daily activities, and any persistent numbness. Treatment proceeds only after review. Bring the symptom passport and check the next appointment. Day 1 usually repeats 14 days after this cycle started, only if confirmed.

YOUR TREATMENT, STEP BY STEP

Your port and treatment pump

From preparing the line to coming home after disconnection. Your team arranges the appointments and shows you what to do.



1 · Before treatment

Choose and prepare the line

The team explains whether a port or PICC suits your treatment. You receive instructions for the insertion appointment.



2 · Insertion and recovery

A separate appointment may be needed

For a port, numbing medicine is used and a small device is placed under the skin. Ask about sedation, transport and wound care.



3 · Day 1

The nurse connects your pump

After the day-ward medicines, the nurse checks and connects the fluorouracil pump. Ask for a demonstration before leaving.

4 · At home

Treatment continues for about 46 hours

Carry the pump as shown. Keep the dressing dry and protect the line from pulling, heat and damage.

5 · As arranged

Return for disconnection

A trained professional checks the infusion, disconnects the pump and cares for the line. The appointment is based on connection time.

6 · Between cycles

The access device usually stays

A port needle can be removed while the port stays under the skin. A PICC remains in the arm with ongoing dressing and line care.

Receiving bevacizumab? Tell the oncology and procedural teams before port insertion, surgery or dental work. They agree treatment timing and wound review.

Port or PICC: what is the difference?

A port is a small chamber beneath the skin, usually on the chest, attached to a tube ending in a large vein. A special needle goes through the skin into the port when it is used. A PICC enters a vein in the upper arm; a short section remains outside under a dressing. Both can deliver a home infusion. Your team helps choose the device and teaches its care.

Before insertion and during recovery

Tell the team about allergies, blood thinners, infections and any other cancer medicines. Follow the instructions about eating, drinking and transport; do not stop medicines yourself. Port placement uses local anaesthetic, sometimes with sedation or general anaesthetic. The team explains the procedure and risks, including infection, bleeding, clots and insertion complications. Ask how to manage discomfort, protect the wound and resume activity. Report increasing pain, redness, swelling, discharge or fever.

Connection: what the nurse does

The nurse checks the line and treatment, secures the connection and dressing, and starts the pump. Before leaving, you should have the disconnection location and time, contact numbers including out of hours, and the written spill instructions. Ask to be shown what normal pump progress looks like. You do not need to calculate a rate.

Carrying it, sleeping and washing

Use the supplied carrier and the position recommended for your pump. Keep tubing free from pulling and kinks. Ask how to place the pump safely for sleep and how to protect the line and dressing when washing. Do not immerse it or swim. Avoid direct heat and cold. An elastomeric pump has a balloon that shrinks slowly and may have no alarm; an electronic pump has different instructions. Follow the instructions for the device you actually receive.

Disconnection is not port removal

Keep the appointment even if the pump looks empty, unless the team changes the plan. Staff check the pump and line and arrange safe disposal. After flushing and other line care as required, the port needle may be removed and a small dressing applied. The implanted port stays in place until a separate removal procedure is arranged. A PICC remains in place and needs scheduled dressing and flushing care.

Leakage, disconnection or unexpected pump progress

Call the team immediately if the pump leaks, is damaged, finishes unexpectedly early, appears not to run, or the line disconnects. Follow the emergency or spill steps you have been specifically taught, including use of a clamp if instructed. Do not reconnect, change settings or disconnect the pump yourself. Keep other people away from leaked medicine. If medicine touches skin, rinse it off and wash with soap and water, then seek advice.

Pain, swelling or feeling unwell

Contact oncology immediately for pain or swelling around the line, a wet/loose dressing, redness, fever, shivering or suddenly feeling unwell. A painful swollen arm or neck needs urgent advice. Severe chest pain, major breathing difficulty or collapse: call 112 / 999. Do not wait for pump disconnection or the next clinic visit.

The drawings show general steps, not instructions for inserting or accessing a line. Your hospital confirms its own arrangements and the care instructions for your device.

Open the complete port and pump guide · **Printable port and pump guide**

Sources: eviQ: port · eviQ: PICC · eviQ: pump. Reviewed against these sources 16 October 2026; local nursing and patient review pending.

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Before the first treatment

Confirm the aim, likely course length, blood-test arrangements and a working urgent contact number. DPD testing is required before starting the fluoropyrimidine unless already completed.

STEP 2

Line and pump arrangements

Confirm venous access, the 46-hour fluorouracil pump and disconnection arrangements before leaving. Know whom to contact for leakage, an alarm, line pain or a pump that finishes unexpectedly.

STEP 3

If treatment is delayed

Ask the unit to confirm the new treatment and blood-test dates. Re-anchor supportive medicines to the treatment actually given; do not take an extra or catch-up anticancer dose yourself.

STEP 4

Getting help between visits

Follow your oncology unit’s urgent advice for fever, infection, severe diarrhoea, chest pain or difficulty breathing. Bring your treatment card and current medicines list.

STEP 5

Operations and dental procedures

Tell both the oncology team and the procedural team that you receive bevacizumab. Holds and restart need an agreed wound-healing plan; do not stop or restart it yourself.

STEP 6

Paper copies, wallet QR and contacts

The example telephone numbers on the wallet card are placeholders. Your team must replace them with working daytime and out-of-hours numbers before patient use. Keep the card with you; the QR opens the regimen information, not your confidential medical record.

Open practical treatment guide · **Printable practical guide**

Sources: NCCP 00446 v7b · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.

MAKE THE PLAN PRACTICAL

Support around the medicines

1 Nausea prevention and rescue

Take the prescribed prevention medicines at the planned times and keep rescue treatment available. Tell the team if nausea or constipation is not controlled; do not duplicate steroid or anti-sickness medicines.

2 Bowel symptoms

Report diarrhoea promptly and follow the written rescue plan. Severe symptoms, fever, dehydration or inability to drink need urgent advice.

3 Counts and recovery

White-cell growth-factor injections are prescribed according to individual risk and previous complications. They are not automatically required for every patient or every cycle.

4 Cold and nerve symptoms

Avoid cold exposure as advised after oxaliplatin. Report persistent numbness or difficulty with buttons, walking or everyday tasks before the next dose.

5 Blood pressure and wound care

Keep the agreed blood-pressure and urine checks. Report bleeding, poorly healing wounds, new swelling or severe headache promptly.

Sources: nccp

TREATMENT REFERENCE · 01

Your treatment aim and review plan

Ask what treatment is intended to achieve for you.

Confirm the cancer setting

Modified FOLFOX-6 + bevacizumab is used for cancer of the colon or rectum that has spread. Treatment aims to control the cancer; the likely benefit and review plan depend on your individual situation. Surgery may be reconsidered in selected cases after multidisciplinary review.

Sources: nccp · patient

Review rather than a fixed promise

Ask when scans and clinical review will assess benefit. Treatment may be adjusted if it is not helping or side effects outweigh benefit. A cycle delay is not itself proof that treatment has stopped working.

Sources: nccp · patient

Your priorities matter

Discuss symptoms you want to improve, daily activities, work, caring responsibilities and time at home. The team can explain alternatives and how they compare in your situation.

Sources: nccp · patient

TREATMENT REFERENCE · 02

Bowel symptoms: act early

Keep the written plan with your medicines.

During the infusion

Tell the nurse about pain, a reaction or feeling unwell during treatment. Oxaliplatin can cause cold-triggered nerve symptoms; new breathing difficulty always needs assessment.

Sources: nccp · patient

At home

Record changes from your usual bowel pattern or stoma output. Follow your prescribed diarrhoea plan and contact oncology if symptoms persist or you cannot replace lost fluids. Do not wait for the next scheduled visit.

Sources: nccp · patient

Get urgent advice

Fever, severe pain, blood in stools, dizziness, reduced urine or inability to drink alongside diarrhoea needs urgent assessment. Severe abdominal pain must not be assumed to be an ordinary treatment effect.

Sources: nccp · patient

TREATMENT REFERENCE · 03

Tests and preparation

Checks support safe treatment even when you feel well.

Blood tests and symptoms

The team reviews blood counts, kidney and liver function and symptoms before another cycle. Missing information is not assumed to be normal.

Sources: nccp · patient

DPD and fluorouracil

DPD helps the body break down fluorouracil. Deficiency can cause severe toxicity. Testing guides the plan but cannot exclude all serious reactions; unexpectedly severe early diarrhoea, mouth sores or illness needs immediate review.

Sources: nccp · patient

Additional checks

Bevacizumab requires blood-pressure and urine-protein checks. Report any wound, bleeding, clot history or planned procedure.

Sources: nccp · patient

TREATMENT REFERENCE · 04

Eating, activity and daily life

Choose manageable steps and ask for help early.

Food and fluids

Take prescribed anti-sickness medicines. Small regular meals may be easier when appetite is low. Tell the team if vomiting, mouth soreness or diarrhoea prevents enough intake; follow any fluid restriction you have.

Sources: hse · patient

Activity and rest

Short, comfortable activity and planned rests can help you organise the day. Avoid activity that pulls the line or risks damaging the pump. Ask about work, driving and travel around your own treatment and symptoms.

Sources: hse · patient · pump

Sex, fertility and pregnancy

Discuss fertility preservation before treatment where relevant. Ask about contraception, pregnancy, breastfeeding and precautions around sex during treatment; the required precautions depend on the medicines.

Sources: nccp · patient

TREATMENT REFERENCE · 05

Medicines and supportive care

Bring a complete list, including supplements.

Before adding medicines

Check new prescriptions, over-the-counter products and supplements with the oncology pharmacist. Do not independently stop regular medicines such as anticoagulants before a procedure.

Sources: nccp · patient

Supportive prescriptions

Keep anti-sickness medicines and the bowel-symptom plan accessible. If growth-factor injections are prescribed, confirm when they are due in relation to the pump; do not infer timing from the graph.

Sources: nccp · patient

Treatment changes

Ask which medicine is being delayed, reduced or stopped and why. Each medicine can change separately; record the agreed plan in your own treatment documents.

Sources: nccp · patient

TREATMENT REFERENCE · 06

Prepare for the next conversation

You do not have to remember everything.

Bring your questions

Use the passport to note symptoms and their effect on eating, drinking, sleep and everyday tasks. You do not need to work out which medicine caused them.

Sources: nccp · patient

Ask for practical support

Ask for a written appointment and contact plan, an interpreter if needed, and help for a family member or carer learning the pump instructions.

Sources: nccp · patient

Tell us what was unclear

Ask the nurse to demonstrate the pump and line, then explain back when and how you would seek help. More explanation is appropriate whenever you need it.

Sources: nccp · patient

TREATMENT REFERENCE · 07

Bevacizumab: vessels, wounds and checks

Some effects are found by monitoring before symptoms appear.

Why blood pressure and urine?

Blocking VEGF can affect normal blood vessels and kidney filtering. Blood pressure and urine-protein checks help the team decide whether bevacizumab needs review.

Sources: nccp · patient

Before a procedure

Tell oncology and the procedural team about bevacizumab before port insertion, dental work or surgery. They agree treatment timing and wound review. Do not apply a generic waiting period yourself.

Sources: nccp · patient

Symptoms that need action

Seek urgent advice for uncontrolled bleeding, severe abdominal pain, severe headache or visual change, new weakness, chest pain, breathlessness or a painful swollen limb. Call 112 / 999 for collapse or severe breathing/chest symptoms.

Sources: nccp · patient

TREATMENT REFERENCE · 08

The nerve-check schedule

Reporting symptoms and scheduled assessment work together.

Formal review every two cycles

This NCCP regimen specifies neuropathy evaluation every two cycles. Report worsening numbness, tingling or trouble with buttons or walking immediately rather than waiting for that review.

Sources: nccp

Cold precautions

Oxaliplatin can cause cold-triggered tingling and throat sensations. Avoid cold triggers and do not use ice chips. New breathing difficulty still requires assessment; it must not be assumed to be only cold sensitivity.

Sources: nccp

TREATMENT REFERENCE · 09

Blood pressure, urine and anticoagulants

Keep the checks even if you feel well.

During every cycle

Blood pressure is measured before each cycle and after treatment. Urine protein is checked during treatment; abnormal results may lead to more testing or a treatment hold.

Sources: nccp

If taking warfarin

INR is checked weekly until the warfarin dose is stable, then before each cycle. This requirement applies to warfarin, not to every patient. Report bleeding or unusual bruising and do not change an anticoagulant yourself.

Sources: nccp

Understanding the evidence

Ask which evidence matches your cancer, treatment aim and medicines.

Adding bevacizumab to oxaliplatin chemotherapy delayed cancer progression in the trial reviewed, but did not establish an overall survival improvement. The study pooled two chemotherapy schedules, neither identical to this modified FOLFOX6 regimen. Your team weighs this evidence against side effects and your priorities.

Sources: trial

PREPARE FOR THE CONVERSATION

Questions worth asking

Does having more side effects mean treatment is working?

No. Side effects do not reliably show whether the cancer is responding. Report them so they can be treated. Your team assesses benefit using the tests and review plan appropriate to your cancer.

Why might a dose be delayed or reduced?

Your team may allow recovery, investigate a symptom or reduce the chance of lasting harm. Ask which drug changes, why, and when review is planned. A delay does not itself mean treatment has failed.

What should I say when calling?

Give the regimen name, last treatment date, symptoms and when they started. Explain whether you can eat, drink, pass urine and manage usual activities. Give your temperature if available, but do not delay calling to collect details.

What belongs in the consent conversation?

The individual aim and likely benefit, uncertainty, reasonable alternatives including not proceeding, common and serious risks, lasting effects, fertility and practical concerns. You can ask for an interpreter, more time or another explanation and can reconsider your decision.

How many cycles should I use this passport for?

The passport provides reusable recording pages, not a prescribed course length. Your oncology team sets the duration and can provide extra pages.

Will I need a port or a PICC?

The team chooses the most suitable access device with you. A port is under the skin; a PICC enters through the upper arm and has an external section. Both require specific care, and the pump can be used with either.

Does disconnecting the pump remove my port?

No. At the arranged visit the pump is disconnected and the line cared for. A port needle may be removed, but the implanted port stays under the skin until a separate removal procedure is arranged.

What if the pump finishes early or still looks full?

Contact the team for advice. Do not change delivery settings or disconnect or reconnect it yourself. Follow only the specific emergency steps you have been taught.

> **Try explaining the plan back:** Can you explain which medicines you are receiving, the schedule, the contact plan and the symptoms that need prompt help?

SOURCE TRANSPARENCY

Where this information comes from

Regimen schedule: NCCP 00446 v7b. Medicine explanations and safety information: sources below. The timeline is an educational illustration, not a measured risk curve or an individual prediction.

- NCCP 00446 v7b · official regimen
- eviQ: patient explanation (NCCP defines your regimen schedule)
- HSE: chemotherapy side effects and urgent advice
- NCI: nausea and vomiting
- eviQ: treatment through an implanted port
- eviQ: treatment through a PICC
- eviQ: elastomeric infusion pump care
- NO16966 (2008)

SACTCheck · Developed by Dr Paul O’Brien · Sources checked 2026-11-04.
This information supports your treatment discussion. Your oncology team confirms your personal treatment and urgent-care plan.

Written source addresses

https://healthservice.hse.ie/documents/6478/446_v7b_Bevacizumab_5_plus_mFOLFOX-6.pdf

https://www.eviq.org.au/medical-oncology/colorectal/metastatic/986-colorectal-metastatic-folfox6-modified-fluo/patient-information

https://www.eviq.org.au/p/507

https://www.eviq.org.au/p/543

https://www.eviq.org.au/p/502

https://www2.hse.ie/conditions/chemotherapy/side-effects/

https://www.cancer.gov/about-cancer/treatment/side-effects/nausea-vomiting

https://pubmed.ncbi.nlm.nih.gov/18421054/