NCCP 00284 · clinician draft

Gemcitabine alone · non-small-cell lung cancer

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Gemcitabine alone · Days 1, 8, 15

Active medicines: gemcitabine

Emetogenic risk: low

Medicine	Dose / route	When
Dexamethasone	8 mg orally	Before each gemcitabine dose if prescribed NCIS low-risk reference; individualise and reconcile with any other steroid course.

Additional phase-specific support

- Gemcitabine has low sickness risk. Follow the prescribed preventive or rescue medicine plan; do not take a platinum anti-sickness course for gemcitabine alone.
- Do not dismiss fever as a flu-like drug effect. Call the oncology team immediately for fever, shaking chills or sudden illness.
- Follow your individual drinking advice. Report reduced urine, swelling, breathlessness or trouble drinking, and ask before new medicines or supplements.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00284 v5 · pNCCP 00284p3; NCISv8 low risk · NCCP NCIS antiemetics V 8

Sources checked 2026-11-02 · SACTCheck v0.84.29 · Regimen-level drafts; prescriber and oncology-pharmacy verification required