SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

Docetaxel alone for previously treated non-small-cell lung cancer

Lung · previously treated advanced NSCLC · NCCP 00203 · version 7 · 21-day cycle

Patient name	DOB	Hospital no.	Date

Treatment context
Treatment aims to slow or shrink lung cancer and relieve symptoms after previous chemotherapy. Expected benefit, side effects, other options and your priorities guide the decision.

Schedule
Docetaxel by drip on Day 1, repeated every 21 days, for up to six cycles. Treatment may stop earlier if it is not helping or side effects are unacceptable. Your team confirms the prescribed course.

1. TREATMENT DISCUSSION

- ☐ Purpose, expected benefit and alternatives explained
- ☐ Indication, medicines, sequence and duration confirmed
- ☐ Blood tests and any individually indicated heart checks discussed
- ☐ Support medicines and phase changes explained
- ☐ Fertility, contraception and breastfeeding discussed where relevant
- ☐ Common, serious and lasting drug-specific risks discussed below
- ☐ Urgent symptoms and local contact details confirmed

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Docetaxel
TAXANE CHEMOTHERAPY

- ☐ Low blood counts, infection and sepsis
- ☐ Hair loss, mouth soreness and fatigue
- ☐ Numbness / tingling; sometimes lasting
- ☐ Fluid retention and nail changes
- ☐ Allergic infusion reaction
- ☐ Rare serious bowel inflammation or perforation
- ☐ Rare serious lung or skin reactions; vision changes

Discuss treatment aim, alternatives including supportive care, uncertainty, serious and lasting effects, fertility and contraception, premedication, IV access and the urgent contact plan. Explain the six-cycle NSCLC maximum and the possibility of stopping earlier.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00203 v7.

SACTCheck v0.84.29
Developed by Dr Paul O'Brien