

# Modified FOLFOX-6

Gastrointestinal · NCCP 00209 · version 10a

## Your medicines

## Your treatment schedule

## A picture of side effects

Numbers connect each symptom box to the relevant anatomical area. Tell your team about new or worsening symptoms.

## Oxaliplatin · nerve and cold effects

## 5-FU / chemotherapy · other effects

### 1 Mouth / throat

## 5-FU / oxaliplatin

Mouth ulcers or soreness; sudden throat tightness needs prompt assessment, including after cold exposure.

### 2 Heart / breathing

## 5-fluorouracil (5-FU)

New chest pain, pressure, fainting or breathlessness needs urgent help, including during the infusion.

### 3 Blood / infection

## Oxaliplatin / 5-FU

Fever, shivers or sudden illness: contact oncology immediately.

### 4 Stomach / bowel

## 5-FU / oxaliplatin

Persistent diarrhoea, vomiting or inability to drink needs advice.

### 5 Hands / nerves

## Oxaliplatin

Cold-triggered tingling or numbness; report symptoms that persist between cycles.

### 6 Feet / nerves

## Oxaliplatin

Numbness, altered sensation or difficulty walking should be reported before the next dose.

### 7 Skin / mouth


## 5-fluorouracil (5-FU)

Report significant skin changes or mouth soreness that limits eating or drinking.

### 8 Line / home pump

## 46-hour 5-FU infusion

Call for leakage, disconnection, pain, redness, swelling or a pump alarm. Follow local pump instructions.



Selected effects only: the same symptom can have more than one cause, and not every possible side effect is shown.

# Modified FOLFOX-6

Gastrointestinal · NCCP 00209 · version 10a

## TREATMENT JOURNEY

|                                                                                           |                                                                                    |                                                                                           |                                                                                 |                                                                  |
|-------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------|------------------------------------------------------------------|
| <b>Day 1</b><br><br>Oxaliplatin and folinic acid are given by drip, then 5-FU is started. | <b>Days 1-3</b><br><br>A portable pump gives 5-FU continuously for about 46 hours. | <b>Around day 3</b><br><br>Pump disconnect. Follow your local line and pump instructions. | <b>Days 4-14</b><br><br>Recovery and monitoring. Symptoms can occur on any day. | <b>Day 15</b><br><br>Next cycle if review and blood tests allow. |
|-------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------|------------------------------------------------------------------|

## IMPORTANT MEDICINE-SPECIFIC EFFECTS

|                                                                                                                                                                                                                                                                                                         |                                                                                                                                                                                                                                                                                       |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Oxaliplatin</b> <ul style="list-style-type: none"><li>• Cold-triggered tingling or throat/jaw sensations can occur soon after treatment.</li><li>• Persistent peripheral neuropathy can develop with cumulative treatment.</li><li>• Infusion hypersensitivity can occasionally be severe.</li></ul> | <b>5-fluorouracil (5-FU)</b> <ul style="list-style-type: none"><li>• Severe toxicity can occur with reduced DPD activity.</li><li>• Diarrhoea and mouth/gut inflammation can become severe.</li><li>• New chest pain needs urgent assessment; hand-foot soreness can occur.</li></ul> |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

**OTHER / SHARED TREATMENT EFFECTS**

The anatomy page highlights important symptom patterns for this exact regimen, but it cannot show every possible effect. Fatigue, nausea, appetite change, mouth/bowel symptoms or blood-count changes may occur depending on the medicines used. Report new or worsening symptoms even if they are not listed.

|                                                                                                                                                                                                                                                                                                                                |                                                                                                                                                                                                                                                                                                                                                                                                                          |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Monitoring / medicines</b> <ul style="list-style-type: none"><li>• DPD status is checked before fluoropyrimidine treatment.</li><li>• Before each cycle: blood count, kidney/liver bloods and assessment of nerve symptoms.</li><li>• Follow your own labelled anti-sickness, mouth-care and diarrhoea medicines.</li></ul> | <b>Bring / track</b> <ul style="list-style-type: none"><li>• Your current medicines, including non-prescription products.</li><li>• A note of new symptoms, when they started and what they stopped you doing.</li><li>• Any temperature, blood-pressure or bowel-count readings your team asked you to record.</li><li>• Questions about the purpose of treatment, scan/review timing or changes to the plan.</li></ul> |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

**WHEN TO CALL - DO NOT WAIT FOR THE NEXT APPOINTMENT**

Call your oncology SOS service for fever or shivers, sudden illness, 4 or more diarrhoea episodes in a day, vomiting despite medicines, mouth sores stopping fluids, bleeding, new chest pain or breathlessness. Severe chest pain or breathing difficulty: call 112/999. If your team has given you different local thresholds or emergency instructions, follow those instructions.

## Consent discussion support

Use the approved consent process to document treatment aim, expected benefit, material risks, alternatives, questions and the patient's decision.

**Scan to open the regimen-specific patient portal**

<https://sactcheck.com/patient/00209/>

Official NCCP 00209 v10a

SACTCheck v0.81.1 · patient information prototype for clinical review. Follow your oncology team and local emergency instructions.

