

Pemetrexed + carboplatin

Lung · NCCP 00318 · version 6

Your medicines

Your treatment schedule

A picture of side effects

Numbers connect each symptom box to the relevant anatomical area. Tell your team about new or worsening symptoms.

Pemetrexed · kidney / mouth / skin

Carboplatin · blood counts / allergy / nausea

1 Blood / infection

Pemetrexed / carboplatin

Low blood counts - especially neutrophils and platelets - can cause infection or bleeding. Fever or sudden illness needs prompt contact.

2 Kidneys

Pemetrexed

Kidney function affects pemetrexed suitability and is checked before treatment. Report reduced urine or dehydration.

3 Nausea / bowel

Carboplatin / pemetrexed

Nausea, vomiting or diarrhoea can occur. Use prescribed supportive medicines and call if you cannot maintain fluids.

4 Mouth

Pemetrexed

Mouth soreness or ulcers can affect eating and drinking; report significant symptoms.

5 Skin


Pemetrexed

Rash can occur; prescribed steroid prophylaxis helps reduce this risk.

6 Infusion / allergy

Carboplatin

Rash, wheeze, swelling, dizziness or breathing difficulty during infusion needs immediate assessment.



Selected effects only: the same symptom can have more than one cause, and not every possible side effect is shown.

Pemetrexed + carboplatin

Lung · NCCP 00318 · version 6

TREATMENT JOURNEY

Day 1 Pemetrexed 500 mg/m ² IV followed by carboplatin AUC 5 IV.	Every 21 days Treatment repeats every 21 days if review and blood tests allow.	Mesothelioma Usually up to 4-6 cycles unless progression or unacceptable toxicity occurs.	NSCLC Usually 4-6 cycles, then maintenance pemetrexed (NCCP 00222) where appropriate.
---	--	---	---

IMPORTANT MEDICINE-SPECIFIC EFFECTS

Pemetrexed • Low blood counts can increase infection, anaemia or bruising risk. • Mouth/gut inflammation and skin rash can occur. • Kidney function can deteriorate and is monitored.	Carboplatin • Low blood counts, especially neutrophils and platelets, can occur. • Allergic infusion reactions can occur, particularly after repeated exposure. • Nausea and vomiting may occur despite preventative medicines.
--	--

OTHER / SHARED TREATMENT EFFECTS

The anatomy page highlights important symptom patterns for this exact regimen, but it cannot show every possible effect. Fatigue, nausea, appetite change, mouth/bowel symptoms or blood-count changes may occur depending on the medicines used. Report new or worsening symptoms even if they are not listed.

Monitoring / medicines • Folic acid and vitamin B12 support are required with pemetrexed. • A corticosteroid is used around pemetrexed to reduce skin reactions. • Blood count, kidney and liver profile are checked before each cycle.	Bring / track • Carboplatin dosing depends on renal function. • Follow the regimen-specific antiemetic prescription.
--	---

WHEN TO CALL - DO NOT WAIT FOR THE NEXT APPOINTMENT

Contact oncology urgently for fever or sudden illness, significant bleeding, new breathing difficulty or infusion reaction, uncontrolled vomiting/diarrhoea, severe mouth soreness, marked reduction in urine or rapidly worsening symptoms. For a medical emergency call 112/999.

Consent discussion support

Use the approved consent process to document treatment aim, expected benefit, material risks, alternatives, questions and the patient's decision.

Scan to open the regimen-specific patient portal

<https://sactcheck.com/patient/00318/>

Official NCCP 00318 v6

SACTCheck v0.81.1 · patient information prototype for clinical review. Follow your oncology team and local emergency instructions.

